



# HIERACIUM HOVER FLIES

*Cheilosia* spp.



## History in New Zealand

---

The hieracium root-feeding hover fly (*Cheilosia urbana*) and the crown hover fly (*Cheilosia psilophthalma*) are native to Europe and were imported from Switzerland by Manaaki Whenua - Landcare Research, in conjunction with the Hieracium Control Trust, in the late 1990s for testing.

Permission to release the hover flies was granted in 2001 and a further shipment imported to begin a rearing colony. A limited number of releases have been made to date, owing to rearing difficulties, and establishment has not yet been confirmed.

## How would I find/recognise it and what is its lifecycle?

---

These hover flies look similar to other species of hover fly that are already present in New Zealand. You would not be able to tell the two hieracium hover flies apart without a microscope. Both are medium-sized (7-8 mm long) stout-bodied blackish flies with large heads. In males, the compound eyes meet on the top of the head, whereas in females, there is a small gap between the eyes.



Root-feeding hover fly

Females of both species lay their eggs in the leaf axils of hawkweed plants in spring. Upon hatching, the larvae of the root-feeding hover fly move underground, so you are unlikely to see them. However, you may see the creamy coloured larvae of the crown hover fly as they feed on plant material above the ground - they often burrow into the centre of rosette plants. The larvae of both species are quite mobile and can easily move from plant to plant. Both species overwinter as brown pupae - the crown hover flies pupate on the soil surface whereas root-feeding hover flies pupate just below the soil surface. New adults of both species emerge in spring and both have only one generation per year.

## How does it damage hawkweeds?

---

Larvae of the root-feeding hover fly feed underground externally on the roots of the plant. Larvae of the crown hover fly feed on the centre of rosettes, at the base of the stolons, in the leaves and leaf axils, and in the stolon tips. Feeding by these hover flies stunts the growth of affected plants.

## Will it attack other plants?

---

No, it is extremely unlikely that hieracium hover flies will attack plants other than mouse-ear hawkweed (*Pilosella officinarum*), king devil hawkweed (*P. praealta*), field hawkweed (*P. caespitosa*), tussock hawkweed (*H. lepidulum*), and orange hawkweed (*P. aurantiaca*).

**How effective is it?**

---

NA.

**How can I get the most out of it?**

---

NA.

**How do I select a release site?**

---

Read *Guidelines for selecting release sites for biocontrol agents*.

**How do I collect it for release at other sites?**

---

NA.

**How do I manage the release sites?**

---

NA.

**For further information contact:**

Paul Peterson  
Manaaki Whenua - Landcare Research  
Private Bag 11052  
Manawatu Mail Centre  
Palmerston North 4442  
NEW ZEALAND  
Email: [petersonp@landcareresearch.co.nz](mailto:petersonp@landcareresearch.co.nz)  
Ph (06) 353 4871