

FIELD HORSETAIL WEEVIL

Grypus equiseti

The history of field horsetail weevils in New Zealand

The field horsetail weevil is native to Eurasia and North America. They were first imported into containment in New Zealand for study in 2013 by Landcare Research on behalf of the Lower Rangitikei Horsetail Control Group. Permission to release the weevils was granted by the EPA in mid-2016, and the first field release is expected to be made in late 2016. The weevil has not been used as a biocontrol agent anywhere in the world before, and is likely to be the only agent needed to provide adequate control for field horsetail, but if not there are other agents that can be considered.

How would I find field horsetail weevils?

The adults are 6-8mm long with mostly black wing cases, but creamy at the tips, and black and brown speckled undersides. Females are generally a bit larger than the males. They readily drop from the plant when disturbed which will make them difficult to see in the field. The adults are likely to be relatively long-lived as they have been observed to live for at least several months in the lab.

You are also unlikely to see the eggs despite them being orange and relatively large (0.5mm), as the female deposits 1-6 of them into a feeding cavity within the horsetail stem and then often seals the opening.



Larva



Adult weevil feeding

The larvae are grub-like and creamy-yellow in colour with tan head capsules. They hatch within the horsetail stems in late summer and then burrow, or mine, down into the below ground rhizomes. The larvae overwinter in the rhizomes and then pupate, emerging as new adults the following summer. Only one generation is completed each year.

Since all stages of the insect will be difficult to see in the field the best thing to look for will be their damage in spring and summer (see photo). There is no other insect on field horsetail that you are likely to confuse with this weevil. However, natural dieback that occurs in autumn/winter could be confused with weevil damage so look instead in spring/summer.

How do field horsetail weevils damage field horsetail?

Both the adults and the larvae feed on horsetail. The adults eat their way into horsetail stems, hollowing out large cavities, and effectively killing the stem above the feeding area. Larvae may kill the entire stem as they burrow down into the roots, where they continue to feed and also damage the rhizomes.

Will field horsetail weevils attack other plants?

In its native range the horsetail weevil attacks a range of horsetail species and host-range testing



OTHER AGENTS



has indicated that the other horsetail species also present in New Zealand, *Equisetum hymale*, is also a potential host. However, no other plants are expected to be at risk.

How effective are field horsetail weevils?

It is too soon to know what impact the horsetail weevil will have in New Zealand. However, in the lab potted field horsetail plants are severely damaged by the weevil and it is expected to be capable of having a substantial impact.

How can I get the most out of field horsetail weevils?

If the horsetail weevil establishes at initial release sites it would be worth helping to establish them in all areas where they are needed. The weevil is likely to disperse fairly slowly, possibly only a few kilometres per year.



Adult feeding damage

How do I choose a release site?

Read *Guidelines for selecting release sites for biocontrol agents*.

How do I collect field horsetail weevils?

When adults are present in good numbers either collect them by sweeping infested horsetail with a butterfly net or by sucking them up with a garden leaf vacuum machine. Modify the tube of the garden-leaf vacuum so that the weevils are collected and not sucked through. Take a sleeve of coarsely-woven material, with one end sewn or tied shut and the other end open (old socks or pantyhose could also be used), and fit it securely around the end so that it forms a bag in the mouth of the tube. Use a pooter to separate the weevils from other material collected during the vacuuming process, which many include pasture pests.

Place collected adults into a chillybin to transport to a new site. Shift at least 100 individuals to each new site at any time during mid to late summer. This will allow time for development of new larvae before horsetail dies back for winter. Sheltered release sites are recommended.

How do I manage the release sites?

Avoid any activities that will interfere with the weevils, such as herbicide application. If you need to undertake control measures then avoid the release site.

For further information contact:

Lindsay Smith
Landcare Research
PO Box 69040
Lincoln 7640
NEW ZEALAND
Email: smithl@landcareresearch.co.nz
Ph (03) 321 9805
Fax (03) 321 9998