

# Fauna of New Zealand

## **Editorial Advisory Group**

(appointments made on a rotational basis)

**MEMBERS AT ENTOMOLOGY DIVISION**  
Department of Scientific and Industrial Research  
Mount Albert Research Centre  
Private Bag, Auckland, New Zealand

*Ex officio*  
Director - Mr J. F. Longworth  
Leader, Systematics Group - Dr B. A. Holloway

*Co-opted from within Systematics Group*  
Dr T. K. Crosby, Dr G. W. Ramsay

---

### **UNIVERSITIES REPRESENTATIVE**

Dr R. M. Emberson  
Entomology Department, Lincoln College  
Canterbury, New Zealand

### **MUSEUMS REPRESENTATIVE**

Dr J. C. Yaldwyn  
Director, National Museum of New Zealand  
Private Bag, Wellington, New Zealand

### **OVERSEAS REPRESENTATIVE**

Dr J. F. Lawrence  
CSIRO Division of Entomology  
P.O. Box 1700, Canberra City, ACT 2601, Australia

---

—◎—

## **Series Editor**

Mr C. T. Duval  
Systematics Group, Entomology Division  
Department of Scientific and Industrial Research  
Mount Albert Research Centre  
Private Bag, Auckland, New Zealand

DIC XXV DIC XXV

XXV DIC XXV

XXV DIC XXV

XXV DIC XXV

# Fauna of New Zealand

Number 14

## Lepidoptera

– annotated catalogue, and  
keys to family-group taxa

J. S. Dugdale

Entomology Division

Department of Scientific and Industrial Research  
Private Bag, Auckland, New Zealand

• SYSTEMATICS GROUP 1963–1988 •

TWENTY-FIVE YEARS' SERVICE  
TO NEW ZEALAND ENTOMOLOGY

### Cataloguing-in-publication citation

DUGDALE, J. S.

Lepidoptera – annotated catalogue, and keys to family-group taxa /  
J. S. Dugdale. – Wellington : DSIR Science Information Publishing  
Centre, 1988.

(Fauna of New Zealand, ISSN 0111-5383; no. 14)  
ISBN 0-477-02518-8

I. Title            II. Series

UDC 595.78(931)

*Date of publication: see back cover of subsequent numbers*

### Suggested form of citation

Dugdale, J. S. 1988: Lepidoptera – annotated catalogue, and keys to  
family-group taxa. *Fauna of New Zealand [number]* 14.

—⊗—

This publication was produced by offset lithography. The author's MS.  
was transcribed on to a text processor, and after revision, editing, and  
style coding was phototypeset at the N.Z. Government Printing Office.  
Times New Roman type is used for most of the text; major headings  
and figure labels are set in Univers; Garamond and Geneva are used  
in the titles.

The Editorial Advisory Group and the Series Editor acknowledge  
the following co-operation.

*DSIR Research Orchard, Havelock North:*

Mrs T. Crockford – assistance with text processing

*Mount Albert Research Centre, DSIR:*

Mr D.W. Helmore – habitus drawings and front cover motif

Mrs L.S. Howey – transcription of manuscript text

Mrs M.L. Lessiter – photoreduction of line figures

Mrs H.A. Whelan – computer file management

*Science Information Publishing Centre, DSIR:*

Dr N. Hawcroft – supervision of production and distribution

Mr C. Matthews – assistance with production and publicity

—⊗—

**Front cover:** The insects depicted are, from the top down, representative  
of Geometridae, Oecophoridae, Pyralidae, and Tortricidae. These, with  
Noctuidae and Tineidae, are by far the most species-rich families in  
New Zealand's Lepidoptera fauna. *Artist: D.W. Helmore.*

© Crown Copyright

Published by Science Information Publishing Centre, DSIR  
P.O. Box 9741, Wellington, New Zealand

## FOREWORD

In New Zealand, terrestrial invertebrate systematics began in a concerted way when a group dedicated to systematics was formed in Entomology Division, DSIR, in the early 1960s. In 1988 we mark the twenty-fifth anniversary of the establishment of that group, and it is timely to reflect on past events and achievements.

In the early years the group was based in Nelson, on the South Island. It moved in 1973 to the Mt Albert Research Centre, in Auckland, where the New Zealand Arthropod Collection grew and developed and, in 1982, the 'Fauna of New Zealand' was begun.

Most of the group's early members are still associated with it – three as full-time scientists, and three as research associates – and this has ensured a continuity of expertise and of esprit that has been of tremendous benefit. We are confident that the Systematics Group and the 'Fauna' will continue to have a central role in entomology and invertebrate systematics for the next twenty-five years.

The group has always sought to develop strong links with other institutions and individuals working in this field, in New Zealand and overseas. In particular the 'Fauna' has provided a vehicle for publication of definitive taxonomic studies on New Zealand insects and other terrestrial invertebrates, by specialists in New Zealand and by colleagues throughout the world. Perhaps the closest of the special relationships that have been developed over the years is that with the Entomology Department of the British Museum (Natural History). It is appropriate that Systematics Group's jubilee year should be marked with the publication of two commemorative 'Fauna' contributions that reflect this particular relationship: John Noyes's Encyrtidae and John Dugdale's Lepidoptera catalogue. The groundwork for each volume was laid during reciprocal study visits, Noyes to NZAC and Dugdale to BMNH.

It is five years since the 'Fauna' series began. Thirteen volumes comprising 1800 pages are now in print, and we are assured of continued support from other contributors and from subscribers. We are confident therefore that the 'Fauna' was well conceived, is making a significant contribution to biology, and has a very definite future. Indeed Entomology Division is firmly committed to the objective of providing authoritative and comprehensive guides to identification of insects and other terrestrial invertebrates through the medium of the 'Fauna' series.

It is a pleasure to acknowledge the achievements of Systematics Group in its first twenty-five years, in particular the establishment of the 'Fauna' series, and I wish both the group and the 'Fauna' well for the future.

*J.F. Longworth*

Director  
Entomology Division  
DSIR

Dedicated to the memory of three amateur lepidopterists:

**George Vernon Hudson**

1867–1946

whose life work this publication  
attempts to keep evergreen

**Alfred Philpott**

1871–1931

whose pioneering studies in Lepidoptera morphology  
are now assuming their true significance

**Kenneth John Fox, FRCOG**

1936–1986

who died before this catalogue reached full term  
but whose enthusiasm and insistence  
ensured its completion



*“... I would advise you to get a knowledge of facts from actual observation. Facts looked at directly are vital; when they pass into words half the sap is taken out of them.”*

—John Tyndall, ‘Fragments of Science’ (1876, p. 287)



## ABSTRACT

This annotated synonymic catalogue represents an attempt to verify and define nomenclaturally the species of Lepidoptera recorded from New Zealand since 1769. It is based on a thorough re-examination of type material and recorded data. Types of nearly all nominal species held by overseas institutions were examined during 1980–81. Of the 2150 published names, type specimens of about 1570 (73%) are in the Northern Hemisphere; over 1460 of these (68%) are at the British Museum (Natural History). The 1761 recognised species are assigned to modern family-group concepts, and are listed under superfamilies, which are placed in 'systematic' sequence. Within each superfamily, all subordinate taxa are listed alphabetically. Synonyms are listed in date order. For each species and synonym, basic nomenclatural data concerning the type locality, collector, type status, condition of specimens (if noteworthy), and repository are given. Where relevant, species are cross-referenced to G.V. Hudson's illustrated monographic works. References are also given, as appropriate, to genitalia descriptions by A. Philpott and to first records of adventive species. Any additional remarks are given under 'Notes' for a taxon entry. Names are proposed for three misidentified species, i.e., for which invalid concepts of previously described species have entered current usage: *Dumbletonius sylvicola* for *Trioxycanus enysii* (not of Butler); *Heterocrossa rubophaga* for *H. adreptella* (not of Walker); and *Stathmopoda horticola* for *S. skelloni* (not of Butler). Brief introductory sections outline the history of study, classification, and composition of New Zealand's lepidopteran fauna. Keys to superfamilies, families, and (where possible) subfamilies are presented, with illustrations of diagnostic features. There is also a key to the brachypterous Lepidoptera. These keys conform to the New Zealand situation; any wider relevance is coincidental. The 69 species reported from the Kermadec Islands are excluded from the main catalogue, but are listed in an appendix. There are over 700 references.

## CONTENTS

	Page
Introduction .....	8
Historical notes .....	9
Classification .....	12
Relationships of Lepidoptera .....	12
Higher classification within Lepidoptera ...	12
Superfamilies of Lepidoptera, and their families .....	13
Composition of Lepidoptera fauna of N.Z. ....	15
Degree of endemism .....	15
Proportion of non-ditrysian groups .....	15
Ditrysia representation .....	15
'Microlepidoptera / macrolepidoptera' proportions .....	15
Fossil representation .....	16
Relationships with other biogeographical areas .....	16
Concepts and conventions used in catalogue .	17
Keys to higher taxa in N.Z. Lepidoptera	
(A) Key to superfamilies .....	19
(B) Keys to families and subfamilies .....	22
(C) Key to brachypterous Lepidoptera .....	27
Illustrations (to the keys) .....	28

	Page
Taxonomic catalogue .....	52
References .....	215
Appendix: Lepidoptera recorded from the Kermadec Islands .....	235
Taxonomic index .....	237

## ACKNOWLEDGMENTS

A work of this nature cannot be done in isolation, and I am deeply indebted to the following – whom it is my great pleasure to thank – for assistance with specimens, nomenclatural problems, access to types, collections, and/or published or unpublished records and information, hospitality in several countries, expeditiously reviewing sections of the manuscript, and in general by their enthusiasm and generosity of spirit prodding a sometimes lethargic and distracted behemoth towards the finishing line. New Zealand: R.C. Craw, T.K. Crosby, the late Kenneth J. Fox, G.W. Gibbs and family, C.J. and O. Green, A.C. Harris, N. Hudson, P.M. Johns, G. Kuschel, P.A. Maddison, C. Muir, R.G. Ordish, R. Palma, B.H. and C. Patrick, A. Savill, and K.A.J. Wise.

Australia: I.F.B. and J. Common, M. Horak, P.B. McQuillan, E.D. Edwards, and E.S. Nielsen.

Canada: D.E. Gaskin.

Denmark: N.P. Kristensen and family.

France: C. Dufay, J. Minet, and P. Viette.

The Netherlands: A. Diakonoff, and C. Wilkinson and family.

United Kingdom: P. Ackery, J.D. Bradley, D.S. Fletcher, the late Allan Hayes, J.D. Holloway, L.A. Mound, I.W.B. Nye, G.S. Robinson, K.R. and E. Sattler, M.J. Scoble, M. Shaffer, K. Tuck, A.K. Walker, A. Watson, and P.S. Whalley.

U.S.A.: D.R. Davis, J.B. Heppner, and R.W. Hodges.

For proofreading I am very grateful to Jo Berry and Grace Hall; and for the habitus illustrations I am especially grateful to Des Helmore.

For continued support I am grateful to my Director, J.F. Longworth, and to DSIR Head Office for a study award in 1980–81, enabling me to visit the British Museum (Natural History). I am especially grateful to the present Keeper of Entomology, Dr L.A. Mound, for the copious facilities he so freely placed at my disposal.

To all the above, and to my long-suffering family, my sincere thanks.

## INTRODUCTION

With over 1760 species the order Lepidoptera is the third largest in New Zealand, after Coleoptera (*c.* 5000) and Diptera (2000+). Lepidoptera occupy all biotopes except caves in New Zealand; there are species living in the rocky coast spray zone, and even in the nival zone – an undescribed geometrid is known from the Cheval Ridge, Malte Brun Range, at 3000 m, 700 m altitude above the summer snowline. Some are ubiquitous (e.g., *Orocrambus flexuosellus* Doubleday), and others are very restricted (e.g., "*Tortrix antichroa* Meyrick, only on Mt Taranaki). Some are so rarely encountered as to raise the question of extinction (e.g., the large *Titanomis sisyrota* Meyrick). Some have become extinct locally – e.g., *Aoraia dinodes* (Meyrick) at Invercargill, and *Hydriomena arida* (Butler) at Dunedin (B. Patrick, pers. comm.).

Lepidoptera in New Zealand commanded the attention of early naturalists, whether Maori or European. There are specific Maori terms for several species; and European culture has imposed yet another, more comprehensive system.

The New Zealand Lepidoptera fauna stands out on three counts: (a) the high level of endemism; (b) the distinctive nature of many forms in relation to those in Australia; (c) the strange absences. It is a workable-sized fauna, even if some elements are

enigmatic, and this publication, I hope, will give essential nomenclatural information about this important and in many ways elegant group, and summarise its classification.

Since G.V. Hudson published his monumental "Butterflies and Moths of New Zealand" (1928) and its "Supplement" (1939) there has been no comprehensive catalogue of New Zealand Lepidoptera. At all taxonomic levels, Lepidoptera classification has changed greatly in the intervening years, and requirements for distinguishing and describing new species and higher taxa have become more scientifically rigorous and time-consuming. After 1950 (the year Dr J.T. Salmon published Hudson's posthumous "Fragments of Entomology") new species have been recognised, classification of the Lepidoptera as a whole has been radically altered, and additional exotic species have either established themselves or are now known to arrive in New Zealand with great regularity.

The greatest stumbling block in the way of taxonomic work by New Zealanders on New Zealand Lepidoptera is the fact that, for the 2150 names proposed for species found in New Zealand, 1569 or 73% of the type specimens are in Northern Hemisphere institutions. The British Museum (Natural History) alone houses some 1460, or 68%, which were not readily accessible to workers in New Zealand. Earlier New Zealand workers had to take Meyrick's and Hudson's concepts on trust; or, after Hudson's death, have the types examined for them (e.g., Dumbleton 1966, Gaskin 1971, 1975), an arrangement conducive to error and confusion (Dugdale 1986).

This catalogue is an attempt at solving three problems: (1) how to relate the classification changes to Hudson's illustrated monographs; (2) how to present the relevant information concerning each type specimen (the specimen(s) on which the author of a name based the original description of that species); and (3) how to identify a specimen to one of the higher taxonomic categories in the catalogue, so that the identifier has a fighting chance of recognising it in 'Hudson'.

The catalogue therefore has four functions:

- to place New Zealand's Lepidoptera in currently accepted classifications, and to cross-reference the species to Hudson's illustrated monographs, i.e., his works of 1898, 1928, 1939, and 1950;
- to provide basic nomenclatural information for the species and genera reported or known to be in New Zealand;\*

\*The Kermadec Islands are excluded from the scope of the catalogue, and hence the index, but a checklist of Kermadec species is given as an appendix.

- to provide keys to higher taxa (superfamilies, families, and where possible subfamilies) known in New Zealand; and
- to provide some historical notes on the study of Lepidoptera in New Zealand, and give some idea of the group's representation here, in contrast to other countries or regions.

It is not intended to give a general account of Lepidoptera; for that, the reader is directed to Common (1970, 1975) and Nielsen & Common (1988, in preparation).

This catalogue and set of keys is dedicated primarily to G.V. Hudson, as a complement – and, I intend, a compliment – to his pioneering works, but not a replacement of them. As the likelihood of a replacement for 'Hudson' is remote, this catalogue is offered as a way of keeping his works up to date. It is my wish also to acknowledge the debt that, as a working lepidopterist, I owe to Alfred Philpott and to my late co-explorer, Kenneth Fox.

In all, some 2280 names have been applied to moths and butterflies collected in New Zealand. Most of the type specimens have been seen by me. Those held overseas were examined during a visit to the British Museum (Natural History), London, in 1980–1981, with the support of a study award. In the catalogue I recognise over 1760 species as 'valid', but for some groups with hosts in Asteraceae and Scrophulariaceae there is evidence for the presence of cryptic species. For all groups new methods of looking at species will uncover other cryptic species assemblages, as pheromonal and electrophoretic methods already have in Tortricidae (Foster *et al.* 1986) and Hepialidae.

## HISTORICAL NOTES

The first New Zealand Lepidoptera brought to Europe were two nymphalid butterflies collected by Banks and his assistants during Cook's first voyage, and were described by Fabricius in 1775. Their capture did not excite Banks (Gibbs 1980b, p. 110). A copper butterfly obtained by Drury – probably from a member of one of Cook's later voyages (Gibbs 1980b, p. 141) – and also described by Fabricius was the third. Drury thought it was from India. All three were depicted precisely by Jones in his "Icones; Papilioes Nymphales ..." of 1785; only the nymphalid specimens, the yellow and red admiral respectively, are known to survive. The copper butterfly may be in the MacLeay Museum, Sydney (Andrews 1986, p. 41).

Entomological collections made during the voyage of Dumont D'Urville were described by Boisduval (1832). One New Zealand moth – the magpie moth – was described (as from Papua New Guinea) and depicted.

None of the voyages yielded nocturnal species, either of Lepidoptera or of Coleoptera (e.g., *Prionoplus*). Can we presume that the collectors spent each night aboard, and that the ship's lights were not bright enough to attract moths or large beetles from the shore?

With land-based exploration and the beginnings of settlement came an increase in specimens. Yet by 1844 only four Lepidoptera collections had reached Britain: a puriri moth collected by J.G. Children (stated to be from "Van Diemens Land"); two small collections presented by Captain W. Parry and by J. Clarke Ross of the Antarctic Expedition, described by Edward Doubleday; and specimens collected by the Rev. Richard Taylor.

The earliest account of New Zealand Lepidoptera natural history is that of Taylor (1843). He included much information provided by his Maori companions, and recorded the presence (and Maori names) of adults of the then hostless monarch butterfly and moon moth. The respective hosts, asceliapiads and wattles, were introduced by Europeans.

The period 1845–1880 saw increasing collecting by resident natural historians such as Percy Earl (Wellington and coastal Otago), Lt Col. D. Bolton, Dr Andrew Sinclair (Auckland), R.W. Fereday, J.D. Enys (Canterbury, Hawkes Bay), Thomas Oxley (Nelson), Dr J. Hector and, later, Prof. F.W. Hutton (Otago), William Colenso (Hawkes Bay – Taupo), and the Rev. J.F. Churton (Wellington briefly, then Auckland). There were also visitors who collected, such as Dr F. Knaggs. Contact with London was strong; Oxley, Fereday, Enys, and Hector all knew F. Walker or A.G. Butler at the British Museum in Russell Square. The Imperial Austrian frigate 'Novara' visited New Zealand on expedition, but relied largely on local collectors (Sinclair in Auckland, Oxley in Nelson), although the geologist Ferdinand Hochstetter also provided some entomological specimens.

Most specimens from New Zealand were presented or sold to the British Museum (as befitting Britons in a British colony). The earlier material was described by Francis Walker and the later by A.G. Butler. Some of Fereday's early material was described by Achille Guenée in Paris, where some types remain; others are now in BMNH. The most extensive collections were those of Lt-Col. D. Bolton around Auckland and T.R. Oxley around Nelson, providing the bulk of the species described from New Zealand by Walker.

Lt-Colonel D. Bolton commanded the Royal Engineers in Auckland between March 1850 and November 1853. He had previously toured New Zealand with Governor Grey in 1847, and was then stationed in Wellington during 1848–1850. As his collection does not include *Wiseana cervinata*

Walker, and many of his specimens (Lepidoptera and cicadas) show characters distinctive of Auckland populations, it is likely that Bolton made his collection during his term in Auckland. He had two houses, an official one at Onehunga and another in Emily Place, at the foot of Symonds Street. He was briefly on the executive council of New Ulster (one of three provinces of New Zealand, a system abandoned in 1852). One species he collected – *Stigmella maoriella* (Walker) – has not been found since. Bolton embarked for England via Sydney in November 1853, and his collection reached the British Museum early in 1854, hence the museum accession number “54.4”. He later died in South Africa (Auckland Institute and Museum Library records; Sir Charles Fleming, pers. comm.).

T.R. Oxley, a professional photographer from London, was our first extensive resident collector. He had previously collected in Victoria (Australia), returning in 1855 to London, where he was befriended by Henry Stainton, the leading British microlepidopterist. Oxley then went to Nelson and made three collections: one he despatched to Walker (mislabelled at BMNH as from Auckland), one to Stainton, and – by arrangement – one to Dr Felder, as part of the ‘Novara’ Expedition results. Oxley’s specimens to Felder are illustrated in the ‘Lepidoptera’ section of the ‘Novara’ results (Felder & Rogenhofer 1875). Oxley later became, in turn, a brewer and an Inspector of Nuisances; he died at Nelson in 1887 (M. Watson, pers. comm.).

Late in 1879 Edward Meyrick came to New Zealand as a classics master at Cathedral Grammar School, Christchurch. Over the next seven years he divided his time between New Zealand and Australia, where he taught at the King’s School, Parramatta, Sydney, building up a large collection – mainly of “Microlepidoptera” – from each country. Home leave on two occasions during this period allowed him to check his captures with the specimens described by Walker and Butler. For each of those species that his friend Fereday had got Guenée to describe, Meyrick was able to examine “an exactly similar one” (Meyrick 1884b, p. 50).

Meyrick travelled extensively, and recorded his daily captures in a diary (now in the BMNH, microlepidoptera floor). He was in contact particularly with Fereday, Enys, and Hutton, and towards the end of his stay with G.V. Hudson. By 1887–1890 Meyrick had revised virtually all families in New Zealand, except for Rhopalocera. Also by this time he was publishing extensively on microlepidoptera from Australia and the central Pacific. The fauna of the latter area – to which he was introduced by G.F. Mathew, a naval officer based at Sydney and in Fiji – particularly excited him, and in subsequent papers he remarked on the disjunct

relationships between New Zealand and some Pacific islands, a topic of great interest today.

Meyrick gathered together the scattered descriptions of previous workers, and put the classification of Lepidoptera in New Zealand on a relatively sound footing. Only Felder & Rogenhofer’s and two of Butler’s contributions were illustrated; Meyrick did not illustrate any of his New Zealand works, but the meeting with Hudson over a weekend in January 1886 in Wellington laid the foundations for Hudson’s massive contribution. New Zealand and, to a certain extent, Australia were the only places for which Meyrick maintained an interest in macrolepidoptera.

After Meyrick’s departure for England, G.V. Hudson, A. Philpott, G. Howes, C.C. Fenwick, and later C.E. Clarke and M.O. Pasco amassed large collections, and these have stayed in New Zealand. Also there was extensive entomological exploration, from the subantarctic islands (1909) to the Kermadec Islands (1908), and general overviews of the fauna were produced. All this was done by resident amateurs. During the late 1920s Alfred Philpott became – at his own request – the honorary lepidopterist at the Cawthron Institute, where previously he had drawn a salary (Tillyard 1931, p. 173).

The first overview of our Lepidoptera, a synoptic catalogue, was compiled by Fereday (1898). It is marred by typographical errors, but formed the basis (unacknowledged) for F.W. Hutton’s “Index Faunae Novae-Zealandiae” in 1904. Fereday’s list was based on Meyrick’s advice. Fereday’s catalogue was largely overshadowed by G.V. Hudson’s first monograph, on the macrolepidoptera (Hudson 1898). It too was based on Meyrick’s advice and guidance.

By 1900 Hudson was in regular correspondence with Meyrick, sending numbered specimens and a ledger sheet; Meyrick kept the specimens and returned the ledger sheets with names beside the relevant numbers. Hudson also guided Meyrick’s papers through the New Zealand Institute’s *Transactions*. The other lepidopterists resident in New Zealand also sent material to Meyrick for identification or description. Only S. Lindsay (at the Canterbury Museum) had the types of new species based on that material returned, labelled as such.

Over the period 1900–1938, Meyrick ruled the roost. His view of what was or was not a species was promulgated by Hudson in his 1928 and 1939 books, and was largely accepted. Meyrick had seen the Doubleday, Walker, Butler, and now the Felder types in the BMNH (and may have had scant regard for types anyway, as suggested by Clarke 1956, p. 9). We had no option but to interpret the New Zealand Lepidoptera by Meyrick’s and Hudson’s

concepts, which in New Zealand were based on Hudson's collection.

Nevertheless, there was some independent thought. G. Howes in Dunedin described several noctuid and geometrid species, and L.B. Prout and G.B. Longstaff (at BMNH) were revising Meyrickian concepts in macrolepidoptera.

The period 1920–1930 is characterised by the significant pioneering efforts of Alfred Philpott, who studied the structure of many moth families. He examined mouthparts, thoracic and leg structures, and genitalia in detail. His work on Hepialidae, Mnesarchaeidae, and Micropterigidae was detailed to a degree well beyond contemporary work in Europe and North America. Philpott also pioneered the representative world collection while he was with the Cawthron Institute, exchanging with A.J. Turner in Australia, E. Hering in France, F. Schaus in Costa Rica, and A.S. Packard, A. Busck, and J.R. Eyer in the United States. One result is the extensive exotic collections now in NZAC. Another notable pioneer was M.N. Watt, in Wanganui, whose illustrations and descriptions of Nepticulidae and Gracillariidae (in the 1920s) are equal to present-day standards. He continued to study leaf-miners when he moved to Dunedin, and his extensive collections are now at NMNZ.

By the time of Hudson's death, in 1946, there were three extensive collections of New Zealand Lepidoptera in the country: Hudson's (at the then Dominion Museum, Wellington), Philpott's (at the joint Cawthron Institute / Entomology Division, DSIR Entomological Research Station at Nelson), and Clarke's (at the Auckland Museum). There were also four local collections – M.O. Pasco's from Queenstown (at Invercargill), C.C. Fenwick's at the Dominion Museum, G. Howes's (split between the Auckland Museum and Dominion Museum), and S. Lindsay's (at the Canterbury Museum). Luckily, most of Fereday's collection was also preserved in the Canterbury Museum. All seven collections include types.

The period 1950 to the present day is still characterised by a high degree of amateur participation. J.T. Salmon, at first the lone professional, set out after Hudson's death to augment the Dominion Museum's holdings separate from the Hudson Collection, and described the exciting Three Kings Islands fauna, but after 1956 his interests turned largely to Collembola. Amateurs and others started to concentrate on local and national collecting, and comprehensive collections were amassed by the Forest Biology Survey of the Forest Research Institute at Rotorua, T.H. Davies (Hawkes Bay), K.J. Fox (Taranaki), C.J. Green (Auckland area), N. Hudson (South Auckland – Bay of Plenty), C. Muir (Riccarton Bush, Christchurch), and B. Patrick

(Otago – Southland). Some have concentrated on a group, e.g., Noctuidae (Fox), Geometridae, Psychidae, Tortricidae (Patrick), or on an ecological entity, e.g., migrant Lepidoptera (Fox).

Professional lepidopterists have dealt either with taxonomic problems (K.A.J. Wise) or with groups of systematic or economic importance (L.J. Dumbleton – Agathiphagidae, Hepialidae; J.S. Dugdale – Tortricidae, forest Geometridae; D.E. Gaskin – Crambidae; G.W. Gibbs – Micropterigidae, Mnesarchaeidae, Rhopalocera). Island faunas have been investigated by Salmon and Bradley, and later by Dugdale (subantarctic islands) and Wise (White Island, and others). The influence of Philpott is reflected in the work of Gibbs, Dumbleton, and Dugdale on the Micropterigidae, Mnesarchaeidae, Hepialidae, Tortricidae, and female genital systems.

Lately the Nepticulidae have been worked on by a group at the Vrije Universiteit, Amsterdam, with local help, as a contribution to the *Fauna of New Zealand* series (Donner & Wilkinson, in press). This may be a forerunner of other such 'off-shore' revisions, based on the extensive collections at NZAC, AMNZ, NMNZ, CMNZ, and FRNZ; and the large, privately held collections of B. Patrick (Dunedin) and the late K.J. Fox (Manai, Taranaki).

The above account briefly acknowledges the better known names. There were many others: Ambrose Quail, who first worked on the life history of the puriri moth and variation in the tortricid *Epalxiphora*; A.V. Chappell, who described eggs and life histories of several species; A. Purdie, who started a study of host-plant moth associations now in urgent need of revival; W.L. Buller, who described a now unrecognised hepialid; and many collectors such as R.M. Sunley, who assisted Hudson; W. Smith, who accompanied Howes to Fiordland; the painfully shy W. Heighway, who assisted Philpott and Lindsay; A. Hamilton, who sent his father "novelties" from beyond Lake Wakatipu; F.S. Oliver and Averil Lysaght, whose collections have largely disappeared.

All have contributed, and, creditably, nearly all the early material is still preserved. That which has been unwittingly destroyed or allowed to disintegrate comes from areas where re-collection is possible. The fine, careful, continuous collecting at a local level now being practised is showing up the presence of many more species than were hitherto expected. Such collecting, often involving elucidation of host-plant associations and periodicity, is a field very much open to amateurs. Where such work can be tied in with electrophoretic or pheromonal studies, then Lepidoptera systematics in New Zealand is even more exciting.

## CLASSIFICATION

### Relationships of Lepidoptera

The order Lepidoptera, with between 100,000 and 200,000 known species, is currently regarded as a member of the "Panorpoid complex" or Mecopteroidea, which comprises the superorders Antliophora (Mecoptera + Siphonaptera + Diptera) and Amphiesmenoptera (Trichoptera + Lepidoptera) (see Kristensen 1984, p. 145).

Lepidoptera are distinguished from their sister-group Trichoptera by the following apomorphies or unique character states:

- vestiture of wings and body largely made up of broad, overlapping scales;
- head lacking a median ocellus;
- apical segment of labial palpus with a group of receptors in a depression (vom Rath's organ).

Some other structures, such as the fore-tibial epiphysis, are not always present; even vom Rath's organ may not be in a pit, as in Southern Hemisphere Roeslerstammiidae (I.F.B. Common, pers. comm.). There is, in fact, only a very small suite of character states that distinguish all Lepidoptera from all Trichoptera, and Kristensen (1984, p. 156) has pointed out the similarity in habitus between less advanced Trichoptera and the lepidopteran *Agathiphaga* Dumbleton.

### Higher classification within Lepidoptera

The higher classification within this order is currently undergoing extensive re-examination, following on from the last intensive summary (Common 1975). The situation is rendered difficult because of two factors:

- around 98% of known Lepidoptera belong to one division (Ditrysia), of relatively uniform structural organisation;
- the remaining 1–2% show a great variety of structural and genital organisation, with often profound differences between groups.

Work by Kristensen, Nielsen, Davis, Minet, and others referred to hereunder is gradually moving to a synthesis, but taxa above superfamily level are still interpreted differently by the various workers (cf. Nielsen 1982, 1985, Davis 1986, and Minet 1984; also the approaches by Heppner 1986, p. 20, and Scott 1986, pp. 30–38).

There is a consensus (Kristensen 1984, Nielsen 1985, Davis 1986) that Lepidoptera phylogeny is an example of Hennig's "additive typogenesis" (Nielsen 1985, p. 1 and fig. 1). This implies a gradual acquisition of group characters from Zeugloptera onwards, culminating in the endoporian ditrysian suite of characters. However, "this relatively simple picture has become modified and much more complicated by the considerable

amount of new information ... available during the last two decades" (Nielsen 1985, p. 2).

The following group-names have been used above the superfamily level (see Kristensen 1984, p. 186, fig. 13): Zeugloptera (includes Micropterigoidea); Aglossata (includes Agathiphagoidea); Heterobathmiina (includes Heterobathmioidea); Glossata, divided into Dacnonypha (haustellum without intrinsic musculature) and Myoglossata (with intrinsic musculature); the Myoglossata including the Neopseustina (Neopseustoidea); and the Neolepidoptera, including the Exoporia (Mnesarchaeoidea, Hepialoidea) and the Heteroneura (Nepticuloidea, Incurvarioidea, Palaephatoidea); the Eulepidoptera including the Etimonostryia (Tischerioidea); and Ditrysia (all other superfamilies). Note that Glossata, Myoglossata, Neolepidoptera, and Eulepidoptera (Minet 1984, p. 147) are clades. Opinion is divided (cf. Minet 1984, Nielsen 1982, 1985, and Davis 1986) over the monophyly or otherwise of such categories as Heteroneura and Neolepidoptera, and, for that matter, of the position of the taxa Heterobathmiina and Exoporia (left hanging by Minet 1984, fig. 26), and whether the Zeugloptera alone, or Zeugloptera + Aglossata are the sister-group of all other Lepidoptera (see discussion by Kristensen 1984, p. 166).

Comprehensive discussions of Heteroneura phylogeny are given by Common (1975), Davis (1986, pp. 55–61), and Nielsen (1982, 1985a, 1985b). By and large the argument involves two assumptions: (a) that changes in frenular structures arose once; and (b) that the internal change from a ventral to a dorsal common oviduct in relation to the copulatory chamber (Dugdale 1974) arose independently in Exoporia and Ditrysia (Common 1975).

The classificatory outline given below is largely derived from Nielsen (1985a, pp. 15–16; 1985b, p. 142), Minet (1986), and Nielsen & Common (in prep.), but no categories above superfamily are listed except where the higher category includes only one superfamily. In such instances the higher term is given only as a recognition term, of possible use to the reader in literature searches.

Within the division (Davis 1986) or phalanx (Minet 1984, 1986) Ditrysia some 28 superfamilies are recognised (Minet 1983; Nielsen & Common, in prep.). Brock (1971) noted two groupings of superfamilies; these were refined by Kyrki (1983) and Minet (1983), largely on the basis of the structure of the thoracic / abdominal junction in adult moths. The term Apoditrysia (Minet 1983, p. 201) includes all ditrysian superfamilies with a tortricoid basisternite, and excludes those with a tineoid basisternite, i.e., Tineoidea, Yponomeutoidea, and Gelechioidea. As a concept, it still requires testing over a greater range of species, and its relation to

metafurcasternal structures (Davis 1986) and prothoracic structures (Minet 1984) needs more precise evaluation.

Minet (1986) gives a revised outline of Lepidoptera classification, and reduces the number of ditrysian superfamilies to 26. His removal of Choreutidae from Sesioidea – as “*Ditrysia à affinités inconnue ou uncertaine*” (p. 292, and note 61, p. 306) – is of relevance to the New Zealand fauna. Minet’s outline does not include Hedyloidea, newly recognised as a group related to Papilionoidea (Scoble 1986).

### **Superfamilies of Lepidoptera, and their families**

Those superfamilies and families absent from New Zealand have a minus sign following them; where the sign is in parentheses, the family is represented by introduced species only; e.g., Cossidae (-).

**Micropterigoidea:** Micropterigidae [Zeugloptera]

**Agathiphagoidea:** Agathiphagidae – [Aglossata]

**Heterobathmioidea:** Heterobathmiidae –

**Eriocranoidea:** Eriocraniidae –, Lophocoronidae –, Acanthopteroctetidae – [Dacnonypha]

**Neopseustoidae:** Neopseustidae –

**Nepticuloidea:** Nepticulidae; Opostegidae –

**Mnesarchaeoidea:** Mnesarchaeidae [Exoporia]

**Hepialoidea:** Hepialidae in the strict sense; “primitive Hepialidae” –, Anomosetidae –, Neotheoridae –, Palaeosetidae –, Prototheoridae – [Exoporia]

**Incurvarioidea:** Prodoxidae [?]; Adelidae –, Cecidoseidae –, Crinopterigidae –, Incurvariidae –, Heliozelidae –

**Palaephatoidea:** Palaephatidae –

**Tischerioidea:** Tischeriidae –

[**Ditrysia:** “non-apodityrsian” superfamilies]

**Tineoidea:** Tineidae, Psychidae, Arrhenophanidae –, Eriocottidae –, Pseudarbelidae –, Roeslerstammiidae, Gassicillariidae, Bucculatrigidae –, Douglasiidae –

**Yponomeutoidea:** Yponomeutidae (including Ypsolophidae, Plutellidae in the sense of Nielsen & Common, in prep.), Glyphipterigidae, Lyonetiidae (including Bedelliinae), Heliodinidae –

**Gelechioidea:** Gelechiidae, Coleophoridae, Batrachedridae (in the sense of Nielsen & Common, in prep.), Cosmopterigidae, Blastobasidae (including Symmocidae, in the sense of Hodges 1978, p. 7) (-), Momphidae, Oecophoridae (Depressariinae, Oecophorinae, Stathmopodinae, Stenomatinae, Xyloryctinae, Peleopodinae in the sense of Hodges 1974 (-)), Lecithoceridae, Scythrididae, Holcopogonidae –, Elachistidae, Agonoxenidae –

[**Ditrysia:** Apodityrsia]

**Coccoidea:** Cossidae (-), Metarbelidae –, Dudgeoneidae –, Ratardidae –

**Castnioidae:** Castniidae – (Minet 1986 includes Castniidae in Sesioidea)

**Sesioidea:** Sesiidiae (-), Choreutidae, Brachodidae –

**Tortricoidea:** Tortricidae (Tortricinae, Chlidanotinae, Olethreutinae)

**Zygaenoidea:** Zygaenidae –, Heterognidae –, Phaudidae –, Megalopygidae –, Chrysopolomidae –, Somabrachyidae –, Limacodidae –, Dalceridae –, Epipyropidae –, Cyclotornidae –

**Immoidea:** Immidae –

**Copromorphoidea:** Copromorphidae, Carposinidae

**Epermenioidea:** Epermeniidae

**Schreckensteinioidea:** Schreckensteinidae –

**Alucitioidea:** Tineodidae –, Alucitidae –

**Pterophoroidea:** Pterophoridae

**Hyblaeoidea:** Hyblaeidae –

**Hedyloidea:** Hedyliidae –

**Hesperoidea:** Hesperiidae –

**Papilionoidea:** Nymphalidae, Lycaenidae, Papilionidae –, Pieridae (-)

**Bombycoidea:** Bombycidae (-), Saturniidae (-), Anthelidae –, Apatelodidae –, Endromidae –, Eupterotidae –, Hibrididae –, Lacosomidae –, Lasiocampidae –, Lemoniidae –, Brahmaeidae –, Carthaeidae –, Mirinidae, Oxytenidae –, Cercophanidae –

**Sphingoidea:** Sphingidae (now regarded as bombycoids by Minet 1986)

**Thyridoidea:** Thyrididae

**Calliduloidea:** Callidulidae –, Pterothysanidae –

**Pyraloidea:** Pyralidae (Galleriinae, Phycitinae, Pyralinae), Crambidae (Crambinae, Pyraustinae, Nymphulinae, Scopariinae, Musotiminae)

**Geometroidea:** Geometridae (Ennominae, Geometrinae –, Larentiinae, Oenochrominae, Sterrhinae)

**Axioidae:** Axiidae –

**Drepanoidea:** Drepanidae (Thyatirinae) (-)

**Uranioidea:** Uraniidae (including Epipleminae) –

**Mimallonoidae:** Mimallonidae –

**Noctuoidea:** Arctiidae, Ctenuchidae, Noctuidae (Agaristinae, Acronictinae, Cuculliinae, “Catocalinae” (-), Chloephoriinae (-), Euteliinae –, Hadenniinae, Heliothinae (-), Herminiiinae –, “Hypeninae”, Hypenodinae, Noctuinae, Nolinae, “Ophiderinae”, Plusiinae (-), Stictopterinae –), Notodontidae –, Thaumetopoidea –, Diptidae –, Thyretidae –, Lymantriidae (-), Aganaidae –)

(Families of unknown position: Sematuridae –, Apogonidae –, Galacticidae –, Pterolonchidae –, Lathrotelidae –, Epicopeidae –)

Superfamily unknown: ***Titanomis*** Meyrick; “***Lysiphragma*** ***argentina*** Salmon (both apodityrsian; see p. 214).

“Classification of the Lepidoptera reflects widely differing states of knowledge in different taxa” (Hodges 1983, p. xiv), a statement as true for New Zealand as for North America. What follows is a personal view of taxonomic treatments relevant to families represented in New Zealand by endemic or indigenous taxa. An asterisk denotes families with economically important members either in New Zealand or with a history of interception at our ports of entry.

**Table 1** New Zealand Lepidoptera; numerical status, by family, of valid species (Spp.), endemics (End.), species naturally shared with Australia (NZ / Aus.), and vagrant and man-adventive species, both established (E) and non-establishing (N).

	Spp.	End.	NZ / Aus.	Vagrant (E)	Vagrant (N)	Man-adv. (E)	Man-adv. (N)
Micropterigidae	20	20					
Nepticulidae	32	31				1	
Mnesarchaeidae	10	10					
Hepialidae	28	28					
Prodoxidae	1	1					
Tineidae	101	86	2	2		11	
Psychidae	51	47		1		3	
Roeslerstammiidae	2	2					
Gracillariidae	23	18					5
Yponomeutidae	45	42	2	1			
Glyptopterigidae	34	34					
Lyonetiidae	5	1	1		1	2	
Gelechiidae	50	42		3		5	
Batrachedridae	9	9					
Blastobasidae	2			1		1	
Coleophoridae	3					3	
Cosmopterigidae	10	7	3				
Lecithoceridae	1					1	
Momphidae	3	2	1				
Oecophoridae	271	259	2	3		7	
Scythrididae	6	6					
Elachistidae	25	23	1		1		
Cossidae	1						1
Sesiidae	1						1
Choreutidae	24	23	1				
Tortricidae	185	174	4	1		6	
Copromorphidae	3	3					
Carposinidae	25	25					
Epermeniidae	1	1					
Pterophoridae	21	18	1	1			1
Lycaenidae	9	7	2				
Nymphalidae	19	8	2		7	2	
Pieridae	2				1	1	
Bombycidae	1					1	
Saturniidae	6					2	4
Sphingidae	3		1		2		
Thyrididae	1	1					
Pyralidae	22	7	4	1	2	8	
Crambidae	233	226	3	3	1		
Geometridae	285	277	5	1	1	1	
Thyatiridae	1						1
Arctiidae	8	4	(1)	1	2	1	
Ctenuchidae	1						1
Lymantriidae	1						1
Noctuidae	174	138	8	9	17	1	1
Affiliation unknown	2	2					
Total:	1761	1582	43	28	35	61	12
% of total:		89.8	2.4	1.6	2.1	3.5	0.6

(a) Families in which revisionary work has been or is being done in a *world* context: Micropterigidae, Nepticulidae, Mnesarchaeidae, Hepialidae\*, Tineidae\*, Tortricidae\*, Gelechiidae (part), Crambidae (Crambinae\*).

(b) Families in which some revisionary work has been or is being done in a *local* context: Noctuidae\*, Geometridae\*, Yponomeutidae (in a broad sense)\*, Choreutidae, Momphidae, Tineidae\*, Psychidae, Elachistidae, Arctiidae,

- Nymphalidae.
- (c) Families newly recognised in New Zealand. Epermeniidae, Roeslerstammiidae, ?Prodoxidae.
  - (d) Families most in need of revision once overseas classification makes this practicable (usually families with a confused or contradictory subfamily classification or little recent overseas information): Oecophoridae\* (including Depressariidae\*), Noctuidae\*, and the taxa *Titanomis* and "*Lysiphragma*" *argentaria*.
  - (e) Families that practicably can be revised now: Psychidae\*, Tineidae\*, Gracillariidae\*, Yponomeutidae\*, Glyphipterigidae\*, Crambidae (Scopariinae\*), Gelechiidae\*, Carposinidae\*, Elachistidae, Cosmopterigidae\*, Choreutidae, Pyralidae\*, Geometridae\*.

## COMPOSITION OF LEPIDOPTERA FAUNA OF NEW ZEALAND

(see Table 1)

### Degree of endemism

Of the 1760 or so species recognised in this catalogue, 1582 (89.8%) are endemic. This high degree of endemism is also characteristic of the Hawaiian Islands, and is approached by Madagascar and St Helena.

### Proportion of non-ditrysian groups

New Zealand has five of the eleven recognised non-ditrysian superfamilies. Nielsen (1985a) has pointed out that about 98–99% of the known Lepidoptera fauna is in one structural group, the (endoporian) Ditrysia, characterised by having the common oviduct dorsal to the bursa copulatrix, and these joined by an internal ductus seminalis. The remaining 1%, or 'non-Ditrysia', are classed in several profoundly distinct divisions. These 'primitive' groups are well represented in South America and Australia (Nielsen 1985a), and at the species level in New Zealand form 5% of our total fauna.

One superfamily, the Mnesarchaeoidea, is endemic. Various features, including the porrect antennae, pseudofrenular hindwing structures, and porrect, scaled maxillary palpi, are distinctive within Exopia.

A feature of New Zealand's non-ditrysian fauna is the lack of the superfamilies Agathiphagoidea (but host *Agathis* is present), Heterobathmioidea (but host *Nothofagus* is present), and Eriocranoidea (but one host family, Fagaceae, is present). Palaephatoidea are also absent, although present in Australia (Nielsen 1985). Heterobathmioidea, Tischerioidea, and Neopseustoiidea are also absent from Australia, and in Incurvarioidea the sole New Zealand representative has characters of the Northern

Hemisphere and southern South American family Prodoxidae (cf. *Lampronia*) rather than of the Heliozelidae, Adelidae, and Incurvariidae, which are present in Australia.

### Ditrysia representation

New Zealand has 11 of the 28 ditrysian superfamilies recognised by Minet (1983) and Nielsen & Common (in prep.). Some major absences are Cossoidea, Zygaenoidea, Hesperioidae, Drepanoidea, and Bombycoidea; these account for a significant proportion of the lepidopteran fauna of North America (Hodges 1983) and Australia (Common 1970). Largely tropical superfamilies such as Immidoidea, Alucitoidea, Hyblaeoidea, Calliduloidea, and Uranioidae are also absent, although other superfamilies are represented in New Zealand by largely tropical groups, e.g., in Thyridoidea, and Copromorphidae in Copromorphaeidea. *Vanicela* in Tineoidea and *Lopharcha* in Tortricoidea are also tropical elements in the New Zealand fauna.

Poor representation of Papilionoidea (with no Papilionidae) is an enigmatic feature of the New Zealand fauna, and at the family level, in contrast to eastern Pacific island faunas, the very small number of species in Cosmopterigidae is particularly striking. Conversely, the eastern Pacific is weak in Oecophoridae, a family well represented in New Zealand, Australia, and New Caledonia.

As in Australia, the major superfamilies are Gelechoidea, Geometroidea, Pyraloidea, and Tortricoidea. In both countries, within Gelechoidea, the Oecophoridae overwhelmingly predominate.

New Zealand's Lepidoptera fauna is dominated by a few large genera: *Tingena* in Oecophoridae, with over 80 genetically distinct species; *Orocrambus* in Crambidae, with over 50; and the *Eudonia* – *Scoparia* complex, with over 100. Most of our Noctuidae are in one subfamily, Hadenniae, and three-quarters of our Geometridae are in Larentiinae (with at least 15% in one genus, *Asaphodes*). In Choreutidae and Glyphipterigidae all but one species each are in a single genus.

Presence in New Zealand of suitable host families or genera is no guarantee of the presence here of phytophagous Lepidoptera which are restricted to them elsewhere, as noted above. Moraceae genera support Copromorphidae here, as they do in Australia and Fiji, but we lack the choreutid genera *Eutromula* and *Tortyra*, and the danaine genus *Euploea*. Rubiaceae are well represented here, but we lack Sphingidae, which are a feature of the fauna on Pacific Rubiaceae.

### 'Microlepidoptera / macrolepidoptera' proportions

The artificial (but still practical) division into micros and macros shows some striking differences between

faunas. Figures derived from this catalogue, Common (1970), Kloet & Hincks (1972), Karsholt & Nielsen (1976a), and Hedges (1983) indicate that in New Zealand, Australia, Great Britain, and Denmark respectively 'micros' form over 50% of lepidopterous faunas, contrasting curiously with North America, where 'micros' form only 34%. This apparent disparity should perhaps be viewed more as an artefact than a reflection of reality.

### Fossil representation

Only one lepidopterous fossil has been reported, a pink scale embedded in Oligocene coal from Glen Afton (WO) (Evans 1931, p. 99). Tillyard (*in Evans 1931*) considered it to be lepidopterous, and noted that Sphingidae and Hepialidae have similar 'fish-tail' wing-scales. Because of the pink colour, Tillyard favoured *Porina* (= *Wiseana*). Searches of fossil leaves for mines (Wilkinson, *in prep.*) have yielded at least one conclusive New Zealand leaf mine, and probably more.

### Relationships with other biogeographic areas

New Zealand biogeography is undergoing critical study from a panbiogeographic viewpoint (Craw 1985). For Lepidoptera, a major survey of migrants across the Tasman (Fox 1978) showed that (a) over the past 100 years, the species composition of the migrant fauna in any one decade was invariable, and (b) all species were widely distributed, either in Australia or in the Old World Tropics. The inference must be that dispersal over sea has made a minor contribution to our fauna.

The Kermadec Lepidoptera show no solely New Zealand relationships; all species common to both are present in Australia, and most species are found extensively in the subtropical Pacific. Species presumed to be endemic to the Kermadecs belong to groups with no close New Zealand relationships (see Appendix, p. 235); but the question of endemism is bound up with the state of interpretation of the relevant groups.

The two island groups in the Tasman Sea between Australia and New Zealand show contrasting degrees of New Zealand representation in Lepidoptera. Lord Howe Island (169°E, 32°S) is not known to have Lepidoptera with solely New Zealand relationships, whereas Norfolk Island (168°E, 29°S) has (Holloway 1977). Examples are in Geometridae (*Pseudocoremia*), Gelechiidae (*Anisoplaca*), and Momphidae (*Zapyrastra calliphana*, on Polygonaceae). The geometrid genus *Austrocidaria* is also present, as it is in eastern Australia. Neither island appears to support *Nyctemera* (Arctiidae). Other examples may be found as interpretation of microlepidoptera is refined.

As may be seen in Table 1 (p. 14), New Zealand shares with Australia about 6.1% of its Lepidoptera species, of which some 2.4% are assumed to be naturally shared, having been collected at the start of European exploration (e.g., *Bassaris itea*, collected during Cook's first voyage) or European settlement, and associated with an indigenous host (e.g., *Strepsicrates ejectana*, on *Kunzea* and *Lepidospermum*). Around 64 species (3.7%) are known or presumed to be migrant from Australia, and half of these have become established, either on adventive plants common to both countries or on Australian endemic plants naturalised in New Zealand. Species shared with Australia are in Arctiidae, Noctuidae, Geometridae, Crambidae, Sphingidae, Nymphalidae, Lycaenidae, Tortricidae, Oecophoridae, Gelechiidae, and Cosmopterigidae (Table 1).

The Chatham Islands are unusual in the virtual absence of Oecophoridae; in Tortricidae, the preponderance of taxa related to *Merophysa*; and the relatively high degree of local endemism (about 30%).

Above the species level, with its very high degree of endemism, the degree of autochthony is related to the degree of interpretation of each group worldwide. Only for Micropterigoidea is interpretation well advanced (Gibbs 1983); close relationships are shown with New Caledonia, and with Australia, and our representation mirrors that of Japan. Gaskin (1986) has discussed possible roles of the inner and outer Melanesian Arcs in regard to Diptychophorini (Crambidae) in New Zealand. Initial studies on Geometridae indicate strong Australian relationships for some groups (Craw 1986) and strong South American relationships for others.

Although trans-Tasman species pairs are known in several families, only the *Nyctemera annulata* - *N. amica* pair has been adequately analysed (Kay 1980). Other candidate pairs are known in Tineidae, Yponomeutidae, Glyphipterigidae, Batrachedridae, Cosmopterigidae, Elachistidae, Momphidae, Oecophoridae, Geometridae, and Noctuidae, but in each family they are the exception rather than the rule.

In some instances (*Tanaoctena* in Yponomeutidae, *Vanicela* in Roeslerstammiidae, "Horisme" in Geometridae), relationships include the Pacific as well as Australia, and it is worth noting that two New Zealand *Kessleria* species (Yponomeutidae) have seemingly identical counterparts on New Caledonia. Povolny (1977) demonstrated the close relationship between Nepalese *Empista* and New Zealand *Kiwaia* (Gelechiidae), and the oriental *Terricula* has genital and external similarities with *Ochetarcha* (Tortricidae).

## CONCEPTS AND CONVENTIONS USED IN THE CATALOGUE

The catalogue is essentially a chronicle of nomenclatural and historical actions. Its construction was guided by the principles listed in the Introduction to the third edition of the International Code of Zoological Nomenclature (1985) 1983. It is also based on the principle "see for yourself"; a principle that, given the accumulative nature of systematics collections practices, is possible to act on in nearly all instances. Its necessity is the basis of the Principle of the Name-bearing Type.

For the consequences of revisers not being enabled to see for themselves, see Duggdale (1986). Additional instances are found in D.E. Gaskin's revisions of New Zealand Crambinae, under *Glaucocharis* (several species) and *Orocrambus simplex*. All the wrong citations could have been averted had Gaskin been able to see for himself.

The basic unit of taxonomy is the type specimen, and the basic unit of systematics is the concept of a type population (the population in the locality at which the type specimen was captured). Therefore, we have to ascertain that the specimen indicated as the name-bearing type is in fact that entity as published, and that it did come from the type locality claimed.

**Recognition of the type.** To be a type, the specimen has to fit the following criteria. It, and its labels, must agree with the original published description, and the labels (to be original) must be of the period in which the species was described, and show idiosyncrasies restricted to the author and/or collector of the specimen. We can ask three questions about a type specimen:

- Is there additional information (in diaries, correspondence, reports) that can verify that this must be the type, and that the locality is correct?
- Who placed the 'Type' or 'Holotype' label on the specimen?
- Is the type status of a specimen implicit in some idiosyncrasy of the original author's way of labelling or positioning a type specimen?

**Neotypes.** Where the type is missing, or cannot be found, the temptation to choose a neotype is best ignored. No neotypes can be proposed in a catalogue; for *Gymnobathra dinocosma* (Meyrick) (Oecophoridae; p. 92) possible candidates are listed, one of which could replace the specimen stated by Meyrick (in a letter to G.V. Hudson) to have been demolished. In general, a neotype is chosen only if (a) the chooser personally saw the original destroyed, or if there is a statement to the effect that the original was destroyed, and (b) in the context of revisionary work.

**Type localities.** Pinpointing type localities can be difficult (e.g., Dr Hector's locality "Otago"), for a number of reasons: (1) the collector travelled or collected widely over a diverse province; (2) the locality name fell into disuse (e.g., Meyrick's "Taranaki" for New Plymouth); (3) a mistake was made (e.g., T.R. Oxley's collection stated to be from "Auckland", in reality came from Nelson). Difficulty is often resolvable by consulting old maps and gazetteers.

Difficulties 1 and 3 can be resolved by finding relevant letters (e.g., T.R. Oxley's collections), diaries (e.g., E. Meyrick's precise collecting localities), or a duplicate collection, with the same species representation and collecting date, and with authentic labels (i.e., in the collector's handwriting).

Another method is to compare present-day representation with that in the original collection. Oxley's "Auckland" collection included: (a) *Wisenana cervinata* (Walker), never collected in Auckland subsequently; (b) several species never subsequently (or previously) collected in the North Island; and (c) his *Graphania mutans* (Walker) and synonyms, differing considerably in size and colour pattern intensity from those of collectors previous (e.g., D. Bolton) and subsequent. Clearly, "Auckland" was the wrong recorded locality (and in the wrong island); a collection from Oxley labelled "Nelson 1860" in the Stainton Collection, and a letter from Oxley to Dr Monro, have made it clear that Nelson, in the South Island, was the general locality.

Some lapses were obvious; some of Meyrick's material from the subantarctic and the (subtropical) Kermadec Islands had the other's label data. As the two faunas are so distinct, no difficulty was encountered. Other lapses are more difficult (see *Proditrix megalyntha* (Meyrick), Yponomeutidae; *Mallobathra globulosa* Meyrick, Psychidae).

Haste may have caused Philpott to record the type locality of *Astrogenes insignita* Philpott (Tineidae) as Woodhaugh, Dunedin DN, and place his type designation on the one specimen in the type series from Waikaraka, Whangarei ND (see p. 60).

**Informal 'paratype' series.** Because Philpott, Fereday, Hudson, and Howes each sent to Meyrick parts of a series they regarded as representing a species, and retained the balance, specimens still present in their collections (at NZAC, CMNZ, NMNZ, and AMNZ respectively) are informal members of the type series. They have no nomenclatural significance except that their label data are often fuller (Meyrick and Guenée generally relabelled material). Systematically they are important as they may show a better representation of variation in the type population.

## Conventions

(1) Institutions housing types are indicated using the four-letter codes of Watt (1982), as follows.

AMNZ	Auckland War Memorial Museum, Auckland, New Zealand
ANIC	Australian National Insect Collection, CSIRO, Canberra, Australia
BMNH	British Museum (Natural History), London, England
CMNZ	Canterbury Museum, Christchurch, New Zealand
CNCI	Canadian National Collection of Insects, Ottawa, Canada
HCOE	Hope Entomological Collections, Oxford, England
IZWP	Institute of Zoology, Academy of Sciences, Warsaw, Poland
MNHN	Museum National de l'Histoire Naturelle, Paris, France
NHMW	Naturhistorisches Museum, Wien [Vienna], Austria
NMNZ	National Museum of New Zealand (formerly Dominion Museum), Wellington, New Zealand
NMVA	National Museum of Victoria, Melbourne, Australia
NRSS	Naturhistoriska Riksmuseet, Stockholm, Sweden
NZAC	New Zealand Arthropod Collection (formerly part of Cawthron Institute, Nelson), DSIR, Auckland, New Zealand
SMNZ	Southland Museum, Invercargill, New Zealand
VUNZ	Zoology Department, Victoria University of Wellington, New Zealand
ZILR	Zoologicheski Instityut Akademii Nauk, Lenigrad, Russia
ZMDK	Zoologisk Museum, Copenhagen, Denmark

(2) Type locality information, where differing from or a refinement of the author's published statement, is enclosed by brackets [ ]; a published emended type locality is enclosed by parentheses ( ).

(3) The symbol § denotes a name first published in summaries or abstracts in 'The New Zealand Journal of Science (Dunedin)' 1882–1885. This journal gave summaries of descriptive papers prior to their publication in full in the 'Transactions & Proceedings of the New Zealand Institute'. One stated aim of the journal was to ensure priority (*N.Z. Jl Sci (Dunedin)* 1 (1), p. 2). By and large the abstracts as printed are full of typographical errors.

(4) Superfamilies and families are arranged as in the list on p. 13. Within each family subordinate taxa are listed alphabetically (i.e., subfamilies within families, genera within subfamilies (or tribes), species within genera).

(5) Citations in the text are by author, date, page, and sometimes plate and/or figure number; all citations are given in full under References, pp. 215–234.

(6) The catalogue gives, for species, the species name, authority, date, page and/or figure numbers, and original genus (in parentheses). For each species name, whether synonymised or not, then follows: type locality, and area code (as in Crosby *et al.* 1976); collector; type status (HT, holotype; LT, lectotype; NT, neotype; PLT, paralectotype; PT, paratype, including allotype; ST, syntype) and gender; if only one specimen had been seen by the describer, the term 'unique' is added. Where the specimen is distinctively damaged, details are noted. Finally, the institution housing the type is listed (see section 1, above). Then follows cross-reference to Hudson's monographs and any other publications that are nomenclaturally relevant. Lastly, under Notes, any relevant further information is given.

For genera, the name, author, date, page number, type species, and status of the type designation are given.

With synonymies, reference to the first synonymiser is given. Although this information is not usually given in catalogues, it is necessary for interpretation of the nomenclatural history of a taxon.

Higher taxonomic categories, especially those recently recognised or redefined, usually have 'in the sense of' and an author, date, and page number, so that readers can (a) be certain whose concept is adopted, and (b) check for themselves.

(7) The Taxonomic Index covers all nominal taxa mentioned in the catalogue, with those regarded as synonyms in lighter type. Thus, the name *Agriophara coricopa* (in Stenomidae in previous publications) can be found in the index by looking for *Agriophara* or *coricopa*, and turning to the indicated page, where that binomen will be found to be a synonym of *Agriophara colligatella*, in Stenomatinae. Where a specific name is used in many genera, the generic names are in alphabetical sequence.

Good hunting.



## KEYS TO HIGHER TAXA IN NEW ZEALAND LEPIDOPTERA

**Cautionary note:** limitations of the keys. The following keys refer strictly to the groups found in New Zealand. In constructing them I have tried to answer the question "What characters used to recognise this group are least fallible?" Apart from facies (habitus, general appearance), which is often useful when dealing with greatly incomplete faunas such as New Zealand's, very few 'traditional' characters are uniformly present throughout the taxa they supposedly encompass, under the present classifications. The reader's attention is drawn to keys to more 'complete' faunas, which will be of more use to those identifying taxa intercepted in quarantine - e.g., Common (1970), Holloway *et al.* (in press), Nielsen & Common (in press). Conversely, the following keys cannot reliably be employed for other faunas.

Some genital and venation characters can be exposed by brushing away the vestiture with a fine sable brush or snipe feather. Otherwise, genitalia should be prepared as described by Robinson (1976). Gross venational characters can be briefly exposed by using a solvent (absolute alcohol, acetone, or xylene [carcinogenic!]) and examining with transmitted light. For more detailed study the techniques outlined by Zimmerman (1978, pp. 73-86), involving the use of bleaching agents, should be used. Wing vein notation is that recommended by Wootton (1971, pp. 91-92).

There are three sets of keys: (A) to superfamilies; (B) to families and subfamilies; (C) to brachypterous adults. Immature stages are not keyed.



### (A) KEY TO SUPERFAMILIES

- 01 Antenna apically clubbed (Fig. 40), the flagellar segments cylindrical, never pectinate or long-setose; forewing (Fig. 74a) lacking a jugum or retinaculum; hindwing (Fig. 74b) lacking a frenulum, with dorsal margin expanded, and often with a humeral vein; metascutellum (Fig. 95, 96) perpendicular, sunken dorsally; diurnal, broad-winged, brightly patterned Lepidoptera (butterflies: Fig. 171-173)  
... (p. 135) ... PAPILIONOIDEA
- Antenna apically tapering, the flagellar segments various, often pectinate, serrate, or long-setose; forewing with a jugum (Fig. 45a, 48a, 49a) or retinaculum (Fig. 78a, insert); hindwing usually with a frenulum (Fig. 50b; but not in groups with a jugum, nor in Bombycoidea), with dorsal

margin not expanded (except in Bombycoidea), and lacking a humeral vein; metascutellum horizontal, usually convex dorsally (Fig. 90-92) ... 02

- 02 (01) Forewing and hindwing (Fig. 45, 47-49) with similar venation, i.e., with 4 or 5 radial veins; forewing with a jugum; hindwing lacking a frenulum, and with 11 or 12 veins reaching the margin ... 03
- Forewing and hindwing with dissimilar venation, i.e., hindwing radial veins 2 or less; forewing with a retinaculum; hindwing usually with a frenulum, and with 10 or less veins reaching the margin ... 05
- 03 (02) Large to very large, stout moths (Fig. 142) with wingspan over 30 mm, or exceeding 70 mm; body thickly clothed in long, pilose scales; antennae less than  $0.25\times$  as long as forewing; middle and hind tibial spurs and bristles absent; hind leg shorter than middle leg; maxillary palpi and haustellum rudimentary or absent (New Zealand species)  
... (p. 56) ... HEPIALOIDEA
- Small to very small, slender moths (Fig. 139, 141) with wingspan 10 mm or less; body with sparse vestiture; antenna more than  $0.3\times$  as long as forewing; hind leg longer than middle leg; maxillary palpi either short, porrect, 3-segmented (often only 2 segments visible), or long, folded between segments 1 and 2 and 3 and 4 ... 04
- 04 (03) Maxillary palpi longer than labial palpi, 4-segmented, folded, the 4th segment prehensile; antennal scape dumbbell-shaped; mouthparts (Fig. 1) mandibulate; head loosely tufted with hair-like scales, appearing broader than thorax; mesoscutum without a midline suture; middle and hind tibiae with spine-whorls, but only hind tibiae with spurs; forewing pattern usually metallic; venation as in Fig. 45 ... (p. 52) ... MICROPTERIGOIDEA  
(Micropterigidae)
- Maxillary palpi shorter than prominent labial palpi, 3-segmented, porrect; antennal scape barrel-shaped; mouthparts (Fig. 3) haustellate; head thickly scaled, not appearing broader than thorax; mesoscutum with a midline suture (e.g., Fig. 92, 95, 96); middle and hind tibiae with spurs; forewing pattern not metallic; venation as in Fig. 47  
... (p. 55) ... MNESARCHAEOIDEA  
(Mnesarchaeidae)

05 (02) Ovipositor a dorsoventrally flattened, laterally serrate needle formed from abdominal tergite 8 (Fig. 117); vinculum an elongate 'V', over 5× as long as valva; valva with a pectinifer (information based on 2 incomplete pharate adults) ... (p. 59)	INCURVARIOIDEA (Prodoxidae)	
—Ovipositor not as above, but if acuminate then formed from papillae anales; vinculum not as above; valva without a pectinifer	... 06	
06 (05) Tympanal organs (e.g., Fig. 97–115) present, externally visible in region of thoraco-abdominal junction; usually a scaled 'collar' (e.g., Fig. 30, 31), divided on dorsal midline, between head and metathorax	... 07	
—Tympanal organs absent, or not externally visible at thoraco-abdominal junction; 'collar' indistinct, or appearing as an unbroken scale series (e.g., Fig. 14) between head and mesonotal scaling	... 09	
07 (06) Haustellum scaled at least basally (Fig. 26); hindwing with veins $Sc+R1$ and $Rs$ approximated or fused beyond discal cell (Fig. 76, 77); maxillary palpus porrect or ascending, scaled, often conspicuously so (Fig. 26); antennal socket often separated from eye margin by a scale band (Fig. 26)	... (p. 141)	PYRALOIDEA
—Haustellum nude; hindwing with veins $Sc+R1$ and $Rs$ divergent beyond discal cell; maxillary palpi minute or absent	... 08	
08 (07) Chaetosemata present (Fig. 27); tympanal organ complex formed entirely from basal abdominal pleurite and sternite 2; tympanum directed ventrolaterally (Fig. 104–108)	... (p. 163)	GEOMETROIDEA
—Chaetosemata absent (Fig. 28, 29); tympanal organ complex involving metathoracic (metepimeral) and basal abdominal structures; tympanal membrane on metathorax, directed posteriorly, and with a counter-tympanal hood or process on the basal abdominal pleura; often with a lobate hindwing axillary cord	... (p. 194)	NOCTUOIDEA
09 (06) Very large moths (over 60 mm wingspan) with stout bodies densely scale-covered; metanotum scaled; antennae pectinate (Fig. 136) or with a hook-tip (Fig. 39)	... 10	
—Slender, large or small or very small moths; metanotum usually naked or largely so; antennae pectinate or simple, never hook-tipped	... 11	
10 (09) Antenna pectinate, less than one-quarter as long as forewing; body stout, short, densely clad in woolly scales; forewing and hindwing both broad, similarly marked, and usually with an ocellate or lunate discal mark (Fig. 136)	... (p. 140)	BOMBYCOIDEA (Bombycidae, Saturniidae)
—Antenna hook-tipped (Fig. 39); body elongate, fusiform, streamlined; forewing narrow; hindwing less than three-quarters as long as forewing (Fig. 137)	... (p. 140)	"SPHINGOIDEA" (Sphingidae)
11 (09) Very small moths (4–8 mm wing span); fore tibia lacking an epiphysis; maxillary palpi 4-segmented, folded; labial palpi smaller than maxillary palpi, porrect to upcurved, drooping in dried specimens (Fig. 2); forewing with a strong, setose cubital retinaculum and a weak subcostal series of hooked scales (Fig. 46), and with $Rs$ stem vein strongly curved towards dorsum in the middle; short, stout moths (Fig. 140); antennal scape expanded into an eyecap; head with erect hair-scales; ovipositor lacking anal papillae, broadly truncate-triangular, flat (Fig. 116) (N.Z. species only) ... (p. 53)	... (p. 53)	NEPTICULOIDEA (Nepticulidae)
—If small, fore tibia with an epiphysis (Fig. 43, ep) and labial palpi porrect or ascending, projecting beyond maxillary palpi; forewing with a subcostal retinacular process (e.g., Fig. 75) in male and $Rs$ stem more or less straight, roughly parallel with costa	... 12	
12 (11) Haustellum naked, or absent	... 13	
—Haustellum scaled	... 20	
13 (12) Wings scaled only along veins (Fig. 67), i.e., membrane nude (ocelli prominent; wings each with a line of hooked setae, along $1A$ – $2A$ on forewing and along $Sc+R1$ on hindwing, the lines interlocking; body smoothly clothed in metallic scales) ... (p. 112)	... (p. 112)	part SESIOIDEA (Sesiidae)
—Forewings at least without nude areas; forewing without anal retinacular hooks and hindwing without such hooks on $Sc+R1$	... 14	

- 14 (13) Wings cleft (Fig. 73), the forewing with 2 lobes, the hindwing with 3; slender-bodied, long-legged moths (Fig. 170); resting posture T-shaped; maxillary palpi 1-segmented; hindwing ventrally often with specialised scales along *CuA*; venation as in Fig. 73  
... (p. 132) ... PTEROPHOROIDEA  
(Pterophoridae)
- Wings entire; hindwing without specialised ventral scaling ... 15
- 15 (14) Abdomen in male with a process on pleura of segment 2 (Fig. 94); slender moths with forewing termen emarginate (Fig. 169) ... (p. 132) ... EPERMENIOIDEA  
(Epermeniidae)
- Abdomen in male lacking a pleural process on segment 2 ... 16
- 16 (15) Forewings triangular, vein *Rs3* ending at truncate apex (Fig. 75); male retinaculum a long, slender hook (Fig. 75); aculum a long, slender hook (Fig. 75); both pairs of wings brown or orange-brown, with a netted pattern; wingspan 18–25 mm; facies as in Fig. 174; venation as in Fig. 75 ... (p. 141) ... THYRIDOIDEA  
(Thyrididae)
- Forewings not as above; *Rs3* extending only to costa; male *Sc* retinaculum a short thumb (Fig. 70, dotted); facies not geometroid ... 17
- 17 (16) Maxillary palpi minute or appressed; labial palpi either porrect or subascending, expanded dorsally and ventrally by scales (e.g., Fig. 22); ocelli usually present ... 18
- Maxillary palpi obvious, porrect or long and folded; labial palpi elbowed or roughly arcuate, at most only segment 2 expanded apically by scales ... 19
- 18 (17) Chaetosema and ocelli present (Fig. 21; ocelli sometimes rudimentary); lower frons with short, ascending scales; forewing (Fig. 69a) with *M* vestigial in discal cell, vein *CuA2* from three-quarters of cell length; male forewing (Fig. 70a) often with a costal fold; mesoscutum with a midline suture at least posteriorly (Fig. 92); ovipositor appearing as paired, leaf-like pads (Fig. 127), protruded (with short apophyses) at rest; hindwing sometimes with a broad cubital pecten (Fig. 69b, 70b) ... (p. 114) ... TORTRICOIDEA
- Chaetosemata absent, ocelli present or absent (Fig. 24); lower frons with descending scales; forewing (Fig. 71a, 72a) with *M* absent from discal cell, vein *CuA2* from close to discal cell apex; male forewing with no costal fold; mesoscutum lacking any trace of a midline suture; ovipositor more or less cylindrical, narrow, retracted at rest, greatly protrusible (apophyses long) (Fig. 128); hindwing with a linear cubital pecten (Fig. 71b, 72b), at least in female  
... (p. 129) ... COPROMORPHOIDEA
- 19 (17) Haustellum usually pallid, with galeae dissociated (Fig. 5); maxillary palpi porrect and ascending (Fig. 7) or long and folded (Fig. 5), or, if absent, haustellum obsolete; labial palpi often with bristles on segment 2 (and maxillary palpi long, folded) (Fig. 5), or, if up-curved (and maxillary palpi porrect), then smooth-scaled; head often with tufts of hair-like scales; male lacking pleural lobes on 8th abdominal segment; female often with post-cubital retinacular scales (Fig. 51)  
... (p. 59) ... TINEOIDEA
- Haustellum with galeae associated (locked); maxillary palpi porrect, or folded over haustellum base, or rudimentary (apparently absent); labial palpi lacking bristles, geniculate or ascending, often tufted ventrally on segment 2 (Fig. 8), the apical segment often longer than segment 2 (Fig. 8, 11); male with 8th abdominal pleura expanded (Fig. 124), or with large lobes enclosing the genitalia (Fig. 125); female with retinacular scales at discal cell base (Fig. 53)  
... (p. 73) ... YPONOMEUTOIDEA
- 20 (12) Haustellum usually clothed with scales over most of its length (except Elachistidae); maxillary palpi scaled, folded over haustellum base (Fig. 18); labial palpi recurved (if segment 2 expanded with scales then apical segment slender and usually acute) (Fig. 12–20); if haustellum scaled on basal quarter or less, then wings slender, forewing pointed, with termen very oblique; metascutum not divided by metascutellum (Fig. 91)  
... (p. 80) ... GELECHIOIDEA
- Haustellum clothed with scales on less than basal quarter and maxillary palpi unscaled, minute, pointing ventrally, under haustellum base (Fig. 23); stoutly built, small, dark moths (Fig. 165) with short, ascending labial palpi expanded dorsally and ventrally with black and

white scales (Fig. 23); wings tending oblong (termen scarcely oblique) (Fig. 68); metascutum divided by metascutellum (Fig. 93) ... (p. 113) ... part SESIOIDEA (Choreutidae)

**Notes.** (1) Coccoidea and Drepanoidea are not keyed because (a) coccoids are no longer imported (in logs of the host trees) and (b) the drepanoid imported for evaluation as a biological control agent against blackberry was never released, and the laboratory colony was destroyed.

(2) See catalogue section 'Species of uncertain family position' (p. 214) for notes on *Titanomis sisyruta* (venation as in Fig. 88) and "Lysiphragma" *argentaria* (venation as in Fig. 89).

**(B) KEYS TO FAMILIES AND SUBFAMILIES**  
(of Hepialoidea, Tineoidea, Yponomeutoidea, Gelechioidea, Tortricoidea, Copromorphoidea, Papilionoidea, Bombycoidea, Pyraloidea, Geometroidea, Noctuoidea)

#### Superfamily Hepialoidea

- 01 Forewing veins *Rs3* and *Rs4* arising separately along *Rs1+2* stem (Fig. 49)  
... "Oxycaninae" of Dumbleton  
—Forewing veins *Rs3* and *Rs4* stalked, from *Rs1+2* stem (Fig. 48) ... 02
- 02 (01) Antennal segments flattened, not pectinate (Fig. 33); male hind tibia with a hair pencil; body vestiture densely short-pilose, smooth; patterned in green (sometimes bluish or yellowish) on males, green and brown on females; wingspan sometimes exceeding 100 mm; hindwing in male (Fig. 48b) triangular, in female elongate ovate ... (p. 56) ... *Aenetus* group  
—Antennal segments bipectinate in both sexes (Fig. 32); male hind tibia lacking a hair pencil; body vestiture long-pilose, woolly; forewings patterned in shades of brown, often with whitish maculation; wingspan usually about 60 mm; male hindwing oval; autumn-emerging  
... (p. 56) ... *Aoraia* group

#### Superfamily Tineoidea

- 01 Mouthparts (haustellum, maxillary palpi) obsolete or absent (Fig. 6); (antenna in male pectinate (e.g., Fig. 144) or simple, the segments with dorsal scales only (Fig. 35, 36)); female winged or apterous or immobile, always with a tuft of long hair-scales surrounding ovipositor (Fig. 120)  
... (p. 67) ... Psychidae

—Haustellum and maxillary palpi usually present, but if absent, antennal segments with appressed scales on most of surface (*Lindera*); female without a tuft of long hair-scales surrounding ovipositor (Fig. 118, 119) ... 02

- 02 (01) Maxillary palpi long, folded (Fig. 5) and labial palpi elbowed, usually with bristles on segment 2 (but if maxillary palpi absent and labial palpi not bristled (*Lindera*), then fore and middle tibiae with apical spines); venation as in Fig. 50; ovipositor either protrusible with long apophyses (Fig. 118) or, if not protrusible, then ovipositor lobes pad-like, divergent, and apophyses short ... (p. 59) ... Tineidae  
—Maxillary palpi short, porrect or ascending (Fig. 7); labial palpi without bristles, curved or geniculate; in female, anterior and posterior apophyses very short, and anal papillae lobes appressed, more or less rhomboidal in side view (ovipositor scarcely protrusible) (Fig. 121, 122) ... 03
- 03 (02) Extremely slender, narrow-winged moths (Fig. 51, 147); antenna as long as forewing or longer, with each segment completely scaled, and scape more or less barrel-shaped; male lacking a gnathos (Fig. 130); venation as in Fig. 51  
... (p. 70) ... Gracillariidae  
—Narrow-winged (*Vanicela*) or broad-winged (Fig. 52) moths; if antenna as long as forewing (*Dolichernis*, Fig. 146) then segments scaled dorsally only; antennal scape flattened, with awning and pecten (Fig. 37) obscuring the compound eye; male with a gnathos (Fig. 129)  
... (p. 70) ... Roeslerstammiidae

#### Superfamily Yponomeutoidea

- 01 Ocelli prominent; maxillary palpi microscopic; labial palpi recurved (never geniculate), either dorsoventrally flattened (Fig. 10) or with segment 2 bearing a porrect scale tuft on ventral surface (Fig. 9); usually rather fulgent moths, with an eye-spot or black hook at forewing apex (Fig. 149); pleural area on abdominal segment 8 of male somewhat expanded (Fig. 124)  
... (p. 76) ... Glyphipterigidae  
—Ocelli absent; maxillary palpi rudimentary or 4-segmented, porrect, scaled; labial palpi not as above, or if with a tuft, then geniculate (Fig. 8); pleural area on abdominal segment 8 of male developed into 2 opposing flaps shielding the genitalia (Fig. 125) ... 02

- 02 (01) Antennal scape scales not forming an eye-cap; if moth slender-winged then head with smooth scaling, or with spreading tufts overhanging frons; maxillary palpi various, but usually porrect or ascending, scaled; labial palpi geniculate and tufted, or recurved, segment 3 often longer than segment 2 (exception: *Charixena*, which lacks mouthparts); antenna usually simple (bipectinate in *Tanaoctena*); hindwing with a *1A+2A* fork; male valvae with lateral brushes not projecting beyond valval apices  
... (p. 73) ... **Yponomeutidae**  
in a broad sense
- Antennal scape widened with pecten and awning (Fig. 11) to form a cap largely obscuring compound eye; slender-winged moths with head rough-scaled (Fig. 150); maxillary palpi rudimentary; hindwing lacking a *1A+2A* fork (Fig. 54); male valvae flanked by long, projecting brushes (Fig. 123)  
... (p. 79) ... **Lyonetiidae**
- Superfamily Gelechioidea**
- 01 Hindwing narrower than its fringe; forewing apex acuminate (i.e., termen very oblique, grading into dorsum) (Fig. 55)  
... 02
- Hindwing broader than its fringe; forewing apex rounded or acute, with termen oblique and termen-dorsum transition abrupt (Fig. 58)  
... 10
- 02 (01) Haustellum scaled on basal quarter or less (Fig. 16); forewing lacking a completely tubular anal fork (Fig. 55), and with vein *1A* vestigial  
... 03
- Haustellum scaled on basal half or more (e.g., Fig. 17); forewing with a completely tubular anal fork, or vein *1A* largely tubular  
... 04
- 03 (02) Haustellum not or scarcely exceeding apex of front coxa; forewing (Fig. 55a) lacking retinacular setae on underside of dorsum; hindwing (Fig. 55b) with *Rs* axial (*Sc+R1* diverging from base); male genitalia symmetrical  
... (p. 111) ... **Elachistidae**
- Haustellum extending beyond hind coxae, strongly transversely ribbed; forewing (Fig. 65a) with stiff, rod-like retinacular scales on underside of dorsum; hindwing (Fig. 65b) with *Rs* dorsal to long axis (and parallel to *Sc+R1*); a line of retinacular bristles between *Sc+R1* and *Rs*; male genitalia asymmetrical, often twisted  
... (p. 110) ... **Scythrididae**
- 04 (02) Head with vertex high-domed, postvertex long (Fig. 13, 14); forewings narrow from base, their acuminate apex tending to curve, flexible  
... 05
- Head with vertex planoconvex, not high-domed, postvertex short (e.g., Fig. 15); forewings tapering, if acuminate and flexible at apex then hind tibia and tarsal segments with whorls of spines  
... 06
- 05 (04) Antennal flagellum thickened with scales on basal half; coloration either dark metallic blackish-green or fawn-striped buff  
... (p. 85) ... **Coleophoridae**
- Antennal flagellum not thickened with scales; coloration never metallic (Fig. 152)  
... (p. 84) ... **Batrachedridae**
- 06 (04) Hind tibia and tarsi with apical whorls of bristles (Fig. 159); hind legs displayed, raised in repose; male antennal segments with whorls of long setae  
... (p. 108) ... **Oecophoridae: Stathmopodinae**
- Hind tibia lacking apical whorls of bristles; hind legs hidden in repose  
... 07
- 07 (06) Very small moths (wingspan 5–7 mm); head smooth, clad in depressed, shining scales; forewings with bands or patches of broad, silvery scales on a shining black ground; hindwing termen not emarginate (Fig. 155); abdominal tergites covered with short, close-set spinules; (male genitalia symmetrical)  
... (p. 87) ... **Momphidae**
- Larger moths (wingspan 6–12 mm); head not clad in shining scales, or if clad in coppery scales, then hindwing termen emarginate; abdominal tergites not covered in spinules; forewing without transverse bands of silvery scales  
... 08
- 08 (07) Hindwing with termen emarginate (Fig. 56)  
... (p. 80) ... part **Gelechiidae**
- Hindwing termen entire (e.g., Fig. 65)  
... 09
- 09 (08) Labial palpi slender, recurved, often longitudinally lined with white, and reaching to behind antennal bases (Fig. 15); forewing without a pterostigma (eyes usually red in fresh specimens); antennal scape not forming an eye cap, cylindrical; forewing pattern usually complex, but if banded, then in black and white  
... (p. 86) ... **Cosmopterigidae**
- Labial palpi thickened (with distinctive scale patches on inner face in male – Fig. 12); forewing with a pterostigma (e.g., Fig. 56, pt), elongate-oval; antennal scape forming an eye-cap; or if labial palpi slen-

- der and antennal scape cylindrical, then wing pattern consisting of 4 transverse dark bands on a yellowish ground (*Oegoconia*; adventive) ... (p. 85) ... **Blastobasidae**
- 10 (01) Hindwing termen emarginate; male genitalia usually with tegumen and uncus flattened, valvae articulating ventrolaterally (Fig. 131); wings lacking a tubular vein *CuP*; female forewing with a radial retinaculum, and hindwing with *A* veins straight (Fig. 56b) ... (p. 80) ... **Gelechiidae**  
 —Hindwing termen entire (Fig. 57–64, 66); male genitalia with tegumen (at least) convex (Fig. 132), valvae articulating laterally, rarely ventrolaterally; at least hindwing with vein *CuP* tubular; female forewing with a subcubital (and sometimes anal) retinaculum, or if retinaculum radial only, then at least one hindwing *A* vein sinuous (Fig. 57–64, 66) ... 11
- 11 (10) Forewing apex rounded (Fig. 58) or subacute; hindwing anal veins straight; female with retinaculum on *CuA* and *R* stems; antennal segments completely scaled; male gnathos with apex or apices spinose, knob-like (Fig. 133); ovipositor either narrow and protrusible (*Eutorna*) or a perpendicular oval pad (Fig. 126)  
 ... (p. 87) ... **Oecophoridae: Depressariinae**  
 —Wing apex usually subacute; at least one hindwing anal vein angled or sinuous (Fig. 60–64) (exception: *Donacostola*, *Xyloryctinae*); female with retinaculum radial, or extending from *CuP* or *2A*; wings held flat or angled roofwise; antennae either completely scaled or (more usually) scaled dorsally only; male gnathos arms not ending in 1 or 2 spinose knobs; female ovipositor narrow, protrusible ... 12
- 12 (11) Ocelli present (Fig. 19); forewing with a sparse overlay of long, strap-like scales  
 ... (p. 88) ... **Oecophoridae: Hierodoris group**  
 —Ocelli absent (Fig. 18); forewing lacking an overlay of strap-like scales ... 13
- 13 (12) Hindwing veins *Rs* and *M1* well separated at base, parallel or slightly divergent, often vestigial, and *Rs4* on forewing ending at apex (Fig. 60–62); abdomen often dorsally spinulose  
 ... (p. 90) ... **Oecophoridae: Oecophorinae**  
 —Hindwing veins *Rs* and *M1* approximated at base, connate or stalked (Fig. 63b, 64b), divergent or, if separate, *Rs4* in forewing ending on termen (*Donacostola*, Fig. 64c); if abdomen spinulose, then antenna thickened with yellow scales ... 14
- 14 (13) Hindwing with *Sc+R1* approaching *Rs* before end of discal cell (Fig. 63) (male antennae with long ciliations); forewing vein *Rs4* to apex, veins *CuA1* / *CuA2* separate (Fig. 63)  
 ... (p. 109) ... **Oecophoridae: Stenomatinae**  
 —Hindwing with *Sc+R1* diverging from well before end of discal cell; forewing veins *Rs4* to costa or termen; veins *CuA1* / *CuA2* separate or stalked ... 15
- 15 (14) Antenna as long as forewing, thickened with yellow scales; forewing vein *CuA2* stalked with *CuA1* (Fig. 66); abdomen dorsally spinulose; (a black moth with contrasting yellow antennae)  
 ... (p. 87) ... **Lecithoceridae**  
 —Antenna shorter than forewing, not thickened with scales; forewing vein *CuA2* arising on discal cell separately from *CuA1* (Fig. 64); abdomen without dorsal spinules; (pallid moths)  
 ... (p. 110) ... **Oecophoridae: Xyloryctinae**

#### Superfamily Tortricoidea (Tortricidae subfamilies)

- 01 Male genitalia with uncus area reduced or membranous, sometimes bilobed; hind wing with a dorsal cubital pecten; antennal segments with apparently 1 ring of scales ... (p. 114) ... **Olethreutinae**  
 —Uncus beak-like, projecting, curved, as a flattened strap, or paddle-shaped, apically bifid, or truncate, rarely as a slender hook; hind wing usually lacking a dorsal cubital pecten (present in *Ctenopseustis* (Fig. 69b, 70b) and *Epalxiphora*); antennal segments with 2 obvious scale rings (basal and apical) ... 02
- 02 (01) Male with uncus sigmoid, slender, and valva with an external pocket enclosing an elaborate scale tuft (which arises on segment 8) (Fig. 135); female with part of ductus bursae annulate; forewings with scattered tufts of raised scales  
 ... (p. 114) **Chlidanotinae (Polyorthini)**  
 —Male uncus not sigmoid, and valva without an external pocket (Fig. 134); female without an annulate section in ductus bursae; scale tufts, if present, restricted to basal quarter of wing ... (p. 117) ... **Tortricinae**

### **Superfamily Copromorphoidea**

- 01 Ocelli absent (Fig. 24); antennae simple or subserrate, long-ciliate in male; labial palpi long, porrect in female, usually obliquely ascending and truncate in male; long-winged moths (Fig. 167), often with raised patches of scales on forewing (Fig. 71a), and in males with an area of modified scales on hindwing; venation (Fig. 71) with hindwing  $Rs+M1$  fused, and  $M2$  vestigial ... (p. 130) ... **Carposinidae**  
—Ocelli present; antennae simple, short-ciliate in male and labial palpi long, porrect (*Isonomeutis*, Fig. 166), or antennae unipectinate (Fig. 38) and labial palpi short, ascending (*Phycomorpha*); venation (Fig. 72) with hindwing  $Rs$  and  $M1$  ending separately, and  $M2$  tubular ... (p. 129) ... **Copromorphidae**

### **Superfamily Papilioidea**

- 01 Eyes emarginate by antennal base (Fig. 25), or antennal base and eye margin contiguous; maxillary palpi absent; forelegs unmodified in female; wings largely blue, violet, or orange; wing span not exceeding 25 mm, usually less ... (p. 135) ... **Lycaenidae**  
—Eyes not emarginate, eye margin not contiguous with antennal socket; maxillary palpi 1-segmented or absent; foreleg fully developed in both sexes, or modified, lacking claws in both sexes; wing-span exceeding 30 mm ... 02
- 02 (01) Forelegs with tarsal claws in both sexes, the claws bifid; colour pattern largely white or yellow ... (p. 139) ... **Pieridae**  
—Forelegs lacking claws in both sexes, non-functional (i.e., not used for walking); colour pattern various ... **Nymphalidae** ... 03
- 03 (02) Antenna nude; male with eversible scent tufts flanking genitalia; female fore tarsus 4-segmented ... (p. 136) ... **Danainae**  
—Antenna with appressed scales (Fig. 40); male without scent tufts; female fore tarsus 1- or 5-segmented ... 04
- 04 (03) Discal cell closed apically by a vestigial  $m-m$  cross-vein; female fore tarsus 5-segmented ... (p. 136) ... **Nymphalinae**  
—Discal cell closed apically by a tubular vein (Fig. 74); veins  $Sc$ ,  $Cu$ , and  $A$  usually swollen at base (Fig. 74); female fore tarsus 1-segmented ... (p. 138) ... **Satyrinae**

**Note:** although *Opsiphanes* species have been intercepted in cartons of bananas from Ecuador, subfamily Brassolinae is not keyed.

### **Superfamily Bombycoidea**

(excluding Sphingidae; see p. 20)

- 01 Wings often crumpled, undulate, or hindwing with a fold; hindwing with 2 anal veins; wings without ocellate markings; domestic ... (p. 140) ... **Bombycidae**  
—Wings large, not undulate; hindwing with 1 anal vein; both wings with an ocellate or lunate mark or hyaline patch in discal cell (Fig. 136); feral ... (p. 140) ... **Saturniidae**

### **Superfamily Pyraloidea**

- 01 Forewing vein  $Rs4$  stalked with  $Rs3$  (Fig. 76); abdomen base with tympanic lines posteriorly convergent, forming a V (Fig. 98, 100); conjunctivum and tympanum on different, opposing planes (conjunctivum facing posterolaterally, tympanum anterolaterally) (Fig. 99, 102, 103); median ridge high, compressed, with long, divergent scales ('praecinctiorium'), sometimes with an anterior torulus tympani (Fig. 99, 102, 103) ... (p. 143) ... **Crambidae**  
—Forewing vein  $Rs4$  free or completely fused with  $Rs3$  (Fig. 77); abdomen base with tympanic lines parallel (except in *Achroia*) (Fig. 97, 101); conjunctivum (opaque) and tympanum (glassy) on same plane, i.e., flat (Fig. 97); left and right tympanal areas divided by a low midline ridge ('ventral Medianleiste'), which anteriorly may bear a long-scaled, sclerotised mound ('bourrelet', 'torulus tympani') (Fig. 97, tty) ... (p. 141) ... **Pyralidae**

**Notes.** (1) Minet (1981, pp. 272–278) keys the subfamilies recognised by him in these two families. As the first division of each key for each family involves preparation of the tympanal organs, and as, for instance, the Crambinae key out in four places, I have to regard Minet's system as somewhat impractical for many users of this Catalogue. Nevertheless, it will be worth testing.

(2) At present, Crambinae may be distinguished from other subfamilies in New Zealand Crambidae by the possession of a cubital pecten on the hind wing.

(3) The structures called "bourrelet" (Minet 1981, 1985a) or "torulus tympani" (Maes 1985) and "praecinctiorium" (Munroe 1972, p. 10; Minet 1981; Maes 1985) are developments of the ventral midline intersegmental zone or "ventral Medianleiste" (Kennel & Eggers 1933) between the metathorax and the anterior margin of abdominal sternite 2. In some Crambidae they include the ventral mesal prolongation of the tympanic bullae (e.g., Fig. 103, "*Heliothela*" *atra*).

(4) Another basic difference between the two families is the site of the internal base of the chordotonal nerve ('scoloparium') – anterodorsal on the tympanic bulla in Pyralidae, laterodorsal on a distinct flange ('processus tympani' of Maes 1985) in Crambidae (Kennel & Eggers 1933, pp. 54–55, fig. 12, 13).

### Superfamily Geometroidea

#### (Geometridae subfamilies)

(largely as in McGuffin 1957, p. 7)

- 01 Hind wing vein  $M_2$  absent (or non-tubular) (Fig. 78); forewing with  $Rs_{3-4}$  stalked (Fig. 78) ( $Rs_1$  appressed to  $Rs_{2-4}$  stem in *Declana*, Fig. 79); tympanal organs as in Fig. 104, 105 ... (p. 163) ... Ennominae
- Hindwing vein  $M_2$  tubular (e.g., Fig. 81) ... 02
- 02 (01) Hindwing veins  $Sc+R_1$  fusing with  $Rs$  only at base, with no cross-vein between them (Fig. 85) (p. 193) ... Sterrhinae
- Hindwing veins  $Sc+R_1$  fusing with  $Rs$  for at least half discal cell length; or  $Sc+R_1$  and  $Rs$  free from base, or with a cross-vein ... 03
- 03 (02) Hindwing veins  $Sc+R_1$  and  $Rs$  fused for more than one-quarter of discal cell length (Fig. 81), or separate but with an apical, tubular  $Sc+R_1-Rs$  cross-vein (*Paradetis*, Fig. 80; *Tatosoma*); tympanal organs as in Fig. 106... (p. 170) ... Larentiinae
- Hindwing veins  $Sc+R_1$  and  $Rs$  not fused, but often closely approximated (Fig. 84), or a basal  $Sc+R_1-Rs$  cross-vein present (*Dichromodes*, Fig. 83; *Xyridacma*, Fig. 82); tympanal organs as in Fig. 107 ... (p. 191) ... Oenochrominae

### Superfamily Noctuoidea

**Note.** Subfamily classification within Noctuidae has been critically examined by Kitching (1985). The key below is a temporary solution.

- 01 Haustellum absent; ocelli absent; female wingless, with pectinate antennae ... (p. 195) ... Lymantriidae
- Haustellum present; ocelli usually present; if female wingless, then antennae simple ... 02
- 02 (01) Counter-tympanal hood (where developed) pre-spiracular, i.e., spiracle on outside of hood (Fig. 109, 110); diurnal, black and white or black and yellow (Fig. 179), or white, black, and red-flecked or metallic species, usually with thin, unicolorous, close-scaled legs ... 03

- Counter-tympanal hood post-spiracular, i.e., spiracle on inside of hood (Fig. 111–115); usually nocturnal, but if diurnal (*Phalaenodes*), femora conspicuously furry, bicoloured ... Noctuidae ... 04
- 03 (02) Ocelli present; tympanal organs well developed (Fig. 109); wing pattern not metallic; hindwings as broad (and nearly as long) as forewings ... (p. 194) ... Arctiidae
- Ocelli absent; tympanal organs well developed (Fig. 110) or reduced; hindwings narrower and much shorter than forewings; metallic dark green moths, usually with part of ventral abdomen contrasting silvery white. (p. 195) ... Ctenuchidae
- 04 (02) Hindwing 'trifid' (i.e., veins  $M_3$ ,  $Cu_{A1}$ , and  $Cu_{A2}$  strong), with vein  $M_2$  arising closer to  $M_1$  than to  $M_3$  (Fig. 86) ... 05
- Hindwing 'quadrifid' (i.e., veins  $M_2$ ,  $M_3$ ,  $Cu_{A1}$ , and  $Cu_{A2}$  strong), with vein  $M_2$  arising closer to  $M_3$  than to  $M_1$  (Fig. 87) ... 09
- 05 (04) Eyes haired (Fig. 29); fore tibia lacking spines ... (p. 199) ... Hadeninae
- Eyes naked; fore tibia with or without spines ... 06
- 06 (05) Eyes fringed by long, fine hair-scales (Fig. 28) ... (p. 198) ... Cuculliinae
- Eyes not fringed by fine hair-scales ... 07
- 07 (06) Fore tibia lacking spines ... (p. 195) ... Acronictinae in the broad sense
- Fore tibia with prominent spines (Fig. 43, 44) ... 08
- 08 (07) Fore femur with a patch of spine-scales on mesal face (Fig. 44); pallid species suffused with yellow to orange; male antennae simple ... (p. 210) ... Heliothinae
- Fore femur lacking a spine-scale patch; grey, black-lined fawn, or dingy brownish species; male antenna usually pectinate ... (p. 212) ... Noctuinae
- 09 (04) Abdomen with spiracle 1 exposed, facing laterally; counter-tympanal chamber closed anteriorly by an erect metepimeral lobe (Fig. 111); large (over 35 mm wingspan), diurnal, broad-winged black moths with yellow and white markings ... (p. 196) ... Agaristinae
- Abdomen with no metepimeral lobe closing the counter-tympanal chamber; abdominal spiracle 1 inside counter-tympanal hood, facing anteriorly ... 10


- 10 (09) Ocelli present (large to very large moths, 30–100 mm wingspan) ... 11  
 —Ocelli absent (small moths, 10–18 mm wingspan) ... 13
- 11 (10) Thorax and basal abdomen strongly crested, and forewings held roofwise, apically acute, with termen straight, its dorsum emarginate (Fig. 138); forewing usually with brassy or silvery marks or blotch ... (p. 213) ... *Plusiinae*  
 —Thorax smooth, or weakly tufted, and abdomen smooth; forewings held flat, with termen slightly convex or sinuous or emarginate; forewing without brassy or silvery marks ... 12
- 12 (11) Labial palpi upright, or porrect to oblique, straight, their apical segment tufted, not clavate; antenna pectinate in male; tympanal organ as in Fig. 114, 115 ... (p. 211) ... ‘*Hypeninae*’  
 —Labial palpi upright or recurved, reaching or exceeding vertex, their apical segment slender, clavate; antenna simple; hind tibia with or without spines; hind leg often modified, not functional for walking; forewing often with a lunate discal mark ... (p. 196) ... ‘*Catocalinae*’ (includes *Ophiderinae* of authors)
- 13 (10) Antennal scape lacking anterior scale tuft (Fig. 31); hind tibia smooth-scaled; hind wing veins *Cu42* and *M3* arising from a common point; (slender, long-winged, brownish moths) ... (p. 211) ... *Hypenodinae* in the sense of Forbes (1954)  
 —Antennal scape with anterior scale tuft (Fig. 30); hind tibia with a basal dorsal tuft of hair-like scales; hindwing with a cross-vein between *Cu41* and *M3* bases (i.e., these veins arising separately on discal cell; (triangular-winged, greyish-white moths; Fig. 181) ... (p. 213) ... *Nolinae*

**Note.** As subfamily *Chloeophorinae* is represented by an unrepeatable record of one specimen collected 50 years ago, it is not keyed here. It is presumed not to have established despite the abundance of its adventive host (*Quercus*).


**(C) KEY TO BRACHYPTEROUS LEPIDOPTERA**

- 01 Haustellum (and often maxillary palpi) present in both sexes ... 02  
 —Haustellum and maxillary palpi absent in females ... 07
- 02 (01) Haustellum scaled, at least basally ... 03  
 —Haustellum totally nude ... 06
- 03 (02) Haustellum scaled at base only; labial palpi long, porrect; maxillary palpi erect, long-scaled; a scaled strip between antennal socket and compound eye ... (p. 143) ... *Crambidae*  
 —Haustellum scaled on more than basal fifth; maxillary palpi folded over haustellum base; labial palpi ascending, recurved. 04
- 04 (03) Ocelli absent ... (p. 87) ... *Oecophoridae*  
 —Ocelli present ... 05
- 05 (04) Haustellum very long (extending beyond hind coxae); forewings densely clad in pale scales underneath ... (p. 110) ... *Scythrididae*  
 —Haustellum shorter (and usually tightly coiled); forewings without dense, pale scaling ... (p. 80) ... *Gelechiidae*
- 06 (02) Ocelli present ... (p. 114) ... *Tortricidae*  
 —Ocelli absent ... (p. 163) ... *Geometridae*
- 07 (01) Body entirely covered in woolly or fluffy hair-like scales; fore tibia with an apical, tooth-like process ... (p. 194) ... *Arctiidae*  
 —Body not entirely covered in wooly, hair-like scales; fore tibia normal, i.e., apex truncate ... 08
- 08 (07) Ovipositor protrusible, surrounded laterally and ventrally by a long tuft of crimped or sinuous hair-scales; legs often reduced, often with less than 5 tarsal segments ... (p. 67) ... *Psychidae*  
 —“Ovipositor” area a perpendicular anogenital field, often hidden dorsolaterally by a dense tuft of straight hair-scales; legs always fully formed, always with 5 tarsal segments ... (p. 56) ... *Hepialidae*


## ILLUSTRATIONS




(1) Micropterigidae  
*Micropardalis doroxena*  
(denuded)



(2) Nepticulidae  
*Stigmella laqueorum*



(3) Mnesarchaeidae  
*Mnesarchaea acuta*  
(denuded)










(4) Hepialidae  
*Wiseana cervinata*  
(denuded)

---


**Figures 1–31 Heads.** Key: aos, antenno-ocular scales; ch, chaetosema; el, 'eyelashes'; em, emarginate eye margin; ha, haustellum; lp, labial palpus; md, mandible; mp, maxillary palpus; oc, ocellus; pe, pecten; pv, postvertex; sc, scape of antenna.

---







(11) Lyonetidae  
*Bedellia psamminella*




(12) Blastobasiidae  
*Blastobasis tarda*




(13) Batrachedridae  
*Batrachedra agaura*




(14) Coleophoridae  
*Batrachedra agaura*  
(dorsal view)




(15) Cosmopterigidae  
*Pyroderces apparitella*




(16) Elachistidae  
"Elachista" *gerasmia*




(17) Gelechiidae  
*Anisoplaca achyrota*




(18) Oecophoridae  
*Gymnobathra flavidella*  
(denuded)




(20) Oecophoridae  
*Gymnobathra flavidella*




(19) Oecophoridae  
"Heliostibes" illita




(21) Tortricidae  
*Ctenopseustis obliquana*  
(denuded)




(22) Tortricidae  
*Planotortrix notophaea*




(23) Choreutidae  
*Asterivora nivescens*




(24) Carposinidae  
*Heterocrossa rubophaga*




(25) Lycaenidae  
*Zizina l. labradus*  
(denuded; dorsal view)




(27) Geometridae  
*Epyaxa rosearia*



(26) Crambidae  
*Deana hybreasalis*




(28) Noctuidae  
*Austramathes purpurea*





(29) Noctuidae  
*Graphania ustistriga*  
(note hairy eyes)

(30) Noctuidae  
*Celama parvitis*  
(→ - tuft on scape)




(31) Noctuidae  
*Hypenodes costaestrigalis*




(32) Hepialidae  
*Aoraiia dinodes*, ♂  
(bipectinate)




(33) Hepialidae  
*Aenetus virescens*, ♂  
(simple, compressed)



(34) Hepialidae  
*Wiseana cervinata*, ♂  
(short bipectinate)




(35) Psychidae  
*Mallobathra globulosa*, ♂  
(whorled)




(36) Psychidae  
*Mallobathra lapidosa*, ♂  
(fasciculate)


Figures 32–40 Antennae. Key: aw, awning; pe, pecten.




(37) Roeslerstammiidae  
*Vanicela disjunctella*  
(scape, ventral view)




(39) Sphingidae  
*Agrius convolvuli*  
(hook-tipped)




(42) Oecophoridae: Stathmopodinae  
*Stathmopoda coracodes*  
(note pilose tibial scaling  
and apical bristle whorl)




(38) Copromorphidae  
*Phycomorpha metachrysa*  
(unipectinate)




(40) Nymphalidae  
*Argyrophenga antipodum*  
(clubbed)



(41) Micropterigidae  
*Micropardalis aurella*  
(hind leg)



(43) Noctuidae  
(generalised foreleg)




(44) Noctuidae  
*Heliothis armigera*  
(foreleg)


---

Figures 41–44 Legs. Key: br, bristles; ep, epiphysis; ps, patch of spines; sp, spurs; ts, tibial spines.


---




(45) Micropterigidae  
*Sabatinca aemula*




(48) Hepialidae  
*Aenetus virescens*, ♂



(46) Nepticulidae  
*Stigmella laqueorum*  
(c, ♂ hindwing)



(47) Mnesarchaeidae  
*Mnesarchaea acuta*






(49) Hepialidae  
*Wiseana cervinata*, ♂




---



**Figures 45–89** Wing venation, females, (a) forewing and (b) hindwing (unless otherwise stated). **Key:** **venation** – A, anal; Cu, cubital; M, medial; R, radial; Sc, subcostal; other – cf, costal fold; cp, cubital pecten; ju, jugum; psf, pseudofrenulum; pts, pterostigma; re, retinaculum; st, scale tuft. Lines intersecting wing margin denote venation boundaries, e.g., M / Cu.

---





(52) Roeslerstammiidae  
*Dolichernis chloroleuca*









(56) Gelechiidae  
*Thiotricha tetraphala*





(57) Oecophoridae: Depressariinae  
*Eutorna phaulocosma*





(60) Oecophoridae: Oecophorinae  
*Gymnobathra dinocosma*





(58) Oecophoridae: Depressariinae  
*Proteodes carnifex*




(59) Oecophoridae: Oecophorinae  
"Helostibes" *atychioides*





(61) Oecophoridae: Oecophorinae  
*Izatha peroneonella*





(62) Oecophoridae: Oecophorinae  
*Tingena appertella*




(63) Oecophoridae: Stenomatinae  
*Agriophara colligatella*




(64) Oecophoridae: Xyloryctinae  
*Donacostola notabilis*




(65) Scythrididae  
*Scythris epistrota*




(66) Timyridae  
*Lecithocera leucomela*




(69) Tortricidae  
*Ctenopseustis obliquana*




(67) Sesiidae  
*Synanthedon tipuliformis*




(68) Choreutidae  
*Asterivora combinatana*




(70) Tortricidae  
*Ctenopseustis obliquana, ♂*




(71) Carposinidae  
*Heterocrossa*  
 (generalised)




(74) Nymphalidae  
*Argyrophenga antipodum*








(72) Copromorphidae  
*Phycomorpha metachrysa*



(73) Pterophoridae  
*Platyptilia falcatalis*




(75) Thyrididae  
*Morova subfasciata*  
 (detail: ♂ retinaculum)



(79) Geometridae  
*Declana floccosa*  
(forewing Sc, R, and Rs)



(80) Geometridae  
*Paradetis porphyrias*



(82) Geometridae  
*Xyridacma alectoraria*  
(hindwing *Sc*, *R*, and *Rs*, basal)




(83) Geometridae  
*Dichromodes sphaerista*  
(hindwing *Sc*, *R*, and *Rs*, basal)




(84) Geometridae  
*Samana falcatella*  
(hindwing *Sc*, *R*, and *Rs*, basal)





(85) Geometridae  
*Scopula rubria*  
(hindwing *Sc*, *R*, and *Rs*, basal)





(86) Noctuidae  
*Graphania mutans*  
(hindwing)




(87) Noctuidae  
*Chrysodeixis eriosoma*  
(hindwing)




(88) ?Family  
*Titanomis sisyrta*




(89) ?Family  
"Lysiphragma" *argentaria*




(90) Tineidae  
*Lysiphragma epixyla*




(91) Oecophoridae  
*Phaeosaces coarctatella*




(93) Choreutidae  
*Asterivora nivescens*



(92) Tortricidae  
*Planotortrix orthopis*




(94) Epermeniidae  
*Thambotricha vates*




(95) Lycaenidae  
*Lycaena salustius*

(96) Nymphalidae  
*Argyrophenga antipodum*


**Figures 90–96** Thoracic structures, in dorsal view. Key:  
ds, dorsal spines; mss, mesoscutum; mssl, mesoscutellum;  
mts, metascutum; mtsl, metascutellum; pap, procumbent  
abdominal process; T, tergite.




(97) Pyralidae  
*Ephestia kuehniella*  
(ventral view)




(98) Crambidae  
*Eudonia sabulosella*  
(ventral view)



(99) Crambidae  
*Eudonia hemiplaca*  
(side view)




(100) Crambidae  
*Musotima nitidalis*  
(ventral view)




(101) Pyralidae  
*Ephestia kuehniella*  
(side view)


**Figures 97–115 Tympanal organs. Key:** bt, bulla tympani; bul, bulla; co, conjunctivum; cth, countertympanal hood; ex, euoxa; ep, epaulette; fty, fornic tympani (border); ml, Medianleiste; mssl, mesoscutellum; mtel, metepimeral lobe; mtm, metepimeron; mts, metascutum; mtsl, metascutellum; pr, praecinctiorium; sp1, abdominal spiracle 1; tl, tympanic line; tty, torulus tympani (bourrelet); ty, tympanum; vh, ventral hood.




(102) Crambidae  
*Deana hybreasalis*  
(side view)








(103) Crambidae  
"Heliothela" atra  
(side view)





(104) Geometridae  
*Pseudocoremia suavis*  
(side view)





(105) Geometridae  
*Declana floccosa*  
(side view)







(113) Noctuidae  
*Graphania mutans*  
(dorsal view)




(114) Noctuidae  
*Rhabda scotosialis*  
(side view)




(115) Noctuidae  
*Rhabda scotosialis*  
(dorsal view)




(116) Nepticulidae  
*Stigmella laqueorum*  
(ovipositor)




(117) Incurvarioidea  
'kamahi scribbler'  
(ovipositor, reconstructed)




(118) Tineidae  
*Erechthias hemiclistra*  
(ovipositor, ventral view)




(120) Psychidae  
*Mallobathra lapidosa*  
(abdomen, side view)



(119) Tineidae  
*Erechthias hemiclistra*  
(ovipositor, side view)




(121) Roeslerstammiidae  
*Vanicela disjunctella*  
(ovipositor, side view)




(122) Roeslerstammiidae  
*Vanicela disjunctella*  
(ovipositor, ventral view)


**Figures 116–135** Abdominal structures (129–135, male genitalia). Key: **gn**, gnathos; **ovp**, ovipositor; **pla**, pleural area; **pll**, pleural lobe; **unc**, uncus; **vlv**, valva; **vlp**, valval pocket.




(124) Glyphipterigidae  
*Pantosperma holochalca*, ♂  
(abdominal apex, dorsal view)




(123) Lyonetidae  
*Bedellia psamminella*, ♂  
(side view)




(125) Yponomeutidae  
*Plutella antiphone*, ♂  
(abdominal apex, dorsal view)




(126) Oecophoridae: Depressariinae  
*Proteodes profunda*  
(ovipositor, end view)




(127) Tortricidae  
*Planotortrix excessana*  
(ovipositor, ventral)




(128) Copromorphidae  
*Isonomeutis amauropa*  
(ovipositor, ventral)




(129) Roeslerstammiidae  
*Vanicela* sp.




(130) Gracillariidae  
*Caloptilia linearis*  
(note lack of gnathos)




(131) Gelechiidae  
*Anisoplaca achyrota*




(132) Oecophoridae  
*Gymnobathra dinocosma*




(133) Oecophoridae  
*Proteodes profunda*  
(note spinulose gnathos)




(134) Tortricidae  
*Planotortrix conditana*  
(note uncal neck)




(135) Tortricidae  
*Lopharca insolita*  
(note sigmoid uncus, valval pocket)



(136) Saturniidae  
*Antheraea eucalypti*



(137) Sphingidae  
*Agrius convolvuli*



(138) Noctuidae  
*Ctenoplusia albostriata*

---

**Figures 136–138** Habitus sketches of three distinctive body forms.


---

---

**Figures 139–181** Habitus sketches marking commencement of (indigenous) family-group sections in catalogue

---

**TAXONOMIC CATALOGUE**  
**Suborder ZEUGLOPTERA**  
**Superfamily MICROPTEROGOIDEA**  
**Family MICROPTERIGIDAE**




---

**Micropterigidae**  
(139) *Micropardalis aurella* Hudson

---

• ***Micropardalis*** Meyrick, 1912a, p. 22 (listed); 1912c, p. 124 (listed); 1912f, p. 7, description. Type species *Palaeomicra doroxena* Meyrick, by monotypy (Meyrick 1912f, p. 7).

***aurella*** Hudson, 1918, p. 62 (*Sabatinca*)  
Tararua Range WN, R.M. Sunley; LT ♀ here designated, labelled "804b", "slopes of Tararua Mountains, Sunley, '08 [Hudson's Register]", NMNZ.  
Hudson 1928, pp. 367–368, pl. xlvi fig. 20.

***doroxena*** Meyrick, 1888e, p. 92 (*Palaeomicra*)  
Waitakere Range AK, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 367, pl. xxxix fig. 27.

• ***Sabatinca*** Walker, 1863c, p. 511. Type species *Sabatinca incongruella* Walker, by monotypy.

***Palaeomicra*** Meyrick, §1885i, p. 592; 1886b, p. 180. Type species *Palaeomicra chrysargyra* Meyrick, by subsequent designation (Meyrick 1912f, p. 7). Synonymised by Meyrick (1912c, p. 124; 1912f, p. 7).

***aemula*** Philpott, 1924c, pp. 667–668 (*Sabatinca*)  
Cobb Valley NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 369, pl. xxxix fig. 17.

***aenea*** Hudson, 1923c, p. 181 (*Sabatinca*)  
Governor's Bay, Banks Peninsula MC, S. Lindsay; HT ♂ labelled "1069a", unique, NMNZ.  
Hudson 1928, pp. 370–371, pl. xlvi fig. 12.

***aurantiaca*** Philpott, 1924c, pp. 668–669 (*Sabatinca*)  
Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 369, pl. li fig. 26.

***barbarica*** Philpott, 1918, p. 132 (*Sabatinca*)  
Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 370, pl. xlvi fig. 19.

***calliaracha*** Meyrick, 1912c, pp. 124–125 (*Sabatinca*)  
Blue Cliffs FD, A. Philpott; HT ♂ unique, BMNH.  
Hudson 1928, p. 379, pl. xxxix fig. 23.  
Note. Blue Cliffs is "on Te Waewae Bay, west of Tuatepere" (Dollimore 1962, p. 104). Although Meyrick cited "Invercargill" on the HT label, he published the locality as "Bluecliff, Invercargill".

***caustica*** Meyrick, 1912c, p. 124 (*Sabatinca*)  
[Seaward Moss] SL, A. Philpott; LT ♂ here designated, labelled "Invercargill New Zealand AP 23.10.10", "*Sabatinca caustica* Meyr. 3/4 E. Meyrick det. in Meyrick Coll.", [selected by K.R. Tuck], BMNH.  
Hudson 1928, p. 369, pl. xxxix fig. 18.

***chalcophanes*** Meyrick, §1885i, p. 592; 1886b, p. 180 (*Palaeomicra*)  
Makatoku WA, E. Meyrick; LT ♂ here designated, labelled "Makatoku New Zealand 8.2.83", "*Sabatinca incongruella* Walker 2/14 E. Meyrick det. in Meyrick Coll.", "Lectotype *Palaeomicra chalcophanes* Meyrick teste K.R. Tuck 1978", BMNH.  
Hudson 1928, p. 370, pl. xxxix fig. 19, as *Sabatinca incongruella*.

***chrysargyra*** Meyrick, §1885i, p. 592; 1886b, p. 182 (*Palaeomicra*)  
Lake Wakatipu OL, E. Meyrick; LT ♂ here designated, labelled "Lake Wakatipu New Zealand 15.12.82", "*Sabatinca chrysargyra* Meyr. 4/12 E. Meyrick det. in Meyrick Coll.", [selected by K.R. Tuck], BMNH.  
Hudson 1928, p. 369, pl. xxxix fig. 17.

***demissa*** Philpott, 1923, p. 154 (*Sabatinca*)  
Te Wairoa BP, R.J. Tillyard; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 368, pl. xlix fig. 15.

***highwayi*** Philpott, 1927d, p. 90 (*Sabatinca*)  
Leslie Valley NN, W. Highway; HT ♀ unique, NZAC.  
Hudson 1939, p. 471, pl. lxi fig. 32.

***ianthina*** Philpott, 1921, p. 342 (*Sabatinca*)  
Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 369, pl. xxxix fig. 25.

***incongruella*** Walker, 1863c, p. 511 (*Sabatinca*)  
[Nelson NN], T.R. Oxley, 1860; LT ♂ designated by Kristensen & Nielsen (1979, p. 140), BMNH.  
Hudson 1928, p. 370, pl. xlvi fig. 18, as *eodora*.

***munda*** Felder & Rogenhofer, 1875, pl. clx fig. 38 (*Oecophora*). **New synonymy.**  
[Nelson NN], T.R. Oxley, 1864; HT (?gender) unique, BMNH.

Note. The HT comprises the left forewing and hindwing, stuck to a pin.

**eodora** Meyrick, 1918a, p. 134 (*Sabatinca*). Synonymised by Kristensen & Nielsen (1979, p. 140). [Shedwood Forest], Tapawera NN, G.V. Hudson; LT ♂ designated by Kristensen & Nielsen, BMNH. Hudson 1928, p. 370, pl. xlvi fig. 18.

**lucilia** Clarke, 1920, p. 35 (*Sabatinca*)

Kauri Gully, Northcote AK, C.E. Clarke; HT ♂ designated by Clarke, AMNZ. Hudson 1928, p. 371, pl. xlix fig. 9.

Note. Clarke's published type designation of a specimen from Waitomo WO (Clarke 1934, p. 16) is at odds with the specimen labelled "Sabatinca lucilia Clarke Type T. LII 1919".

**passalota** Meyrick, 1923, p. 169 (*Sabatinca*)

Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 369, pl. xlix fig. 14.

**quadrijuga** Meyrick, 1912c, p. 126 (*Sabatinca*)

Invercargill SL, A. Philpott; HT ♂ unique, BMNH. Hudson 1928, p. 369, pl. xxxix fig. 25.

**rosicoma** Meyrick, 1914a, p. 118 (*Sabatinca*)

Kaeo ND, G.V. Hudson; LT ♂ here designated, labelled "Sabatinca rosicoma Meyr. 3/4 E. Meyrick det. in Meyrick Coll.", "Kaeo New Zealand GVH 1.13", [selected by K.R. Tuck], BMNH. Hudson 1928, p. 368, pl. xxxix fig. 20.


**zonodoxa** Meyrick, 1888e, p. 91 (*Sabatinca*)

[Waitakere Range AK], E. Meyrick; LT ♂ here designated, labelled "Sabatinca zonodoxa Meyr. 11/11 E. Meyrick det. in Meyrick Coll.", "Auckland New Zealand 22.12.85", [selected by K.R. Tuck], BMNH. Hudson 1928, p. 368, pl. xxxix fig. 21.

Also 1 undescribed species (G.W. Gibbs, VUNZ).

—◎—

**Suborder MONOTRYRIA**  
**Superfamily NEPTICULOIDEA**  
**Family NEPTICULIDAE**



**Nepticulidae**

(140) *Stigmella oriastra* (Hudson)

● ***Stigmella*** Schrank, 1802, p. 169. Type species *Phalaena (Tinea) anomalella* Goeze, 1783, p. 168 (= *Tinea rosella* Schrank, 1802, p. 139), by subsequent designation (Walsingham 1907, p. 1008).

***Nepticula*** Heyden, 1843, p. 208. Type species *Tinea aurella* Fabricius, 1775, p. 666, by subsequent designation (Tutt 1899, p. 184). Synonymised by Wilkinson (1978, p. 19).

***cypracma*** Meyrick, 1916b, p. 419 (*Nepticula*) new combination

Wellington WN, G.V. Hudson; HT ♀ unique, BMNH. Hudson 1928, p. 355, pl. xxxiv fig. 4, as *Nepticula cypracma*.

***erechtitus*** Watt, 1924, p. 686 (*Nepticula*) new combination

Dunedin DN, M.N. Watt; LT ♀ here designated, labelled "Dn, Dec. 1910", "Nepticula tricentra", NMNZ. Hudson 1928, p. 356, pl. li fig. 28 (figured from a specimen reared by Watt), as *Nepticula erechtitus*.

***fulva*** Watt, 1921a, p. 215 (*Nepticula*) new combination

Dunedin DN, M.N. Watt; 20 ST, NMNZ. Hudson 1928, p. 356, pl. li fig. 29 (figured from a specimen reared by Watt), as *Nepticula fulva*.

***insignis*** Philpott, 1927d, p. 89 (*Nepticula*) new combination

[Salisbury's Opening], Mt Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC. Hudson 1939, p. 469, pl. lxi fig. 24 (figured from a specimen selected by Philpott), as *Nepticula insignis*.

***laqueorum*** Dugdale, 1971b, p. 117 (*Nepticula*) new combination

The Snares islands, P.M. Johns; HT ♂ designated by Dugdale, NZAC.

➤ Nepticulidae, *Stigmella*

---

***lucida*** Philpott, 1919, p. 225 (*Nepticula*) new combination  
Waitati DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1928, p. 355, pl. xlvii fig. 20 (figured from a ST), as *Nepticula lucida*.

***maoriella*** Walker, 1864b, p. 1008 (*Tinea*) new combination  
. Auckland AK, D. Bolton; LT ♂ here designated, labelled [by H. Durrant] "Tinea maoriella Wkr Cat. Lep. Het. BM 30 p.1008 (1864) Type ♀", "BMNH Microlepidoptera genitalia slide 21,650", BMNH.  
Not mentioned by Hudson.

***microtheriella*** Stainton, 1854, p. 302 (*Nepticula*)  
Palearctic. New Zealand: Nelson NN, ex *Corylus*, NZAC.  
Not mentioned by Hudson.

***ogygia*** Meyrick, 1889b, p. 187 (*Nepticula*) new combination  
Dunedin DN, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 355, not figured, as *Nepticula ogygia*.

***oriastra*** Meyrick, 1917a, p. 247 (*Nepticula*) new combination  
Otira Gorge WD, Stella Hudson; LT ♀ selected by P.A. Brown and here designated, labelled "Nepticula orias tra Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", "Otira Gorge New Zealand S.H. 3000' 1.16", BMNH.  
Hudson 1928, p. 356, pl. xl fig. 6 and 20 (figured from STs), as *Nepticula oriastra*.

***perissopa*** Meyrick, 1919, p. 354 (*Nepticula*) new combination  
Mt Egmont TK, G.V. Hudson; LT ♀ selected by P.A. Brown and here designated, labelled "Nepticula perissopa Meyr. 2/2 E. Meyrick det. in Meyrick Coll.", "Mt Egmont New Zealand GVH 3000' 2.18", BMNH.  
Hudson 1928, p. 355, not figured, as *Nepticula perissopa*.

***progama*** Meyrick, 1924b, p. 662 (*Nepticula*) new combination  
Bold Peak OL, G.V. Hudson; HT ♀ unique, abdomen missing, BMNH.  
Hudson 1928, p. 356, pl. li fig. 30, as *Nepticula progama*.

***progonopis*** Meyrick, 1921, p. 336 (*Nepticula*) new combination  
Mt Arthur NN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 356, pl. xlvi fig. 10, as *Nepticula progonopis*.

***propalaea*** Meyrick, 1889b, p. 187 (*Nepticula*) new combination  
Arthur's Pass NC/WD, 2000', E. Meyrick; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 355, not figured, as *Nepticula propalaea*.

***sophorae*** Hudson, 1939, p. 469 (*Nepticula*) new combination

Christchurch MC, M.N. Watt; LT ♂ here designated, labelled "Kowhai Christchurch 24/ii/26", Watt Coll., NMNZ.  
Hudson 1939, p. 469, pl. xli fig. 25 (figured from a ST), as *Nepticula sophorae*.

***tricentra*** Meyrick, 1889b, p. 187 (*Nepticula*) new combination


[Lyttelton], Christchurch MC, E. Meyrick; HT ♂ unique, BMNH.

Hudson 1928, p. 355, not figured, as *Nepticula tricentra*.

Also 16 undescribed species (Donner & Wilkinson, in press) (NZAC).

—♂—

**Suborder DITRYSIA EXOPORIA**  
**Superfamily MNESARCHAEOIDEA**  
**Family MNESARCHAEIDAE**



**Mnesarchaeidae**  
(141) *Mnesarchaea fusilella* (Walker)

• ***Mnesarchaea*** Meyrick, §1885i, p. 591; 1886b, p. 180. Type species *Mnesarchaea paracosma* Meyrick, by original monotypy.

***acuta*** Philpott, 1929a, p. 304 (*Mnesarchaea*)  
Seven specimens (only 1 with abdomen) labelled "Arthurs Pass 9.1.20" (possibly collected by R.J. Tillyard) under this name, but none labelled as HT, AT, or PT, Philpott Collection, NZAC.  
Hudson 1928, pl. xxxix fig. 24, as *Mnesarchaea hamadelpha*; 1939, p. 470, as *M. acuta*.

***fallax*** Philpott, 1927a, p. 709 (*Mnesarchaea*)  
Mt Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 367, not figured.

***fusca*** Philpott, 1922, p. 82 (*Mnesarchaea*)  
Gouland Downs NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 367, pl. 1 fig. 2.

***fusilella*** Walker, 1864b, p. 1008 (*Tinea*) new combination  
Auckland AK, D. Bolton; HT ♂ unique, genitalia slide no. 21,780, BMNH.  
Not mentioned by Hudson.

***loxoscia*** Meyrick, 1888e, p. 90 (*Mnesarchaea*).  
New synonymy.  
Waitakere Range AK, E. Meyrick; LT ♀ here designated, labelled "Mnesarchaea loxoscia Meyr. 15/15 E. Meyrick det. in Meyrick Coll.", "Auckland New Zealand 22.12.85", BMNH.  
Hudson 1928, p. 367, pl. xxxix fig. 23.

***hamadelpha*** Meyrick, 1888e, p. 91 (*Mnesarchaea*)  
Mt Arthur NN [descent, 3000–1500'], E. Meyrick; LT ♂ here designated, labelled "Mnesarchaea hamadelpha Meyr. 5/3 E. Meyrick det. in Meyrick Coll.", "Mt Arthur New Zealand 19.1.86", BMNH.  
Hudson 1939, p. 470, pl. lxi fig. 8.

***similis*** Philpott, 1924c, p. 667 (*Mnesarchaea*).  
Synonymised by Hudson (1928, p. 367) and Philpott (1929, p. 304).  
Cobb Valley NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 367, as synonym of *Mnesarchaea hamadelpha*.

***paracosma*** Meyrick, §1885i, p. 591; 1886b, p. 180 (*Mnesarchaea*)  
Lake Wakatipu OL, E. Meyrick; LT ♂ selected by J.S. Dugdale and here designated, labelled "Mnesarchaea paracosma Meyrick 7/5 E. Meyrick det. in Meyrick Coll.", "L. Wakatipu New Zealand 15.12.82", BMNH.  
Hudson 1928, p. 366, pl. xxxix fig. 26.

Also 4 undescribed species (VUNZ, NZAC).


—♂—

**Superfamily HEPIALOIDEA**

**Family HEPIALIDAE**

**Subfamily HEPIALINAE**

(in the sense of Dumbleton 1966, p. 927)



---

**Hepialidae**

(142) *Wiseana cervinata* (Walker)

---

• ***Aenetus*** Herrich-Schaeffer, 1855, p. 85. Type species *Hepialus ligniveren* Lewin, by subsequent designation (Kirby 1892, p. 891, as *Oenetus*); Australia.

***Charagia*** Walker, 1856a, p. 1570. Type species *Aenetus virescens* Doubleday, by subsequent designation (Kirby 1892, p. 891). Synonymised by Kirby (1892, p. 891).

Note. Paclt (1953, p. 144) regarded *Aenetus* Herrich-Schaeffer, 1855 as an invalid name, giving as reasons: (a) no species was cited; (b) no diagnosis was given. However, since no violence is done to at least one requirement each in Articles 12 and 16 of the International Code of Zoological Nomenclature (1961), Paclt's action is not followed here.

***virescens*** Doubleday, 1843, p. 284 (*Hepialus*) Waitemata AK, E. Dieffenbach; HT not in BMNH [lost; identity not in doubt].

Hudson 1928, pp. 357–359, pl. xlvi fig. 13 and 14, pl. xlvi fig. 13 and 14, as *Hepialus virescens*.

***rubro viridans*** White in Taylor, 1855, pl. 1 fig. 1 (*Hepialus*). Synonymised by Meyrick (1890, p. 211).

Hudson 1898, p. 129; 1928, p. 357, as synonym of *Hepialus virescens*.

***rubroviridans*** Stephens MSS., Walker, 1856a, p. 1570 (*Charagia*). Synonymised by Kirby (1891, p. 891).

[Auckland AK], A. Sinclair; HT ♀ unique, BMNH. Hudson 1898, p. 129; 1928, p. 357, as synonym of *Hepialus virescens*.

***fischeri*** Felder in Felder & Rogenhofer, 1874, pl. lxxx fig. 1 (*Charagia*). Synonymised by Meyrick (1890, p. 211).

[Auckland AK, ?F.R. Hochstetter]; HT ♀ unique, BMNH. Hudson 1898, p. 129; 1928, p. 357; as synonym of *Hepialus virescens*.

***hectori*** Butler, 1877, p. 380 (*Charagia*). Synonymised by Meyrick (1890, p. 211).

"North Island", J. Hector; HT ♀ unique, BMNH. Hudson 1898, p. 129; 1928, p. 357; as synonym of *Hepialus virescens*.

***virescens ab. albo-extremis*** Quail, 1903, p. 252 (*Charagia*). Synonymised by Hudson (1928, p. 357, footnote).

Palmerston North WI/WN, A. Quail; HT ♂, Illidge Collection, South Australian Museum.

● ***Aoraiā*** Dumbleton, 1966, pp. 930–931. Type species *Porina dinodes* Meyrick, by original designation.

***Trioxycanus*** Dumbleton, 1966, p. 943. Type species *Porina enysii* Butler, by original designation. Misidentified type species; new synonymy.

***aurimaculata*** Philpott, 1914, p. 121 (*Porina*)

Mount Cook MK, F.S. Oliver; HT ♂ unique, not found, not in CMNZ.

Hudson 1928, p. 360, pl. xlvi fig. 12 (HT), as *Porina aurimaculata*.

Note. NZAC and CMNZ hold specimens with golden patches on the forewing that match Hudson's illustration. A specimen matching the type colour pattern, from Mt Cook MK, has been deposited at CMNZ.

***dinodes*** Meyrick, 1890, p. 206 (*Porina*)

Invercargill SL, F.W. Hutton; LT ♂ designated by L.J. Dumbleton, CMNZ.

Hudson 1898, p. 132, pl. xiii fig. 8; 1928, pp. 360–361, pl. xlvi fig. 6 and 7; as *Porina dinodes*.

***enysii*** Butler, 1877, p. 381, pl. xlvi fig. 7 (*Porina*)  
new combination

"North Island", J.D. Enys; HT ♂ unique, antennae missing, BMNH.

Note. Butler's artist drew imaginary, simple antennae on his depiction of the unique specimen. Meyrick applied Butler's name to another entity with simple antennae and oxycanine forewing venation. Because he could not personally examine Butler's HT, Dumbleton (1966, p. 943) had no reason not to adopt Meyrick's misconception. The case has been submitted to the International Commission on Zoological Nomenclature. The entity described by Meyrick (1890, p. 207) – and subsequently Hudson (1898, p. 133, pl. xiii fig. 9 and 10; 1928, p. 361, pl. xlvi fig. 4–10), Philpott (1927c, fig. 19), and Dumbleton (1966, pp. 946–947, fig. 52–56) – is not that of Butler. See *Dumbletonius*, below.

***leonina*** Philpott, 1927a, p. 709 (*Porina*). New synonymy.

Mount Arthur Tableland NN, A. Philpott; HT ♂ unique, NZAC.

Hudson 1928, p. 361, not figured, as *Porina leonina*.

**senex** Hudson, 1908, p. 107 (*Porina*)  
Old Man Range CO, 4000 ft, J.H. Lewis; HT ♂ unique,  
NMNZ.  
Hudson 1928, p. 360, pl. xlivi fig. 1, pl. xlix fig. 20, as  
*Porina senex*.  
Note. Philpott (1923, p. 154) described the short-winged  
female, collected by F.S. Oliver.

**annulata** Hamilton, 1909, p. 48 (*Porina*). Syn-  
onymised by Hudson (1928, p. 360).  
Mount Aurum OL, H. Hamilton; HT ♂ unique, NMNZ.  
Hudson 1928, p. 360, as synonym of *Porina senex*.

**sp.**, Dumbleton 1966, p. 939, fig. 42 and 45–47  
(*Aoraia*)

Cobb Valley, Dun Mountain NN; Gordon's Knob  
NN/MB; Altamarlock MB; Craigieburn Range MC;  
[Rotoiti BR]; Dansey's Pass, DN (NZAC, NMNZ).

Also 8 undescribed species (NZAC).

—◎—

### Subfamily OXYCANINAE

(in the sense of Dumbleton 1966, p. 940)

● **Cladoxycanus** Dumbleton, 1966, pp. 948–949.  
Type species *Porina minos* Hudson, by original  
designation.

**minos** Hudson, 1905a, p. 357 (*Porina*)  
Ophir CO, J.H. Lewis; HT ♂ unique, labelled "595a",  
NMNZ.

Hudson 1928, p. 365, pl. xlvi fig. 3–5, as *Porina minos*.

**autumnata** Hudson, 1920, p. 277 (*Porina*). Syn-  
onymised by Hudson (1928, p. 365).  
Lower Hutt WN, A.V. Clere; HT ♂ unique, NMNZ.  
Hudson 1928, p. 365, as synonym.

● **Dioxycanus** Dumbleton, 1966, pp. 951–952. Type  
species *Porina fusca* Philpott, by original  
designation.

**fusca** Philpott, 1914, p. 121 (as *Porina fusca*)  
Bold Peak OL, C.C. Fenwick; HT ♂ designated by Phil-  
pott, not found in NMNZ (Dumbleton 1966, p. 955);  
PT ♂ labelled "Bold Peak, A. Philpott 27.xii.12",  
"Paratype ♂", available in NZAC.  
Hudson 1928, p. 364, pl. xlvi fig. 10 and 11, as *Porina*  
*fusca*.

**oreas** Hudson, 1920, p. 277 (*Porina*)  
Mount Egmont TK, A. Lysaght; LT ♂ designated by  
Dumbleton (1966, p. 953), NMNZ.  
Hudson 1928, pp. 364–365, pl. xlvi fig. 19 and 20, as  
*Porina oreas*.

**ascendens** Meyrick, 1921, p. 336 (*Porina*). Syn-  
onymised by Dumbleton (1966, p. 952).  
Mount Arthur NN, G.V. Hudson; LT ♀ designated by  
Dumbleton, BMNH.  
Hudson 1928, p. 365, pl. xlii fig. 21, as species.

**descendens** Hudson, 1923c, p. 180 (*Porina*). Syn-  
onymised by Dumbleton (1966, p. 952).  
Arthur's Pass NC/WD, H. Hamilton; LT ♂ designated by  
Dumbleton (1966, p. 954), NMNZ.  
Hudson 1928, p. 364, pl. li fig. 19, as species.

**gourlayi** Philpott, 1931, p. 36 (*Porina*). Synony-  
mised by Dumbleton (1966, p. 952).  
Flora Camp, Mount Arthur NN, E.S. Gourlay; HT ♂  
designated by Philpott, NZAC.  
Hudson 1939, p. 470, pl. lxii fig. 5 and 6, as species.

● **Dumbletonius** Dugdale, new name for *Trioxy-  
canus* Dumbleton, 1966, pp. 942 (key) and 943.  
Type species *Dumbletonius sylvicola* Dugdale, new  
name for *Porina enysii* of authors (see below).  
Note. See discussion under *Aoraia enysii*, above.

**characterifer** Walker, 1865a, p. 594 (*Hepialus*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1898, p. 133, pl. xiii fig. 11; 1928, p. 362, pl. xli  
fig. 11 and 12; as *Porina characterifer*.

**impletus** Walker, 1865a, p. 598 (*Hepialus*). Syn-  
onymised by Meyrick (1890, p. 208).  
[Nelson NN], T.R. Oxley; HT ♀ unique, BMNH.  
Hudson 1898, p. 133; 1928, p. 362; as synonym.

**sylvicola** Dugdale, new name for *Porina enysii* in  
the sense of Meyrick (1890, p. 207), Hudson  
(1898, p. 133, pl. xiii fig. 9 and 10), Philpott  
(1927c, p. 39, fig. 19), Hudson (1928, pp. 361–  
362, pl. xli fig. 4–10), and Dumbleton (1966, pp.  
940–943, fig. 52–56) but not of Butler (1877, p.  
381, pl. xlvi fig. 7).  
Wellington WN, ?collector; HT ♂ here designated, labelled  
"Wellington 25.1.10", NZAC.

Note. Philpott's, Hudson's, and Dumbleton's descrip-  
tions and illustrations are all based on Wellington  
specimens. The HT is a ♂ in good condition, with  
markings resembling those of *P. enysii* Butler but with  
the diagnostic hindwing colour – "pinkish brown,  
tinged with ochreous on the termen" (Hudson 1898,  
p. 133).

**unimaculatus** Salmon, 1948, p. 309 (*Porina*)  
Great Island, Three Kings Islands, E.G. Turbott; HT ♂  
unique, AMNZ.  
Dumbleton 1966, pp. 944–946, fig. 48–51 and 108.

● **Wiseana** Viette, 1961, pp. 38–39, replacement  
name for *Philpottia* Viette, 1950, pp. 72–73, preoc-  
cupied by *Philpottia* Broun, 1915 (Coleoptera). Type  
species *Pielus umbraculata* Guenée, by original  
designation.

>> Hepialidae, *Wiseana*

**cervinata** Walker, 1865a, p. 595 (*Elhamma*)  
[Nelson NN], T.R. Oxley, 1860; HT ♂ unique, BMNH.  
Hudson 1898, pp. 133–134, pl. xiii fig. 12 and 18; 1928,  
p. 362, pl. xlivi fig. 5. Dumbleton 1966, fig. 110 (HT  
♂).

Note. Under *cervinata* I synonymise those names that apply to adults from WO-DN emerging in late spring to early summer and which have no consistent morphological differences either within or between populations. Hudson (1928, pl. xlivi fig. 5–7) illustrates variation in *W. cervinata*.

**despecta** Walker, 1865, p. 594 (as *Hepialus despectus*). New synonymy.

[Nelson NN], T.R. Oxley, 1860; HT ♀ unique, BMNH.  
Hudson 1898, p. 134; 1928, p. 362, as species. Dumbleton  
1966, fig. 112 (HT).

**vexata** Walker, 1865a, pp. 597–598 (*Porina*).  
Synonymised by Meyrick (1890, p. 208).

[Nelson NN], T.R. Oxley, 1860; LT ♂ here designated, labelled “Type” (circular, red-margined label), “New Zealand” // “60-73”, with 39.5 mm wing span, BMNH.  
Hudson 1898, p. 133; 1928, p. 362; as synonym.

**variolarus** Guenée, 1868, p. 1 (*Pielus*). Synonymised by Meyrick (1890, p. 208).

[Fendalton, Christchurch MC], R.W. Fereday; HT ♂ MNHP; [2 ♂♂ labelled “25.10.64 Fendalton flying at dusk” and 1 ♂ labelled “1st week Nov ‘64 Fendalton at light on window”, Fereday Collection, CMNZ.]  
Hudson 1898, p. 133; 1928, p. 362; as synonym.

**copularis** Meyrick, 1912c, p. 123 (*Porina*)  
Invercargill SL, A. Philpott; LT ♂ designated by L.J.  
Dumbleton, BMNH.

Hudson 1928, p. 363, pl. xlvi fig. 8 and 9 (topotypic ♂  
and ♀). Dumbleton 1966, fig. 111 (LT).

Note. This name refers to adults with slender wing scales and oblong antennal pectinations, emerging mid November to February. The name “despecta” was applied in error to this entity by Dumbleton (1966) and in subsequent agricultural literature.

**fuliginea** Butler, 1879a, p. 488 (*Porina*)  
“Otago”, F.W. Hutton; HT ♂ unique, BMNH.  
Hudson 1898, p. 133; 1928, p. 362; as synonym.

Note. This species is removed from synonymy with *cervinata* (cf. Meyrick 1890, p. 208; Dumbleton 1966, p. 959) as it has consistent differences in morphology and in electrophoretic characteristics (G. MacArthur, pers. comm.).

**jocosa** Meyrick, 1912c, p. 124 (*Porina*)  
Invercargill SL, A. Philpott; LT ♂ designated by Dumbleton, BMNH.  
Hudson 1928, p. 363, pl. xlvi fig. 1 and 2 (topotypic ♂).  
Dumbleton 1966, fig. 115 (LT).

Note. This species is removed from synonymy with *cervinata* (cf. Dumbleton 1966, p. 959) as it has consistent morphological differences, and is the only *Wiseana* with short, broad forewing scales.

**mimica** Philpott, 1923, p. 153 (*Porina*)

West Plains, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 362, pl. xlvi fig. 15 (topotypic ♂). Dumbleton 1966, fig. 109 (HT).

Note. This species is removed from synonymy with *cervinata* (cf. Dumbleton 1966, p. 969) as it has consistent morphological differences. The lowland flight period is early spring.

**signata** Walker, 1865a, p. 1563 (*Elhamma*)

[Auckland AK], D. Bolton; LT ♂ designated by Dumbleton (1966, p. 1964), BMNH.  
Hudson 1898, pp. 134–135, pl. xiii fig. 15; 1928, pp. 363–  
364, pl. xlvi fig. 2–4.

**novae zealandiae** Walker, 1856a, p. 1573 (*Porina*).

Synonymised by Meyrick (1890, p. 210).  
[Auckland AK], A. Sinclair; LT ♂ here designated, labelled “45-61” “New Zealand”, “Type” (circular, green-margined label), BMNH.  
Hudson 1898, p. 134; 1928, p. 363; as synonym.

**umbraculata** Guenée, 1868, p. 1 (*Pielus*)

[Fendalton, Christchurch MC], R.W. Fereday; LT ♂ designated by L.J. Dumbleton, MNHP; [1 ♂, 1 ♀ labelled “21.10.64 window at night, Fendalton” in Fereday Collection, CMNZ.]

Hudson 1898, p. 134, pl. xiii fig. 14; 1928, p. 363, pl. xlvi fig. 1–3.

• Also 1 undescribed genus and species.

### Names of uncertain identity

**mairi** Buller, 1873, p. 279 (*Porina*)

Ruahine Range RI/WA, W.L. Buller; HT lost when the barque ‘Assaye’ was wrecked in 1890 (Hudson 1898, p. 132). The illustration in Buller’s paper suggests a large *Dumbletonius characterifer*.

### Excluded species

**Epialus antipoda** “Bd” [Boisduval]: Herrich-Schaeffer 1853, pl. x fig. 44 (illustration); 1855, p. 86 (text)

Wagner & Pfitzner (1911, p. 16) record it from “N. Seeland” (in *Porina*), but the true locality is Australia (E.S. Nielsen, pers. comm.).

**Epiolus hyalinatus** Herrich-Schaeffer 1853, fig. 50 (illustration); 1855, p. 86 (text); as *Abantiades diaphanus* HS

Wagner & Pfitzner (1911, p. 15) record it from “N. Seeland, Tasmanien” (in *Pielus* Walker).

—⊗—

**Phalanx INCURVARIINA**  
(in the sense of Minet 1984, p. 147)

**Superfamily INCURVARIOIDEA**  
(in the sense of Minet 1984, p. 147)


Larval, pupal, and adult remains of a species referable to this superfamily, and possibly to Prodoxiidae as defined by Nielsen & Davis (1985), have been found at the end of cortical mines on *Weinmannia* species (Cunoniaceae).



**Suborder DITRYSIA ENDOPORIA**

**Superfamily TINEOIDEA**  
(in the sense of Minet 1986)

**Family TINEIDAE**



---

**Tineidae**  
(143) *Erechthias charadrota* Meyrick

---

Note. No attempt is made to assign New Zealand taxa to subfamilies, as subfamily concepts differ widely among workers on the Tineidae.

● ***Amphixystis*** Meyrick, 1901, p. 576. Type species *Amphixystis hapsimacha* Meyrick, by original monotypy.

Note. Meyrick (1927b, pp. 315–316) redescribed the genus and species.

***hapsimacha*** Meyrick, 1901, p. 577 (*Amphixystis*)  
Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 333, pl. xxxvi fig. 24 and 25.

● ***Archyala*** Meyrick, 1889b, p. 159. Type species *Archyala paraglypta* Meyrick, by original monotypy.

***culta*** Philpott, 1931, p. 35 (*Archyala*)  
Opho, Dunedin DN, C.E. Clarke; HT ♂ unique, AMNZ.  
Hudson 1939, p. 464, pl. lxi fig. 30.

***lindsayi*** Philpott, 1927a, p. 708 (*Tinea*) new combination  
Mt Grey NC, S. Lindsay; HT ♂ unique, CMNZ.  
Hudson 1928, p. 347, pl. lii fig. 32 (an inaccurate illustration).

***opulenta*** Philpott, 1926a, p. 398 (*Archyala*)  
Upper Maitai NN, E.S. Gourlay; HT ♂ designated by  
Philpott, NZAC.  
Hudson 1928, p. 341, as synonym of *terranea* Butler.  
Philpott (1927e, p. 94, fig. 5) shows genital differences  
between *opulenta* and *terranea*.

***paraglypta*** Meyrick, 1889b, p. 159 (*Archyala*)  
Riccarton Bush, Christchurch MC, E. Meyrick; HT ♂  
unique, BMNH.  
Hudson 1928, p. 341, pl. xxxvii fig. 3.

➤ Tineidae, *Archyala paraglypta*

**tigrina** Philpott, 1930b, p. 15 (*Archyala*). Synonymised by Hudson (1939, p. 464).  
Waikaraka Valley, Whangarei ND, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 464, not figured.

**pentazyga** Meyrick, 1915a, p. 204 (*Archyala*)  
Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 341, pl. xxxvii fig. 4.

**terranea** Butler, 1879a, p. 510 (*Scardia*)  
“Otago” [?Dunedin DN], F.W. Hutton; HT ♀ unique, BMNH.  
Hudson 1928, p. 341, pl. xxxvii fig. 5.

Also 1 undescribed species (NZAC).

● **Astrogenes** Meyrick, 1921, p. 335. Type species *Astrogenes chrysograpta* Meyrick, by original monotypy.

**chrysograpta** Meyrick, 1921, p. 335 (*Astrogenes*)  
Mt Arthur NN, 4,200 ft, Stella Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 347, pl. xxxiv fig. 14, clearly a topotypic specimen – North Island specimens lack the silver spots on the apical third of the forewing costa, i.e., they agree with Whangarei ND *insignita* Philpott.

**insignita** Philpott, 1930b, p. 16 (*Astrogenes*)  
[Waikaraka Valley, Whangarei ND], C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 467, as synonym of *chrysograpta*.  
Note. The specimen labelled “Holotype ♂” by Philpott agrees in wingspan with the published description. The 4 PTs are all from Woodhaugh DN, as published. Philpott may have assumed that all 5 specimens were from Woodhaugh. I have not disturbed the labelling so that a name is available for the North Island population should that be necessary.

● **Bascantis** Meyrick, 1914a, pp. 114–115. Type species *Bascantis sirenica* Meyrick, by original monotypy.

**sirenica** Meyrick, 1914a, p. 115 (*Bascantis*)  
Kaeo ND, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 341, pl. xxxvii fig. 23; 1950, pp. 112–113, pl. vii fig. 2.  
Note. Although Meyrick records 2 localities, he saw only 1 specimen.

● **Crypsitricha** Meyrick, 1915b, p. 235. Type species *Endophthora mesotypa* Meyrick, by original designation.

**agriopa** Meyrick, 1888e, p. 95 (*Endophthora*)  
Wellington WN, E. Meyrick; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 339, not figured.

**generosa** Philpott, 1926a, p. 398 (*Crypsitricha*)  
Manapouri [Hunter Mountains] FD, S. Lindsay; HT ♀ unique, CMNZ.  
Hudson 1928, p. 340, pl. lii fig. 15.

**mesotypa** Meyrick, 1888e, pp. 94–95 (*Endophthora*)  
Auckland AK, E. Meyrick; LT ♀ here designated, labelled “Auckland New Zealand 16.12.85”, “Crypsitricha mesotypa Meyr. 8/19 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 340, pl. xxxvii fig. 6 (pattern agrees well with LT).

**pharotoma** Meyrick, 1888e, p. 94 (*Endophthora*)  
Whangarei [Heads] ND, E. Meyrick; LT ♂ here designated, labelled “Whangarei New Zealand 19.12.85”, “Crypsitricha pharotoma Meyr. 7/10 E. Meyrick det. in Meyrick Coll.”, abdomen in capsule, BMNH.  
Hudson 1928, p. 339, pl. xi fig. 15 (the forewings are drawn too narrow).

Note. There are 6 PJ.Ts, nos. 1–4/10 and 6/10 from Christchurch, no. 5/10 from Palmerston North, also in BMNH.

**roseata** Meyrick, 1913a, p. 28 (*Endophthora*)  
Wadestown, Wellington WN, G.V. Hudson; HT ♀ unique, abdomen missing, BMNH.  
Hudson 1928, p. 340, pl. xxxvii fig. 22.

**stereota** Meyrick, 1914a, p. 114 (*Endophthora*)  
Auckland AK, G.V. Hudson; LT ♂ here designated, labelled “Auckland New Zealand GVH 2.1.13”, “Crypsitricha stereota Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 339, pl. xxxvii fig. 7.

Also 1 undescribed species (NZAC).

● **Dryadaula** Meyrick, 1893, p. 559. Type species *D. glycinopa* Meyrick, by original monotypy; Australia (Blackheath, N.S.W.).

**castanea** Philpott, 1915, p. 201 (*Dryadaula*)  
Bluff SL, A. Philpott; HT ♂ designated without gender by Philpott, abdomen missing, NZAC.  
Hudson 1928, p. 338, pl. xxxvi fig. 26.

**myrrhina** Meyrick, 1905, p. 243 (*Dryadaula*)  
[Wellington] “New Zealand”, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 337, pl. xxxvii fig. 19.

**pactolia** Meyrick, 1901, p. 577 (*Dryadaula*)  
Nelson NN, E. Meyrick; LT ♂ here designated, labelled “Nelson New Zealand 12.1.86”, “Dryadaula pactolia

Meyr. 4/4 E. Meyrick. det. in Meyrick Coll.", BMNH. Hudson 1928, p. 338, pl. li fig. 13.

Note. Recorded as "Esch[atotypal] sp.?" in Meyrick's Diary of Captures for 12 January 1886. Adventive to Great Britain (Meyrick 1916a, pp. 17–18; Morrison 1968 [biology]) and Denmark (Karsholt & Nielsen 1976a).

● **Endophthora** Meyrick, 1888e, p. 93. Type species *Endophthora omogramma* Meyrick, by subsequent designation (Meyrick 1915, p. 235).

**omogramma** Meyrick, 1888e, p. 94 (*Endophthora*) Mt Arthur [descent] NN, E. Meyrick; LT ♀ here designated, labelled "Mt Arthur New Zealand 19/1/86", "Endophthora omogramma Meyr. 2/4 E. Meyrick det. in Meyrick Coll.", BMNH; PLT ♀ from [Waitakere Range] "Auckland, 17.12.85" lacks the abdomen. Hudson 1928, p. 339, pl. xxxvi fig. 23.

**pallacopis** Meyrick, 1918a, p. 134 (*Endophthora*) Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 339, pl. xl fig. 1.

**rubiginella** Hudson, 1939, p. 464 (*Endophthora*) Waiho River WD, G.V. Hudson; HT ♂ unique, labelled by Hudson "1270a", NMNZ. Hudson 1938, p. 464, pl. lvii fig. 13.

**tylogramma** Meyrick, 1924a, p. 206 (*Endophthora*) Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 339, pl. l fig. 15.

● **Erechthias** Meyrick, 1880b, pp. 261–262. Type species *Erechthias charadrota* Meyrick, by subsequent designation (Meyrick 1915b, p. 233).

Note. For extralimital synonymies concerning Pacific taxa, see Robinson (1983).

**Hectacma** Meyrick, 1915b, pp. 233–234. Type species *Erechthias chasmatis* Meyrick, by original designation. **New synonymy.**

**acrodina** Meyrick, 1912c, pp. 122–123 (*Ereunetis*) Wellington WN, G.V. Hudson; HT ♀ unique, BMNH. Hudson 1928, pp. 334–335, pl. xxxvi fig. 19.

**charadrota** Meyrick, 1880b, p. 268 (*Erechthias*) [hills above Port Lyttelton], Christchurch MC, E. Meyrick; LT ♀ here designated, labelled "Christchurch New Zealand 8.1.80", "Erechthias charadrota Meyr. 9/11 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, p. 335, pl. xxxvi fig. 16, from a North Island specimen.

**melanotricha** Meyrick, 1888e, p. 93 (*Erechthias*). **New synonymy.**

[Mt Manaia, Whangarei ND], E. Meyrick; LT ♂ here designated, labelled "Auckland New Zealand 21.12.85", "Erechthias melanotricha Meyr. 2/2 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, p. 335, not figured as such.

Note. Meyrick's Diary of Captures indicates that Meyrick's label on the LT is in error. Specimens from central New Zealand in NZAC are intermediate in colour pattern characters, hence the proposed synonymy.

**chasmatis** Meyrick, 1880b, p. 264 (*Erechthias*) Wellington [Botanic Gardens] WN, E. Meyrick; LT ♂ here designated, labelled "Wellington, New Zealand 31.12.79", "Hectacma chasmatis Meyr. 2/11 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, p. 336, pl. xxxvi fig. 17.

**chionodira** Meyrick, 1880b, pp. 268–269 (*Erechthias*) Auckland [Domain] AK, E. Meyrick; LT ♂ here designated, labelled "Auckland New Zealand 20.1.80", "Hectacma chionodira Meyr. 8/12 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, p. 336, pl. xxxvi fig. 28.

**crypsimima** Meyrick, 1920a, p. 31 (*Hectacma*) Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, pp. 336–337, pl. xlvi fig. 11.

**decoranda** Meyrick, 1925a, p. 274 (*Hectacma*) Manukau, Chatham Islands, S. Lindsay; HT ♂ labelled "Chatham Islands Dec. 1923 Coll. C. Lindsay", "Hectacma decoranda Meyr. TYPE E. Meyrick det. 1924", CMNZ.

Hudson 1928, p. 337, pl. lii fig. 14 (not a very accurate portrayal).

Note. Meyrick returned 3 of the 4 specimens constituting the type series; one, the HT, bears the word "TYPE" in Meyrick's handwriting and is in CMNZ. BMNH has the specimen Meyrick retained (♂ genitalia on slide no. 3779), and this is paratypic, as are the two CMNZ specimens collected on 15 December 1923 and 4–21 January 1924. The presence of an uncus and gnathos may exclude this species from true *Erechthias*.

**exospila** Meyrick, 1901, p. 577 (*Ereunetis*) Whangarei [Heads] ND, E. Meyrick; HT ♂ unique, BMNH. Hudson 1928, p. 335, pl. xxxvi fig. 22.

**externella** Walker, 1864b, p. 841 (*Glyphipteryx*) [Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 334, pl. xxxvi fig. 13 and 14. Note. Walker's record of the collector as Col. Bolton is regarded here as erroneous; the original circular label reads "60-73" and "Auckland, N. Zeal."

**bisignella** Walker, 1864b, p. 1007 (*Tinea*) new synonymy

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH. Not mentioned by Hudson or Meyrick.

Note. Walker made the same error as for *externella*.

**monastræ** Meyrick, 1891, p. 100 (*Decadarchis*). Synonymised by Meyrick (1919, p. 353). Wellington WN, G.V. Hudson; HT ♀ unique, abdomen missing, BMNH.

➤ Tineidae, *Erechthias externella*

**erebitis** Meyrick, 1892, p. 220 (*Erechthias*).  
Synonymised by Meyrick (1915, p. 223).  
Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 334, as synonym.

**fulguritella** Walker, 1863c, p. 548 (*Cerostoma*)  
[Nelson NN, T.R. Oxley]; HT ♀ unique, abdomen missing, BMNH.  
Hudson 1928, p. 336, pl. xxxvi fig. 20; 1950, p. 112.

**hemiclistra** Meyrick, 1911b, pp. 77–78 (*Decadarchis*)  
Makara WN, R.M. Sunley; LT ♀ here designated, labelled  
“Makara New Zealand RMS bred 11.09”, “Erechthias  
hemiclistra Meyr. 7/8 E. Meyrick det. in Meyrick  
Coll.”, BMNH.  
Hudson 1928, p. 335, pl. iii fig. 36 and 37, pl. xxxvi fig.  
21.

**indicans** Meyrick, 1923, p. 168 (*Erechthias*)  
[Karori], Wellington WN, G.V. Hudson; HT ♀ unique,  
BMNH.  
Hudson 1928, p. 335, pl. xliv fig. 33.

**lychnopa** Meyrick, 1927a, p. 702 (*Erechthias*)  
[Sinclair Head], Wellington WN, G.V. Hudson; HT ♂  
unique, BMNH.  
Hudson 1928, p. 334, pl. xl fig. 8.

**macrozyga** Meyrick, 1916b, p. 419 (*Erechthias*)  
Tisbury, Invercargill SL, A. Philpott; HT ♂ unique,  
abdomen missing, BMNH.  
Hudson 1928, p. 336, pl. xlvi fig. 13.

**stilbella** Doubleday, 1843, p. 289 (*Argyrotesia*)  
[Auckland AK], A. Sinclair; type material not in BMNH,  
but identity not in doubt; Meyrick specimen 3/14  
topotypic, BMNH.  
Hudson 1928, p. 336, pl. xxxvi fig. 18.

**terminella** Walker, 1863c, p. 548 (*Cerostoma*)  
[Auckland AK], D. Bolton; LT ♀ here designated, labelled  
“Cerostoma terminella Wkr Ct. Lep. Het. B.M. 28  
p.548 (1863) Type ♂”, “New Zeal. 54.4” (lavender,  
circular label), BMNH. There are 8 PLTs (Walker's  
specimens b–i).  
Hudson 1928, p. 335, pl. xxxiii fig. 21.

**? subpavonella** Walker, 1864b, p. 898 (*Elachista*).  
Synonymised by Meyrick (1915, p. 233).  
[Auckland AK], A. Sinclair; LT lost, according to H. Dur-  
rant, BMNH; Walker's PLT – [Auckland AK], D. Bolton  
– is also missing.  
Note. Meyrick's synonymy was based on his interpre-  
tation of Walker's description, and on a ♂ (now lacking  
the abdomen) labelled “Auckland [Domain] 12.1.80” by  
Meyrick (“*Erechthias terminella* Wlk 2/6 det. E. Meyrick  
in Meyrick Coll.”) and recorded as *subpavonella* in his  
Diary of Captures.

Also 2 undescribed species (NZAC).

● **Eschatotypa** Meyrick, 1880b, p. 256. Type species  
*Eschatotypa melichrysa* Meyrick, by original  
monotypy.

**derogatella** Walker, 1863c, p. 485 (?*Tinea*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1928, p. 338, pl. xxxvii fig. 1.

**halosparta** Meyrick, 1919, p. 354 (*Archyala*) new  
combination  
Wainuiomata WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 341, pl. xlvii fig. 9.  
Note. On external genitalia, colour pattern, and labial  
palpus *halosparta* is placed in *Eschatotypa*.

**melichrysa** Meyrick, 1880b, p. 256 (*Eschatotypa*)  
Dunedin DN, E. Meyrick; LT ♂ here designated, labelled  
“Dunedin New Zealand 4.1.80”, “*Eschatotypa meli-  
chrysa* Meyr. 2/8 E. Meyrick det. in Meyrick Coll.”,  
BMNH.  
Hudson 1928, p. 338, pl. xxxvii fig. 2.

● **Eugenaea** Meyrick, 1915b, p. 232. Type species  
*Decadarchis laquearia* Meyrick, by original  
monotypy.

**laquearia** Meyrick, 1914a, p. 113 (*Decadarchis*)  
Kao ND, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, pp. 333–334, pl. xxxvi fig. 15.

● **Habrophila** Meyrick, 1889b, p. 161. Type species  
*Habrophila compseuta* Meyrick, by original  
monotypy.

**compseuta** Meyrick, 1889b, pp. 161–162 (*Hab-  
rophila*)  
[plateau, 3,700–4,000 ft], Mt Arthur NN, E. Meyrick; HT  
♀ unique, BMNH.  
Hudson 1928, p. 340, pl. xl fig. 19.

● **Lysiphragma** Meyrick, 1888e, p. 104. Type spe-  
cies *Lysiphragma mixochlora* Meyrick, by subse-  
quent designation (Meyrick 1915b, p. 239).

**epixyla** Meyrick, 1888e, pp. 105–106 (*Lysiphragma*)  
[Lower Hutt] “Wellington” WN, E. Meyrick; LT ♂ here  
designated, labelled “Wellington New Zealand  
25.1.86”, “*Lysiphragma epixyla* Meyr. 6/7 E. Meyrick  
det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 349, pl. xxxix fig. 10.

**howesii** Quail, 1901, p. 154 (*Lysiphragma*)  
Invercargill SL, W.G. Howes [“in rotten wood, *Plagian-  
thus betulinus*”]; not located.  
Hudson 1928, p. 349, pl. xxxix fig. 8.

Note. The ♂ labelled "629c" ("Invercargill, Oct. 1900, G. Howes" in Hudson Register, NMNZ) postdates Quail's paper, delivered 4 July 1900, mentioning adults reared by Howes on and after 21 December 1899.

**mixochlora** Meyrick, 1888e, p. 105 (*Lysiphragma*) [Waitakere Range], Auckland AK, E. Meyrick; LT ♂ here designated, labelled "Auckland New Zealand 17.12.85", "Lysiphragma mixochlora Meyr. 2/7 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, pp. 348–349, pl. xxxix fig. 9.

• **Monopis** Hübner, 1825, p. 401. Type species *Tinea rusticella* Hübner, 1796.

**argillacea** Meyrick, 1893, p. 528 (*Blabophanes*) "Melbourne and Warragul, Victoria; Adelaide, South Australia; four specimens in September and December" (Meyrick 1893, p. 528). ST ♂ labelled "Adelaide S. Australia OBL /91", "Monopis argillacea Meyr. 1/5 E. Meyrick det. in Meyrick Coll.", "Syntype", BMNH. New Zealand: various localities AK, after 1970, in birds' nests.

Not mentioned by Hudson.

**crocicapitella** Clemens, 1859, pp. 257 (key) and 258 (description) (*Tinea*) Cosmopolitan.

New Zealand: widespread, around human habitation.

**ferruginella** in the sence of Meyrick (1888e, p. 97). Synonymised by Meyrick (1915, p. 238). Hudson 1928, p. 343, pl. xxxix fig. 22.

**dimorphella** Dugdale, 1971b, p. 150 (*Monopis*) Long Island [Big South Cape Island] SI, J.S. Dugdale; HT ♂ designated by Dugdale, NZAC.

**ethelella** Newman, 1856, p. 288 (*Tinea*) ["Forest Creek, Barker's Creek, and Campbells Creek ... Mt Alexander Range ... about eighty miles from Melbourne" Vict., T.R. Oxley; BMNH.] New Zealand: widespread, largely rural, ?introduced. Hudson 1928, p. 343, pl. xxxix fig. 7.

**rectella** Walker, 1863, p. 482 (*Tinea*). Synonymised by Meyrick (1888, p. 97). [Tasmania; BMNH.] Hudson 1928, p. 343, as synonym.

**namuella** Felder & Rogenhofer, 1875 (*Blabophanes*). Synonymised by Meyrick (1888, p. 97). [Nelson NN], T.R. Oxley (1864); HT ♀ unique, BMNH. Hudson 1928, p. 343, as synonym.

**ornithias** Meyrick, 1888e, p. 97 (*Blabophanes*) Christchurch MC, R.W. Fereday; LT ♂ here designated, labelled "Christchurch New Zealand RWF /82", "Monopis ornithias Meyr. 1/3 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, p. 343, pl. xxxvii fig. 25.

**typhlopa** Meyrick, 1925a, p. 274 (*Monopis*) Mangere Island, Chatham Islands, S. Lindsay; LT ♀ (identification label in Meyrick's writing), CMNZ. Hudson 1928, p. 343, not figured.

Note. Meyrick's labelling gives no indication as to which of the two specimens is the HT, so the series is basically syntypic. Lindsay placed the "Type" label on a specimen, so it is the LT.

• **Nemapogon** Schrank, 1802, p. 167. Type species *Tinea granella* Linnaeus, by original monotypy.

**granella** Linnaeus, 1758, p. 357 (*Tinea*) Cosmopolitan.

New Zealand: adventive, widespread around human habitation. Hudson 1939, pp. 466–467, pl. lxi fig. 23, as *Tinea granella*.

• **Niditinea** Petersen, 1957, p. 134. Type species *Tinea fuscipunctella* Haworth, by original designation.

**fuscella** Linnaeus, 1758, p. 539 (*Phalaena (Tinea)*) Europe.

New Zealand: adventive, widespread, in birds' nests. Hudson 1928, p. 346, pl. xi fig. 4, as *Tinea fuscipunctella*.

**fuscipunctella** Haworth, 1828, p. 562 (*Tinea*). Synonymised by Robinson & Nielson (1983, p. 238).

• **Opogona** Zeller, 1853, p. 507. Type species *Opogona dimidiatella* Zeller, 1853; Java.

**Hieroxestis** Meyrick, 1893, p. 567. Type species *Hieroxestis omoscopa* Meyrick, by original monotypy. Synonymised by Meyrick (1927, p. 315.)

**comptella** Walker, 1864b, p. 1007 (*Tinea*) [Hobart] Tasmania, Smith; HT ♀ unique, BMNH. New Zealand: adventive (Meyrick 1911b, p. 69). Hudson 1928, p. 333, pl. xxxiv fig. 7.

**omoscopa** Meyrick, 1893, p. 567 (*Hieroxestis*) [Sydney N.S.W., E. Meyrick; LT ♂ designated by Davis (1978, p. 23), BMNH.]

New Zealand: first recorded from Kaeo ND and Thames CL areas (Meyrick 1914a, p. 113), now widespread, ND south to 42°S; an isolated population is in Riccarton Bush MC (C. Muir, pers. comm., 1985). Hudson 1928, p. 332, pl. xxxvi fig. 11 and 12.

Note. Davis (1978, pp. 13 and 16–23) gives full generic and specific synonymies and world distribution.

• **Petasactis** Meyrick, 1915b, p. 234. Type species *Ereunetis technica*, by monotypy (Meyrick 1915, p. 234).

» Tineidae, *Petasactis*

**technica** Meyrick, 1888e, p. 92 (*Ereunetis*)  
Whangarei [Heads] ND, E. Meyrick; HT ♀ unique,  
BMNH.  
Hudson 1928, p. 337, not figured.

● (**Pringleophaga** Enderlein, 1905. Type species  
*Pringleophaga kerguelensis* Enderlein, 1905, by  
original monotypy.

**Antipodesma** Salmon & Bradley, 1956, p. 64.  
Type species *Antipodesma turbotti* Salmon & Bradley,  
by original designation. Synonymised by Vari  
(1971, p. 349).

Note. In the absence of precise dating, Vari's publication is regarded as dated "(31 December) 1971". Dugdale's synonymy below, under *Proterodesma*, was published on 10 November 1971. It is probable that neither synonymy reflects the true situation.)

● **Proterodesma** Meyrick, 1909b, p. 74. Type species *Proterodesma byrsopola* Meyrick, by original monotypy.

**Antipodesma** Salmon & Bradley, 1956, p. 64.  
Type species *Antipodesma turbotti* Salmon & Bradley,  
by original designation. Synonymised by Dugdale  
(1971b, p. 152).

**byrsopola** Meyrick, 1909b, p. 74 (*Proterodesma*)  
[Carnley Harbour, Auckland Islands], G.V. Hudson; LT  
♀ here designated, labelled "Kermadec Is GVH '08",  
"Proterodesma byrsopola" Meyr. 3/3 E. Meyrick det.  
in Meyrick Coll.", BMNH.

Hudson 1928, p. 348, pl. xxxvii fig. 17 and 18.

**mysticopa** Meyrick, 1914a, p. 115 (*Tinea*).  
Synonymised by Dugdale (1971b, p. 153).  
Invercargill SL, A. Philpott; LT ♂ here designated, labelled  
"Invercargill New Zealand AP 0.10.10", "Tinea mysti-  
copa" Meyr. 2/3 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 348, pl. xl fig. 7.

**chathamica** Dugdale, 1971a, p. 62 (*Proterodesma*)  
Chatham Island, J.S. Dugdale; HT ♂ designated by Dugdale, NZAC.

**turbotti** Salmon & Bradley, 1956, p. 65 (*Anti-  
podesma*)

Ringdove Bay, Antipodes Island, E.G. Turrott; HT ♂  
designated by Salmon & Bradley, AMNZ.

● **Prothinodes** Meyrick, 1914a, p. 116. Type species  
*Prothinodes lutata* Meyrick, by original designation.

**grammocosma** Meyrick, 1888e, pp. 98–99 (*Tinea*)  
Nelson NN, E. Meyrick; LT ♂ here designated, labelled  
"Nelson New Zealand 22.1.86", "Prothinodes gram-

mocosma" Meyr. 1/3 E. Meyrick det. in Meyrick Coll.",  
BMNH.

Hudson 1928, p. 347, pl. xxxix fig. 5.

**lutata** Meyrick, 1914a, pp. 116–117 (*Prothinodes*)  
Kao ND, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, pp. 347–348, pl. xxxix fig. 4.

● **Sagephora** Meyrick, 1888e, pp. 95–96. Type species  
*Sagephora phortegella* Meyrick, by original monotypy.

**exsanguis** Philpott, 1918, p. 131 (*Sagephora*)  
Bluff SL, A. Philpott; HT ♂ designated by Philpott,  
NZAC.  
Hudson 1928, p. 342, pl. xl fig. 18.

**felix** Meyrick, 1914a, p. 114 (*Sagephora*)  
Kao ND, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 342, pl. xxxvii fig. 14.

**jocularis** Philpott, 1926a, p. 398 (*Sagephora*)  
Tisbury, Invercargill SL, A. Philpott; HT ♂ designated  
by Philpott, NZAC.  
Hudson 1928, p. 342, pl. lii fig. 33.

**phortegella** Meyrick, 1888e, p. 96 (*Sagephora*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂  
here designated, labelled "Christchurch New Zealand  
29.8.82", "Sagephora phortegella" Meyr. 7/11 E. Meyr-  
ick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 342, pl. xxxvii fig. 20 and 21.

**steropastis** Meyrick, 1891, p. 100 (*Sagephora*)  
Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 342, pl. xxxvii fig. 15.

**subcarinata** Meyrick, 1931a, pp. 96–97 (*Sagephora*)  
[Gollan's Valley], Wellington WN, G.V. Hudson; HT ♂  
unique, BMNH.  
Hudson 1939, p. 465, pl. lviii fig. 11.

● **Lindera** Blanchard, 1852, p. 105. Type species  
*Lindera tessellatella* Blanchard, by original mono-  
typy; Chile.

Note. The synonymy with *Setomorpha* Zeller, 1852,  
p. 94 (type species *S. rutella* Zeller) by Gozmany  
& Vari (1973, p. 82) is not recognised by most  
workers.

**tessellatella** Blanchard, 1852, p. 106 (*Lindera*)  
Cosmopolitan.  
New Zealand: adventive, widespread, around human  
habitation.  
Hudson 1928, p. 350, pl. I fig. 24.  
Note. Philpott (1924a, p. 214) gives early records.

● **Tephrosaria** Meyrick, 1915b, p. 234. Type species *Erechthias cimmeria* Meyrick, by monotypy.

**cimmeria** Meyrick, 1914a, pp. 113–114 (*Erechthias*) Waitakere AK, G.V. Hudson; LT ♂ here designated, labelled “Waitakere New Zealand GVH 1.13”, “Tephrosaria cimmeria Meyr. 2/2 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 337, pl. xxxvi fig. 27.

● **Thallostoma** Meyrick, 1913a, pp. 28–29. Type species *Thallostoma eurygrapha* Meyrick, by monotypy.

**eurygrapha** Meyrick, 1913a, p. 29 (*Thallostoma*) Wadestown, Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “Wadestown New Zealand GVH 11.96”, “Thallostoma eurygrapha Meyr. 2/2 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 342, pl. xxxix fig. 3.

● **Tinea** Linnaeus, 1758, p. 534. Type species *Tinea pellionella* Linnaeus, by subsequent designation (ICZN 1957, *Opin. decl. Int. Comm. Zool. Nom. I*, p. 254).

**dubiella** Stainton, 1859, p. 183 (*Tinea*) See Robinson (1979, pp. 88 and 90–91) for full synonymy and details of biology.  
New Zealand: adventive, widespread, around human habitation.

**turicensis** Müller-Ratz, 1920, p. 348 (*Tinea*).  
Synonymised by Robinson (1979, p. 88).

**pallescentella** Stainton, 1851, p. 2 (*Tinea*) See Petersen (1969, pp. 374–375) for synonymy and biology.  
New Zealand: adventive, widespread, around human habitation.  
Not mentioned by Hudson.

**pellionella** Linnaeus, 1758, p. 536 (*Phalaena (Tinea)*) See Robinson (1979, pp. 72–74 and 76–77) for full synonymy and details of biology.  
New Zealand: adventive, widespread, around human habitation.  
Hudson 1939, p. 467, pl. lxi fig. 22.

● **Tinea** in the sense of Meyrick (1915, p. 238)

**accusatrix** Meyrick, 1916b, p. 419 (*Tinea*) [Kaitoke], Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “Wellington New Zealand GVH .15”, “Tinea accusatrix Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 345, pl. xlvi fig. 2.

**aetherea** Clarke, 1926, p. 421 (*Tinea*) Arthur's Pass NC/WD, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.  
Hudson 1928, p. 346, pl. lii fig. 13.

**argodelta** Meyrick, 1915a, p. 204 (*Tinea*) Bluff SL, A. Philpott; HT ♂ unique, BMNH.  
Hudson 1928, p. 346, pl. xxxvii fig. 9

**astraea** Meyrick, 1911b, p. 68 (*Tinea*) [West Plains (Meyrick 1911, p. 68) or Otatara (Hudson 1928, p. 345)], Invercargill SL, A. Philpott; HT ♀ unique, BMNH.

Hudson 1928, p. 345, but pl. xxxvii fig. 13 is not *T. astraea*.

**cymodoce** Meyrick, 1924a, p. 206 (*Tinea*). New synonymy.

Mt Arthur NN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 345, pl. xlvi fig. 13.

**erata** Philpott, 1930b, p. 15 (*Tinea*). Synonymised by Hudson (1939, p. 465).  
Tongariro National Park TO, C.E. Clarke; HT ♀ designated by Philpott, AMNZ.  
Hudson 1939, p. 465, pl. lii fig. 22.

Note. The HT locality label reads “Tongariro, 16.1.30, A. Philpott”.

**atmogramma** Meyrick, 1927b, p. 316 (*Tinea*) Arthur's Pass NC/WD, G.V. Hudson; LT ♀ here designated, labelled “Arthurs Pass New Zealand GVH 3000' 1.27”, “Tinea atmogramma Meyr. 1/1 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1939, p. 466, pl. li fig. 17 and 18.

**belonota** Meyrick, 1888e, p. 99 (*Tinea*) Palmerston North WI/WN, E. Meyrick; HT unique, not found in BMNH (label should read “Palmerston New Zealand 4/3/83” in Meyrick's writing; based on the entry in Meyrick's Diary of Captures).  
Hudson 1928, p. 347; 1939, p. 466, pl. lx fig. 12 (specimen from a series checked by Meyrick, from Pohangina WI, close to Palmerston North).

**zephyrina** Clarke, 1926, p. 419 (*Gymnobathra*).  
Synonymised by Philpott (1931, p. 35).  
Whangarei ND, C.E. Clarke; HT ♂ unique, AMNZ.  
Not mentioned by Hudson.

**conferta** Meyrick, 1914a, pp. 115–116 (*Tinea*) Wellington WN, E. Meyrick; LT ♂ here designated, labelled “Wellington New Zealand 1.1.80”, “Tinea conferta Meyr. 4/15 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 346, pl. xxxix fig. 13.

**conspecta** Philpott, 1931, p. 35 (*Tinea*) The Domain, Auckland AK, A. Philpott; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 466, not figured.

**dicharacta** (not of Meyrick, 1893, p. 536) Meyrick, 1911b, p. 78 (*Tinea*)

➤ Tineidae, *Tinea dicharacta*

[True *T. dicharacta* from Sydney, N.S.W., E. Meyrick; HT ♂ unique, BMNH.]  
Hudson 1928, p. 346, pl. xxxvii fig. 8.  
Note. The New Zealand specimen differs in forewing pattern and in antennal characters.

***dividua*** Philpott, 1928a, p. 370 (*Tinea*)  
Flora River NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 466, pl. lx fig. 9.

***fagicola*** Meyrick, 1921, p. 336 (*Tinea*)  
[Day's Bay], Wellington, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 2.19.", "Tinea fagicola Meyr. 1/7 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 345, pl. xlvi fig. 15.

***furcillata*** Philpott, 1930b, p. 15 (*Tinea*)  
Anderson's Bay, Dunedin DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 465, pl. lxi fig. 20.  
Note. The HT lacks the left forewing.

***margaritis*** Meyrick, 1914a, p. 116 (*Tinea*)  
Tisbury, Invercargill SL, A. Philpott; LT ♂ here designated, labelled "Tisbury New Zealand AP 19.12.11", "Tinea margaritis Meyr. 1/5 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 344, pl. xxxvii fig. 26.

***mochlota*** Meyrick, 1888e, p. 100 (*Tinea*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 13/1/83", "Tinea mochlota Meyr. 1/5 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 346, pl. xxxix fig. 4.

***munita*** Meyrick, 1932, p. 24 (*Tinea*)  
[Wilton's Bush], Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1939, p. 466, pl. lx fig. 13.

***sphenocosma*** Meyrick, 1919, pp. 353–354 (*Tinea*)  
Wellington WN, G.V. Hudson; HT ♀ unique, not in BMNH.  
Hudson 1939, p. 465, pl. lxi fig. 24.

***texta*** Meyrick, 1931a, p. 97 (*Tinea*)  
Whangarei ND, S.C. Patterson; HT ♀ unique, not in BMNH.  
Hudson 1939, p. 465, pl. lxi fig. 21 (specimen from Wellington WN).

Also 3 undescribed species (NZAC), one of them adventive.

● ***Tineola*** Herrich-Schaeffer, 1853, p. 23. Type species *Tineola bisselliella* Hummel, by original monotypy.

***bisselliella*** Hummel, 1823, pp. 6–13 (*Tinea*)  
Cosmopolitan.  
New Zealand: adventive, widespread, around human habitation.  
Hudson 1928, p. 344, pl. xlvi fig. 9, as *biselliella*.

● ***Trichophaga*** Ragonot, 1894, p. 123. Type species *Trichophaga coprobiella* Ragonot, by original designation.


***tapetzella*** Linnaeus, 1758, p. 536 (*Tinea*)  
Cosmopolitan.  
New Zealand: adventive, widespread, around human habitation.  
Hudson 1928, p. 343, pl. xxxix fig. 12, as *tapetiella*.

● ***Trithamnora*** Meyrick, 1913a, p. 29. Type species *Trithamnora improba* Meyrick, by original monotypy.

***certella*** Walker, 1863c, p. 484 (*Tinea*)  
[Auckland AK], ?D. Bolton; HT ♀ unique, BMNH.  
Hudson 1928, p. 348, pl. xlvi fig. 15 and 16.  
***improba*** Meyrick, 1913a, p. 29 (*Trithamnora*).  
Synonymised by Meyrick (1915, p. 239).  
Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 12.11", "Trithamnora certella Meyr. 1/14 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 348, as synonym.

—◎—

**Family PSYCHIDAE**  
**Subfamily PSYCHINAE**  
(in the sense of Kozhanchikov 1956)



Psychidae: Psychinae  
(144) *Orophora unicolor* (Butler), ♂

● **Liothula** Fereday, 1878b, p. 260. Type species *Liothula omnivora* Fereday, by original monotypy.

**omnivora** Fereday, 1878b, p. 260 (*Liothula*).  
Christchurch MC, R.W. Fereday; 10 ♂ STs, CMNZ.  
Hudson 1928, pp. 212–214, pl. xliv fig. 14, as *Oiceticus omnivorus*.


● **Orophora** Fereday, 1878b, p. 261. Type species *Orophora toumatou* Fereday, by original designation.

**unicolor** Butler, 1877, p. 381 (*Psyche*)  
[?Castle Hill MC], J.D. Enys; HT ♂ unique, BMNH.  
Hudson 1928, p. 214, pl. xliv fig. 9.

**toumatou** Fereday, 1878b, p. 262 (*Orophora*).  
Synonymised by Meyrick (1890, p. 212).  
[Rakaia and Waimakariri riverbeds] MC, R.W. Fereday;  
3 ♂ STs, CMNZ.  
Hudson 1928, p. 214, as synonym.  
Note. A single specimen labelled “Meserala seelandica Gn” from “Nouvelle Zélande”, collected by R.W. Fereday and representing a manuscript name, is in BMNH. It agrees well with *O. unicolor*.



**Subfamily PSYCHEODINAE**  
(in the sense of Kozhanchikov 1956)



Psychidae: Psycheodinae  
(145) *Mallobathra perisseuta* Meyrick

● **Cebysa** Walker, 1854, p. 486. Type species *Cebysa leucotelus* Walker, by original monotypy.

**leucotelus** Walker, 1854, p. 486 (*Cebysa*)  
“Sydney. From the collection of the Paris Museum”; HT  
♀ unique, BMNH.  
Australian.  
New Zealand: adventive, Mt Albert AK.

● **Grypotheca** Dugdale, 1987b, p. 107. Type species *Grypotheca pertinax* Dugdale, by original designation.

**horningae** Dugdale, 1987b, p. 110 (*Grypotheca*)  
The Snares islands, C.J. Horning; HT ♂ designated by  
Dugdale, NZAC.

**gen. et sp. indet.** Dugdale 1971, pp. 143–144.

**pertinax** Dugdale, 1987b, p. 109 (*Grypotheca*)  
Riccarton Bush MC, J.S. Dugdale; HT ♂ designated by  
Dugdale, NZAC.

**triangularis** Philpott, 1930b, p. 16 (*Talaeporia*) new  
combination  
Hump Ridge FD, C.E. Clarke; HT ♂ designated by Phil-  
pott, AMNZ.  
Hudson 1939, pp. 467–468, pl. lxi fig. 5.

Also 3 undescribed species (NZAC; B. Patrick Col-  
lection, Dunedin).

● **Mallobathra** Meyrick, 1888e, p. 102. Type spe-  
cies *Mallobathra crataea* Meyrick, by subsequent  
designation (Meyrick 1915b, p. 240).

**abyssina** Clarke, 1934, p. 15 (*Sabatinca*) new  
combination  
Franz Josef WD, C.E. Clarke; HT ♂ unique, AMNZ.  
Hudson 1939, p. 471, not figured.

» Tineidae, *Mallobathra*

---

**angusta** Philpott, 1928g, p. 498 (*Mallobathra*)  
Flora River [Mt Arthur Tableland] NN, A. Philpott; HT ♂ designated by Philpott, unique, NZAC.  
Hudson 1939, p. 469, pl. lxi fig. 4.

**aphrosticha** Meyrick, 1912c, p. 123 (as *Taleporia aphrosticha*; misspelling)  
Hump Ridge FD, A. Philpott; LT ♂ here designated, labelled "Talaeporia aphrosticha Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 350, pl. xxxix fig. 7.

Note. Meyrick (1912, p. 123) mentions a pair. It is possible that the ♀ in NZAC collected at the same time as the HT is a specimen returned by Meyrick. The specimen bears a label in Meyrick's writing "aphrosticha", indicating Meyrick's intended spelling.

**campbellica** Dugdale, 1971b, p. 139 (*Mallobathra*)  
Beeman Camp, Campbell Island, G. Kuschel; HT ♀ unique, NZAC.

**cana** Philpott, 1927d, pp. 89–90 (*Mallobathra*)  
Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 469, pl. lxi fig. 2.

**cataclysmata** Clarke, 1934, p. 15 (*Mallobathra*)  
[bushline] near Harris Saddle FD/OL, C.E. Clarke; HT ♀ unique, AMNZ.  
Hudson 1939, p. 468, as a synonym of *lapidosa* Meyrick.

**crataea** Meyrick, 1888e, p. 102 (*Mallobathra*)  
Mt Arthur NN, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 16/1/86", "Mallobathra crataea Meyr. 4/17 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 352, pl. xl fig. 2 (based on WN specimens).

**fenwicki** Philpott, 1924a, p. 214 (*Mallobathra*)  
Mt Ruapehu TO, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ.  
Hudson 1928, p. 352, as synonym of *crataea* Meyrick.

**homalopa** Meyrick, 1891, pp. 100–101 (*Mallobathra*).  
Wellington WN, G.V. Hudson; HT ♂ unique, not in BMNH.  
Hudson 1928, p. 353, pl. xxxix fig. 12.  
Note. A topotypic specimen collected at Wellington "xii 1890, G.V. Hudson", no. 1387, ex Walsingham Collection, would be suitable if a NT were deemed necessary.

**lapidosa** Meyrick, 1914a, p. 117 (*Mallobathra*)  
Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH bred 1.13",

"Mallobathra lapidosa Meyr. 2/7 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 352, pl. xxxix fig. 15 and 16.

**memotuina** Clarke, 1934, p. 14 (*Mallobathra*)  
Anderson's Bay DN, C.E. Clarke; HT ♂ designated by Clarke, not in AMNZ.  
Hudson 1939, p. 469, pl. lxi fig. 2.  
Note. None of the 3 ♀ (2 without abdomen) can be the holotype, since none were collected in 1928 or 1929.  
All bear the label "Allotype ♀ 1932".

**metrosema** Meyrick, 1888e, p. 103 (*Mallobathra*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 22/9/82", "Mallobathra metrosema Meyr. 1/9 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 352, pl. xlvi fig. 5.

**obscura** Philpott, 1928a, p. 370 (*Mallobathra*)  
West Plains, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 469, not figured.

**perisseuta** Meyrick, 1920a, p. 32 (*Mallobathra*)  
Dunedin DN, [C.E. Clarke] G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 353, pl. xlvi fig. 15.

**petrodoxa** Meyrick, 1923, p. 169 (*Narycia*) new combination  
Otira Gorge WD, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 354, pl. xlvi fig. 26.  
Note. This may prove to be the ♀ of a described species known only from the ♂.

**scoriota** Meyrick, 1909a, p. 16 (*Mallobathra*)  
[?Invercargill SL or ?Dunedin DN, W.G. Howes or A. Philpott]; LT ♂ here designated, labelled "Wellington New Zealand GVH 07", "Mallobathra scoriota Meyr. 1/6 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 353 "I am unable to identify this species", not figured.

**globulosa** Meyrick 1914a, p. 117 (*Mallobathra*).  
Synonymised by Meyrick (1919, p. 354).  
West Plains SL, [A. Philpott]; LT ♂ here designated, labelled "West Plains New Zealand GVH 27.8.08", "Mallobathra scoriota Meyr. 3/6 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 353 "... discovered by Mr Philpott near Invercargill", pl. xxxix fig. 11.

Note. It is possible that Meyrick received a packet of moths collected by Mr George Howes of Dunedin DN. It is also possible that he confused these with ones from Hudson, and with Philpott material (cf. *Proditrix megalynta* Meyrick, Yponomeutidae).

**strigulata** Philpott, 1924a, p. 214 (*Mallobathra*)  
Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 352, as synonym of *crataea*; 1939, p. 468, as species.

**subalpina** Philpott, 1930a, p. 250 (*Mallobathra*)  
Red Lake, Mt Sebastopol MK, S. Lindsay; HT ♂ des-  
ignated by Philpott, CMNZ.  
Hudson 1939, p. 468, pl. ix fig. 11.

**tonnoiri** Philpott, 1927d, p. 90 (*Mallobathra*)  
“Lake Moana” [Lake Brunner] BR, A. Tonnoir; HT ♂  
unique, CMNZ.

Also 1 undescribed species (NZAC).

● **Reductoderces** Salmon & Bradley, 1956, p. 69, in  
the sense of Dugdale (1971b, p. 141). Type species  
*Reductoderces fuscoflava* Salmon & Bradley, by  
original designation.

**araneosa** Meyrick, 1914a, p. 117 (*Mallobathra*)  
Ben Lomond OL, A. Philpott; LT ♂ here designated,  
labelled “Ben Lomond New Zealand AP 25.11.12”,  
“Mallobathra arenosa 1/3 E. Meyrick det. in Meyrick  
Coll.”, BMNH.  
Hudson 1928, p. 351, pl. xxxix fig. 6.

**aucklandica** Dugdale, 1971b, p. 141 (*Reductoderces*)  
Camp Cove, Auckland Island, P.M. Johns; HT ♂ des-  
ignated by Dugdale, NZAC.

**cawthronella** Philpott, 1921, p. 341 (*Taleporia*)  
[Botanical Hill], Nelson NN, A. Philpott; HT ♂ des-  
ignated by Philpott, NZAC.  
Hudson 1928, p. 351, pl. xlvi fig. 9.

**fuscoflava** Salmon & Bradley, 1956, p. 70  
(*Reductoderces*)  
Campbell Island, J.R. Sorensen; HT ♂ designated by  
Salmon & Bradley, NMNZ.

**illustris** Philpott, 1917b, p. 245 (*Mallobathra*)  
Hump Ridge [The Hump] FD, A. Philpott; HT ♂ des-  
ignated by Philpott, NZAC.  
Hudson 1928, p. 352, pl. xlvi fig. 17.

**microphanes** Meyrick, 1888e, p. 103 (*Mallobathra*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; HT ♂  
unique, BMNH.  
Note. Not as described or depicted in Hudson 1928, p.  
351, pl. xxvii fig. 10 and pl. iii fig. 12. Hudson’s treat-  
ment refers to an unnamed species.

**fragilis** Philpott, 1927a, p. 708 (*Mallobathra*). New  
synonymy.

Riccarton Bush, Christchurch MC, S. Lindsay; HT ♂  
designated by Philpott, right forewing missing, CMNZ.  
HT *fragilis* agrees well in structural and pattern charac-  
ters with HT *microphanes*.

Note. Not as described or depicted in Hudson 1928, p.  
352, pl. lii fig. 6, as species. Philpott’s PT material from  
Riccarton Bush includes three species; Hudson’s illustra-  
tion is of an unnamed species.

Also 5 undescribed species (NZAC).

● **Rhathamictis** Meyrick, 1924b, p. 662. Type spe-  
cies *Mallobathra perspera* Meyrick, by original  
monotypy.

**nocturna** Clarke, 1926, p. 421 (*Mallobathra*) new  
combination

Kauri Gully, Northcote AK, C.E. Clarke; HT ♂ des-  
ignated by Clarke, AMNZ.  
Not mentioned by Hudson.

**perspera** Meyrick, 1924b, p. 662 (*Rhathamictis*)  
Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 344, pl. xl fig. 16; 1950, p. 113, pl. iv  
fig. 7.

Also 1 undescribed species (NZAC).

● **Scoriodyta** Meyrick, 1888e, p. 101. Type species  
*Scoriodyta conisalia* Meyrick, by original monotypy.  
Note. Dalla Torre & Strand (1929, p. 19) list this  
genus as *Scoriodytes* and ascribe it to Walsingham.

**conisalia** Meyrick, 1888e, p. 102 (*Scoriodyta*)  
Wellington WN, E. Meyrick; LT ♂ here designated,  
labelled “Wellington New Zealand 1/1/80”, “Scori-  
odyta conisalia Meyr. 2/13 E. Meyrick det. in Meyrick  
Coll.”, BMNH.

Hudson 1928, p. 354, pl. xxvii fig. 11 and 12.  
Also 5 undescribed species (1 NZAC, 4 Hättens-  
chwiler Collection, Uster, Switzerland).

● Genus near *Alytopistis* Meyrick, 1920b, p. 322


Two undescribed species (NZAC; adventive,  
?Australian).

● Genus indeterminate

One undetermined species (NZAC; adventive,  
origin obscure, possibly Taiwan).

—⊗—

**Family ROESLERSTAMMIIDAE**  
(in the sense of Kyrki 1983)



**Roeslerstammiidae**  
(146) *Dolichernis chloroleuca* Meyrick

● ***Dolichernis*** Meyrick, 1891, p. 99. Type species *Dolichernis chloroleuca* Meyrick, by original monotypy.

***chloroleuca*** Meyrick, 1891, p. 99 (*Dolichernis*)  
Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 325, pl. xxxvi fig. 2.

● ***Vanicela*** Walker, 1864, p. 1039. Type species *Vanicela disjunctella* Walker, by original monotypy.  
Note. This genus is removed from Heliodinidae in the sense of Meyrick (1913c, pp. 10–11; 1922c, pp. 1–4) and from Stathmopodidae in the sense of Common (1970, p. 818). It shares characters of larval structure and biology with Roeslerstammiidae (Dugdale, in prep.).


***disjunctella*** Walker, 1864b, p. 1039 (*Vanicela*)  
Auckland AK, D. Bolton; HT ♀ unique, BMNH.  
Hudson 1928, p. 298, pl. xxxii fig. 28.

? ***frontella*** Walker, 1864b, pp. 856–857  
(*Gracilaria*)  
[Auckland], “New Zealand”, D. Bolton; HT (?gender)  
unique, not in BMNH.

Note. No material collected by Bolton (i.e., bearing accession label 54.4) was found other than the HT of *disjunctella*.

—◎—

**Family GRACILLARIIDAE**



**Gracillariidae**  
(147) *Acrocercops leucocyma* (Meyrick)

Vari (1961) has restricted several genera previously used in a broad sense by Meyrick, and reinstated others. Where these latter apply to New Zealand, they are used here.

● ***Acrocercops*** in the sense of Vari (1961, pp. 146–147)

Note. Most species listed below are under *Parectopa* in Hudson (1928, 1939), but Vari (1961, pp. xii and 55–56) restricted *Parectopa* to Holarctic species which in genital characteristics do not resemble New Zealand species. As with South African species, New Zealand species “might have to be transferred to other genera when ... more data on relationship become known” (Vari 1961, p. 147).

***aellomacha*** Meyrick, 1880a, p. 158 (*Gracilaria*)  
[Botanic Gardens and forest], Wellington WN, E. Meyrick;  
HT ♂ unique, BMNH.  
Hudson 1928, pp. 322–323, pl. xxxv fig. 4.

***aethalota*** Meyrick, 1880a, pp. 143–144 (*Gracilaria*)  
[“reserved bush and forest”], Dunedin DN, E. Meyrick;  
HT ♂ unique, BMNH.  
Hudson 1928, p. 322, pl. xlvi fig. 11.

Note. Mr B. Patrick has rediscovered this species in Woodhaugh Gardens, Dunedin DN, on *Parsonsia* (Apocynaceae).

***alytidota*** Meyrick, 1880a, p. 161 (*Gracilaria*)  
Sydney N.S.W., E. Meyrick; LT ♂ here designated,  
labelled “Sydney N.S. Wales 25/10/79”, “*Acrocercops alytidota* Meyr. 12/6 E. Meyrick det. in Meyrick Coll.”,  
BMNH.

***citharoda*** Meyrick, 1916b, p. 418 (*Parectopa*). New  
synonymy.  
Wanganui WI, G.V. Hudson [M.N. Watt]; HT ♀ unique,  
BMNH.  
Hudson 1928, p. 322, pl. xxxv fig. 15.

**eumetalla** Meyrick, 1880a, p. 160 (*Gracilaria*)  
Brisbane Qld, E. Meyrick; LT ♂ here designated, labelled  
“Brisbane Queensland 29/9/79”, “Acrocercops eumetalla Meyr. 6/1 E. Meyrick det. in Meyrick Coll.”,  
BMNH.

Note. Larvae live in galls on *Acacia*. Specimens from New Zealand and Toowoomba (Qld) lack the oblique white mark at one-quarter on the forewing dorsum.

**leucocyma** Meyrick, 1889b, p. 184 (*Gracilaria*)  
[Waitakere Range], “Auckland” AK, E. Meyrick; HT ♂ unique, BMNH.

Hudson 1928, p. 322, not figured. See Wise (1962, pl. 1).

**panacicorticis** Watt, 1920, p. 457, pl. 30 fig. 5  
(*Parectopa*)

Mt Egmont TK, M.N. Watt; LT ♀ here designated, labelled “Egmont corticella”, “Morris N. Watt Coll.”, abdomen missing, NMNZ.

Hudson 1928, p. 322, as synonym of *Parectopa aethalota*. Note. The LT is the moth figured by Watt (1920, pl. 30 fig. 5).

**panacifinens** Watt, 1920, p. 460 (*Parectopa*)  
[Dunedin DN], M.N. Watt; LT ♀ here designated, labelled “M & C”, “6”, NMNZ.

Hudson 1928, p. 322, as synonym of *Parectopa aellomacha*.

Note. “M & C” denotes marginal and central [Panax moth]. LT is the specimen illustrated by Watt (1920, pl. 30 fig. 6); it agrees with Dunedin DN specimens in its brownish coloration and small, white forewing maculation. The 2 Egmont TK specimens are paler, with larger white areas on the forewing.

**panacitorsens** Watt, 1920, p. 449 (*Parectopa*)  
[Flagstaff DN], M.N. Watt; LT ♀ here designated, labelled “g”, “Giant under”, “Morris N. Watt Collection”, NMNZ.

Hudson 1928, p. 322, as synonym of *Parectopa aellomacha*.

Note. “Giant under” denotes giant mines on underside of *Panax* leaf. LT is the moth illustrated by Watt (1920, pl. 30 fig. 3).

**panacivagans** Watt, 1920, pp. 464–465 (*Parectopa*)  
Aberfeldy WI, M.N. Watt; HT ♀ unique, not located in Hudson Collection, NMNZ.

Hudson 1928, p. 322, pl. xlvi fig. 8, as species.

Note. The PT ♀ labelled “Lancewood Aberfeldy, 25/1/20” in NMNZ is not the moth figured by Watt (1920, pl. 30 fig. 10).

**panacivermiformis** Watt, 1920, pp. 452–453  
(*Parectopa*)

Mt Egmont TK, M.N. Watt; LT ♀ here designated, labelled “Egmont vermiformis”, “Morris N. Watt Collection”, NMNZ.

Hudson 1928, p. 322, as synonym of *Parectopa aellomacha*.

Note. The LT is the moth illustrated by Watt (1920, pl. 30 fig. 4).

**zorionella** Hudson, 1918, p. 62 (*Parectopa*)  
Botanical Gardens, Wellington WN, G.V. Hudson; LT ♂ labelled “861a”, “Lectotype ♂ Parectopa zorionella Hudson 861a”, Hudson Collection, NMNZ. Hudson 1928, pp. 321–322, pl. xxxv fig. 5.

Also 1 undescribed species (NZAC).

• **Caloptilia** in the sense of Vari (1961, pp. xi–xii and 3)

Note. All New Zealand species with larvae that spin leaves into a pouch are placed here.

**azaleella** Brants, 1913, p. lxxii (*Gracilaria*)  
[Ex *Azalaea indica* from Japan, Booskop, Netherlands.] New Zealand: adventive, widespread on *Rhododendron* subg. *Azalea*.

Not mentioned by Hudson.

**chalcodelta** Meyrick, 1889b, p. 183 (*Gracilaria*) new combination

Masterton WA, E. Meyrick; LT ♀ here designated, labelled “Masterton New Zealand 11/3/83”, “Gracilaria chalcodelta Meyr. 2/17 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 324, pl. xxxv fig. 7.

**chrysitis** Felder & Rogenhofer, 1875, pl. cxl fig. 143  
(*Gracilaria*) new combination

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1928, p. 324, pl. xxxv fig. 10.

**rutilans** Butler, 1880, p. 561 (*Gracilaria*). Synonymised by Meyrick (1889b, p. 183).

[forests north of] Blenheim MB, W. Skellon; HT ♂ unique, BMNH.

Hudson 1928, p. 324, as synonym.

**adelina** Meyrick, 1880a, p. 142 (*Gracilaria*). Synonymised by Meyrick (1889b, p. 183).

Hamilton WO, E. Meyrick; LT ♂ selected by P.A. Brown, labelled “Hamilton New Zealand 18/1/80”, “Gracilaria chrysitis Feld. 10/17 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 324, as synonym.

**purpurea** Philpott, 1927d, p. 89 (*Gracilaria*). New synonymy.

West Plains [Invercargill] SL, A. Philpott; HT ♂ designated by Philpott, abdomen missing, NZAC.

Hudson 1939, p. 461, pl. lxviii fig. 6.

**elaeas** Meyrick, 1911b, p. 66 (*Gracilaria*) new combination

Castle Hill [homestead] MC, E. Meyrick; LT ♂ here designated, labelled “Castle Hill New Zealand bred 16/2/83”, “Gracilaria elaeas Meyrick 3/14 E. Meyrick det. in Meyrick Coll.”, BMNH.

» Gracillariidae, *Caloptilia elaeas*

Hudson 1928, p. 323, pl. xxxv fig. 3.

Note. Butler (1877, pl. xlivi fig. 16) illustrates a ♀ of this species (cf. *linearis*, below). On 16 February 1883 Meyrick collected at around 2500 ft, i.e., about the altitude of the station homestead, but it is not known from what species of *Coriaria* (Coriariaceae) he collected his *C. elaeas* larvae. The status of *C. elaeas* and *C. linearis* (Butler) requires closer examination. My observations suggest that *C. elaeas* is on summer-green *Coriaria* species, and *C. linearis* on evergreen or arborescent *Coriaria* species.

***linearis*** Butler, 1877, p. 406 (*Gracilaria*) new combination

[Canterbury or Otago], J. Hector; HT ♀ ("Type" in Butler's handwriting), abdomen missing, BMNH.

Hudson 1928, p. 323, pl. xxxv fig. 6.

Note. See note under *elaeas*, above. Hector's collection, though sent from Dunedin, was recorded in the BMNH accession book as largely collected by J.D. Enys.

***selenitis*** Meyrick, 1909a, p. 15 (*Gracilaria*) new combination

Mt Holdsworth WN, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 324, pl. xxxv fig. 1.

Also 1 undescribed species (NZAC).

● ***Conopomorpha*** Meyrick, 1885i, p. 592; 1886b, p. 183. Type species *Conopomorpha cyanospila* Meyrick, by original monotypy.

***cyanospila*** Meyrick, §1885i, p. 592; 1886b, p. 183 (*Conopomorpha*)

"Taranaki" [New Plymouth] TK, E. Meyrick; LT ♀ here designated, labelled "Taranaki New Zealand 28/2/82", "Acrocercops cyanospila Meyr. 2/12 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 321, pl. xxxv fig. 14.

Note. *Conopomorpha* was reinstated by Vari (1961, p. 102). "Taranaki" or "Taranaki Roadstead" is an early name for the port of New Plymouth.

● ***Lithocolletis*** of authors, but not of Hübner (1825).

***hardenbergiella*** Wise, 1957, p. 26 (*Lithocolletis*)  
Epsom AK, K.A.J. Wise; HT ♂ designated by Wise, NZAC.

Note. Adventive; the host plant is Australian. The colony known to Wise has been destroyed, and I have been unable to locate others.

● ***Macarostola*** Meyrick, 1907, p. 62. Type species *Gracilaria formosa* Stainton, 1862, by original designation; Australia.

Note. Kumata (1977, pp. 34–36) lists all known species of *Macarostola*.

***miniella*** Felder & Rogenhofer, 1875, pl. cxl fig. 42  
(?*Stathmopoda*) new combination

[?Auckland AK, A. Sinclair], "N. Seeld"; HT ♂ unique, BMNH.

Hudson 1928, p. 323, pl. xxxv fig. 8 and 9.

***ethela*** Meyrick, 1880a, pp. 152–153 (*Gracilaria*).  
Synonymised by Meyrick (1889, p. 185).

[Hamilton WO], E. Meyrick; LT ♂ selected by P.A. Brown and here designated, labelled "Taranaki New Zealand 18.1.80", "Lectotype teste P.A. Brown", BMNH.

Hudson 1928, p. 323, as synonym.

Note. Meyrick's Diary of Captures records that he collected at Hamilton on 18 January 1880.

● ***Phyllonorycter*** Hübner, 1822, pp. 66–74 and 76–80. Type species *Phalaena (Tinea) rajella* Linnaeus, 1758, by subsequent designation (Fletcher 1928, p. 176); Europe.

***messaniella*** Zeller, 1846, p. 221 (*Lithocolletis*)

Europe.  
New Zealand: adventive; widespread, on *Quercus* and *Fagus*.




**Superfamily YPONOMEUTOIDEA**

(in the sense of Kyrki 1984)

**Family YPONOMEUTIDAE**

(in the sense of Moriuti 1977, pp. 16 and 25–38)



**Yponomeutidae**  
(148) *Orthenches disparilis* Philpott

Note. No attempt is made to assign New Zealand genera to the subfamilial or tribal classification of Moriuti (1977, pp. 28–37), nor to the generic groups delimited by Kyrki (1984).

- **Cadmogenes** Meyrick, 1923, pp. 167–168. Type species *Cadmogenes literata* Meyrick, 1923, by original monotypy.

*literata* Meyrick, 1923, p. 168 (*Cadmogenes*)  
Kauri Gully, Birkenhead AK, G.V. Hudson; LT ♂ here  
designated, labelled “Kauri Gy 27.1.21”, “Cadmogenes  
literata Meyr. 1/1 E. Meyrick det. in Meyrick  
Coll.”, head and abdomen missing, left forewing and  
hindwing in gelatin capsule, BMNH.  
Hudson 1928, pp. 327–328, pl. xl ix fig. 11.

Note. *Cadmogenes literata* is univoltine on flowers of  
*Ackama* and *Weinmannia* (Cunoniaceae). The  
description mentions a specimen each from Silver-  
stream (Hutt Valley WN), where *Weinmannia* is  
abundant, and Kauri Gully; the Silverstream speci-  
men is missing from Meyrick's collection. At Kauri  
Gully *Weinmannia* is now a rare tree. The venation of  
the LT agrees with that of specimens reared from  
*Weinmannia* inflorescences.

- **Charixena** Meyrick, 1921, p. 335, new name for  
*Philpottia* Meyrick, 1916b, p. 416, preoccupied by  
*Philpottia* Broun, 1915 (Coleoptera). Type species  
*Philpottia iridoxa* Meyrick, 1916, by original  
monotypy.

*iridoxa* Meyrick, 1916b, p. 417 (*Philpottia*)  
Mount Burns, [Hunter Mountains] FD, A. Philpott; LT  
♀ here designated, labelled “Mt Burns New Zealand  
AP 29.12.14”, “Charixena iridoxa Meyr. 1/2 E. Meyr-  
ick det. in Meyrick Coll.”, abdomen in gelatin cap-  
sule, BMNH.  
Hudson 1928, p. 317, pl. xxxvii fig. 16, in  
“*Glyptapterigides*”.

- **Doxophyrtis** Meyrick, 1914a, p. 112. Type species  
*Doxophyrtis hydrocosma* Meyrick, by original  
monotypy.

*hydrocosma* Meyrick, 1914a, p. 113 (*Doxophyrtis*)  
Kao ND, G.V. Hudson; LT ♂ here designated, labelled  
“Kao New Zealand G.V.H. 1.13”, “Doxophyrtis  
hydrocosma Meyr. 1/3 E. Meyrick det. in Meyrick  
Coll.”, BMNH.  
Hudson 1928, p. 325, pl. xxxiv fig. 15.

- **Hierodoris** of authors, but not Meyrick (1912d,  
p. 42)

*stellata* Philpott, 1918, pp. 129–130 (?*Hierodoris*)  
Blue Cliff [Te Waewae Bay] FD, C.C. Fenwick; HT ♂  
designated by Philpott, NMNZ.  
Hudson 1928, p. 305, pl. xxxviii fig. 21, in  
“*Glyptapterygides*”.

- **Kessleria** Nowicki, 1864, p. 12, in the sense of  
Moriuti (1977, p. 207). Type species *Kessleria zimmermanni*  
Nowicki, 1864, as reported by Moriuti  
(1977, p. 207).

*Circostola* Meyrick, 1889b, p. 163. Type species  
*Circostola copidota* Meyrick, by original mono-  
typy. Synonymised by Moriuti (1977, p. 207).  
Note. Moriuti (1977, p. 207) gives full synonymy.

*copidota* Meyrick, 1889b, p. 163 (*Circostola*)  
[Otira Gorge, 1,600 ft], Otira River WD, E. Meyrick; LT  
♂ designated by Moriuti (1977, p. 208), BM genitalia  
slide no. 19506, BMNH.  
Hudson 1928, p. 320, pl. xxxv fig. 11.

- **Orthenches** Meyrick, §1885i, p. 591; 1886b, pp.  
173 (key) and 175. Type species *Orthenches*  
*chlorocoma* Meyrick, by subsequent designation  
(Meyrick 1915b, p. 230).

*chartularia* Meyrick, 1924a, pp. 205–206  
(*Orthenches*)  
Mount Ruapehu TO, 4,000 ft, G.V. Hudson; HT ♂  
unique, BMNH.  
Hudson 1928, p. 328, pl. 1 fig. 25.

*nivalis* Philpott, 1927d, p. 89 (*Orthenches*). Syn-  
onymised by Philpott (1931, p. 34).  
Arthur's Pass NC/WD, A. Philpott; HT ♀ designated by  
Philpott, NZAC.

*chlorocoma* Meyrick, §1885i, p. 591, as *chloro-  
croma*; 1886b, pp. 175–176, as *chlorocoma*  
Christchurch MC, E. Meyrick; HT ♂ unique, labelled  
“chlorocoma Meyr.” in Meyrick's handwriting,  
BMNH.  
Hudson 1928, pp. 329–330, pl. xxxviii fig. 9.

➤ Yponomeutidae, *Orthenches*

- dictyarcha** Meyrick, 1927b, p. 315 (*Orthenches*)  
Arthur's Pass NC/WD, G.V. Hudson; HT ♀ unique,  
abdomen missing, BMNH.  
Hudson 1939, p. 462, pl. lxi fig. 29.
- disparilis** Philpott, 1931, p. 34 (*Orthenches*)  
Kauri Gully, [Birkenhead] AK, C.E. Clarke; HT ♂ unique,  
AMNZ.  
Hudson 1939, p. 462, not figured.
- drosochalca** Meyrick, 1905, p. 242 (*Orthenches*)  
Wellington WN, G.V. Hudson; LT ♂ selected by P.A.  
Brown and here designated, labelled "Wellington New  
Zealand GVH /03", "Orthenches drosochalca Meyr.  
2/5 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 328, pl. xxxvi fig. 4.  
Note. Hudson was misled by larvae pupating on "silvery  
tree fern"; the true hosts are *Prumnopitys ferruginea*  
and *P. taxifolia* (Podocarpaceae; species formerly in  
*Podocarpus*).
- glypharcha** Meyrick, 1919, p. 353 (*Orthenches*)  
Mount Egmont TK, 3,000 ft, G.V. Hudson; LT ♀ here  
designated, labelled "Mt Egmont New Zealand GVH  
3000' 2.18", "Orthenches glypharcha Meyr. 2/2 E.  
Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 330, pl. xlvi fig. 19.
- homerica** Salmon, 1956, p. 576 (*Archyalia*) new  
combination  
Homer Cirque FD, J.T. Salmon; HT ♂ designated by  
Salmon, NMNZ.  
Salmon 1956, pl. 2 fig. 4 and 5.  
Note. This species is transferred from Tineidae. It is  
scarcely distinguishable from *O. dictyarcha*.
- polita** Philpott, 1918, p. 131 (*Orthenches*)  
[Tisbury], Invercargill SL, A. Philpott; HT ♂ designated  
by Philpott, abdomen missing, NZAC.  
Hudson 1928, p. 329, pl. xlvi fig. 4.
- porphyritis** Meyrick, §1885i, p. 591; 1886b, pp. 176–  
177 (*Orthenches*)  
Dunedin DN, E. Meyrick; LT ♀ here designated, labelled  
"Dunedin New Zealand 28.9.82", "Orthenches porphyritis  
Meyr. 8/15 E. Meyrick det. in Meyrick Coll.",  
BMNH.  
Hudson 1928, pp. 328–329, pl. xxxvi fig. 6 and 29, pl.  
xlvi fig. 17.
- cuprea** Meyrick, 1901, pp. 575–576 (*Ypono-  
meuta*). Synonymised by Meyrick (1923, p. 168).  
?Christchurch MC, R.W. Fereday; HT ♀ unique, abdo-  
men missing, BMNH.
- prasinodes** Meyrick, §1885i, p. 591; 1886b, p. 176  
(*Orthenches*)  
Riccarton Bush, Christchurch MC, E. Meyrick; HT ♂  
unique, BMNH.
- Hudson 1928, p. 329, pl. xxxvi fig. 5.
- saleuta** Meyrick, 1913a, p. 28 (*Orthenches*)  
Waiouru TO, G.V. Hudson; LT ♀ here designated,  
labelled "Waiouru New Zealand GVH 2.12",  
"Orthenches saleuta Meyr. 2/3 E. Meyrick det. in  
Meyrick Coll.", BMNH.  
Hudson 1928, p. 328, pl. xxxvi fig. 3.
- semifasciata** Philpott, 1915, pp. 200–201 (*Orth-  
enches*)  
Queenstown OL, C.C. Fenwick; HT ♀ designated by  
Philpott, NMNZ.  
Hudson 1928, p. 329, pl. xl fig. 3.
- septentrionalis** Philpott, 1930b, p. 14 (*Orthenches*)  
Kauri Gully, Birkenhead AK, C.E. Clarke; HT ♂ desig-  
nated by Philpott, AMNZ.  
Hudson 1939, p. 462, pl. lvii fig. 21.
- similis** Philpott, 1924a, p. 211 (*Orthenches*)  
Dun Mountain NN/MB, A. Philpott; HT ♂ designated  
by Philpott, NZAC.  
Hudson 1928, p. 329, as synonym of *O. semifasciata*.  
Note. Philpott (1931, p. 35) gave distinguishing features.
- vinitincta** Philpott, 1917b, pp. 244–245 (*Orthenches*)  
Rowallan FD, C.C. Fenwick; HT ♂ designated by Phil-  
pott, abdomen missing, NMNZ.  
Hudson 1928, p. 329, pl. xxxviii fig. 20.  
Note. Externally this species appears identical to *O.  
prasinodes*.
- virgata** Philpott, 1920, p. 44 (*Orthenches*)  
Auckland AK, A.J. Turner; HT ♀ designated by Philpott,  
abdomen missing, NZAC.  
Hudson 1928, p. 330, pl. xlvi fig. 18.
- Also 2 undescribed species (NZAC).
- **Phylacodes** Meyrick, 1905, pp. 241–242. Type  
species *Phylacodes cauta* Meyrick, by original  
monotypy.
- cauta** Meyrick, 1905, p. 242 (*Phylacodes*)  
[Ida Valley CO], J.H. Lewis; HT ♂ unique, BMNH.  
Hudson 1928, p. 327, pl. xxxiv fig. 16.
- **Plutella** Schrank, 1802, p. 169, in the sense of  
Moriuti (1977, p. 52). Type species *Phalaena Tinea  
xylostella* Linnaeus, as reported in Moriuti (1977,  
p. 52).
- antiphona** Meyrick, 1901, p. 576 (*Plutella*)  
Wellington WN, G.V. Hudson; HT ♀ unique, BM gen-  
italia slide no. 18897, BMNH.  
Hudson 1928, p. 331, not figured; Dugdale 1973b, p. 1013,  
fig. 2, 8, 13, 15, 17, and 19.

Note. *P. antiphona* is another ‘mystery species’ supposedly collected by Hudson but of which he appeared to have no knowledge (see *Mallophthora globulosa*, p. 68). There is some evidence that W.G. Howes (Dundin DN) sent specimens to Meyrick, who may have mistaken them for Wellington material. See *Proditrix megalynata*, below.

***xylostella*** Linnaeus, 1758, p. 538, no. 265 (*Phalaena Tinea*). ICBN Opinion 1002 (1973, p. 86). Cosmopolitan.

New Zealand: recorded throughout.

Hudson 1928, p. 331, pl. xxxvi fig. 7 and 8, as *Plutella maculipennis* Curtis. Moriuti (1977, pp. 53–54) gives full synonymy.

Also 1 undescribed species (NZAC).

• ***Plutella*** in the sense of Meyrick (1915, p. 230)

***psammochroa*** Meyrick, §1885i, p. 591; 1886b, p. 179 (*Plutella*)

[Arthur’s Pass NC/WD, at light], “Otira River”, E. Meyrick; LT ♀ selected by K. Sattler, designated by Dugdale (1973b, p. 1019), BMNH.

Hudson 1928, p. 331, pl. xxxvi fig. 10.

Note. Meyrick’s Diary of Captures notes “*P. catapela* 1 at light” at Arthur’s Pass on 29 January 1883, and another at Castle Hill NC on 17 January 1883. The latter specimen is labelled “*Plutella psammochroa* Meyrick 3/3 E. Meyrick det. in Meyrick Coll.”, “Castle Hill New Zealand 17.1.83”. The Australian specimens differ in 6 colour pattern characters from the New Zealand specimens, and are doubtfully conspecific.

• ***Prays*** Hübner, 1826, p. 413. Type species [*Phalaena curtisella* Donovan, 1793], as reported by Moriuti (1977, p. 116).

***nephelomima*** Meyrick, 1907b, pp. 75 (key) and 76 (*Prays*)

Murrurundi and Sydney N.S.W., Australia; ST series, including BM genitalia slide no. 3275 ♂, BMNH.

New Zealand: adventive, recorded from Auckland AK and Tauranga BP 1976–78, Kerikeri ND 1981–82.

Note. Classification of the *Citrus*-infesting *Prays* species is inadequate. Although superficially *P. nephelomima* closely resembles the type (and topotypic) specimens of *P. citri* Millière from Corsica, there are genital differences. It should be noted that both *Citrus* and *P. citri* are Oriental adventives to Corsica.

• ***Proditrix*** Dugdale, 1987a, pp. 99–100. Type species *Plutella megalynata* Meyrick, by original designation.

***chionochloae*** Dugdale, 1987a, p. 106 (*Proditrix*) Pouakai Range TK, J.S. Dugdale; HT ♂ designated by Dugdale, NZAC.

***gahniae*** Dugdale, 1987a, p. 108 (*Proditrix*) Titirangi AK, C.R. Thomas; HT ♂ designated by Dugdale, NZAC.

***megalynata*** Meyrick, 1915a, pp. 203–204 (*Plutella*) [Otago–Southland, W.G. Howes], “Wellington, Philpott”; HT ♂ unique, BMNH.

Hudson 1928, p. 330, pl. xxxvi fig. 1.

Note. Philpott (1917a, p. 235, footnote) states “The type specimen [i.e., the one Philpott sent to Meyrick] was given to me by Mr W.G. Howes. It was among some Wellington Lepidoptera, but as there is a little uncertainty as to the locality it is advisable, for the present, to record it as a southern form only”. Philpott (1927g, p. 318) had difficulty interpreting the highly modified ♂ genitalia.

***tetragona*** Hudson, 1918, p. 62 (*Titanomis*)

Mt Egmont TK, M.N. Watt; HT ♀ (recorded as ♂) unique, NMNZ.

Hudson 1928, p. 350, pl. xxxv fig. 28.

Note. Structure of the ♂ genitalia and larva indicate a close relationship with *P. megalynata*. Dugdale (1987a) assigned *tetragona* to *Proditrix*.

• ***Protosynaema*** Meyrick, §1885i, p. 591; 1886b, p. 174. Type species *Protosynaema eratopis* Meyrick, by subsequent designation (Meyrick 1915b, p. 229).

***eratopis*** Meyrick, §1885i, p. 591; 1886b, p. 174 (*Protosynaema*)

Otira Gorge WD, 1,600 ft, E. Meyrick; LT ♂ here designated, labelled “Otira Gorge New Zealand 26.1.83”, “*Protosynaema eratopis* Meyr. 6/16 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 327, pl. xxxiv fig. 1.

***hymenopis*** Meyrick, 1935, p. 304 (*Protosynaema*)

Gollan’s Valley WN, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1939, p. 461, pl. lx fig. 25.

***matutina*** Philpott, 1928g, p. 489 (*Protosynaema*)

Mount Arthur NN, 4,500 ft, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 461, pl. lx fig. 15.

***quaestuosa*** Meyrick, 1924a, p. 205 (*Protosynaema*)

Mount Aurum OL, G.V. Hudson; LT ♂ here designated, labelled “Mt Aurum New Zealand GVH bred .21”, “*Protosynaema questuosa* [sic] Meyr. 3/7 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 326, pl. 1 fig. 13, pl. xxxiv fig. 3.

***steropucha*** Meyrick, §1885i, p. 591; 1886b, pp. 174–175 (*Protosynaema*)

Kaiapoi NC, [sandhills], E. Meyrick; LT ♂ here designated, labelled “Kaiapoi New Zealand 31.3.82”, “*Protosynaema steropucha* Meyr. 8/14 E. Meyrick det. in Meyrick Coll.”, abdomen in gelatin capsule, BMNH.

» Yponomeutidae, *Protosynaema steropucha*

Hudson 1928, p. 326, pl. xxxiv fig. 2.

Also 1 undescribed species (B. Patrick Collection, Dunedin).

● **Rhigognostis** Zeller in Staudinger, 1857, pp. 273–274. Type species *Plutella dalella* Stainton [by implication?].

**Caunaca** Wallengren, 1880, p. 45. Type species *Cerostoma annulatella* Curtis, 1832, by subsequent designation (Fletcher 1929). Synonymised by J. Kyrki (unpublished).

Note. Moriuti (1977, p. 59) gives synonymy.

**sera** Meyrick, §1885i, p. 591; 1886b, p. 178 (*Plutella*) Makatoku HB, E. Meyrick; LT ♂ selected by K. Sattler, designated by Dugdale (1973, p. 1017), BMNH. Hudson 1928, p. 330, pl. xxxvi fig. 9, as *Plutella sera*.

● **Tanaoctena** Turner, 1913, p. 204. Type species *Tanaoctena ooptila* Turner, 1913, p. 205, by monotypy; Australia.

**Tanaoctenota** Meyrick, 1918b, p. 188; replacement name for *Tanaoctena* Turner (possible confusion with *Tanaoctenia* Warren – Geometridae). Synonymised by Clarke (1971, p. 167).

**dubia** Philpott, 1931, p. 34 (*Tanaoctenota*) Auckland AK, C.E. Clarke; HT ♂ designated by Philpott, AMNZ. Hudson 1939, p. 441, pl. lx fig. 8, as *Tanaoctenota dubia*.

● **Zelleria** of authors, in the sense of Meyrick (1915b, p. 227) but not Stainton (1849, p. 22)


**maculata** Philpott, 1930b, pp. 13–14 (*Zelleria*) Mount Maungatua DN, C.E. Clarke; HT ♀ designated by Philpott (as ♂), AMNZ. Hudson 1939, p. 459, pl. lxi fig. 27.

**porphyraula** Meyrick, 1927b, p. 314 (*Zelleria*) Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1939, p. 460, pl. lxi fig. 6.

**rorida** Philpott, 1918, p. 130 (*Zelleria*) Bluff SL, in coastal forest, A. Philpott; HT ♂ designated by Philpott, this and AT ♀ lacking locality labels, NZAC. Hudson 1928, pp. 320–321, pl. xxxviii fig. 14.

**sphenota** Meyrick, 1889b, p. 162 (*Hofmannia*) [Riccarton Bush], Christchurch MC, E. Meyrick; HT ♂ unique, BMNH. Hudson 1928, p. 321, not figured.

**Family GLYPHIPTERIGIDAE**  
(in the sense of Heppner 1982, p. 38)



**Glyphipterygidae**  
(149) *Glyphipteryx zelota* Meyrick

● **Glyphipteryx** not of Hübner, [1825], p. 421, but in the sense of Meyrick (1913, p. 41)

**Phryganostola** Meyrick, 1880b, p. 248. Type species *Phryganostola drosophae* Meyrick, by subsequent designation (Meyrick 1907b, p. 115). Synonymised by Meyrick (1913c, p. 41).

**Circica** Meyrick, 1888e, p. 88. Type species *Circica cionophora* Meyrick, by subsequent designation (Meyrick 1913c, p. 41). Synonymised by Meyrick (1913c, p. 41).

**achlyoessa** Meyrick, 1880b, p. 252 (*Phryganostola*) [Botanic Gardens], Wellington WN, E. Meyrick; HT ♂ unique, BMNH. Hudson 1928, p. 312, pl. xxxiii fig. 11 and 12.

**acronoma** Meyrick, 1888e, p. 86 (*Glyphipteryx*) Mount Arthur NN, E. Meyrick; LT ♂ here designated, labelled “Mt Arthur New Zealand 18.1.86”, “*Glyphipteryx acronoma* Meyr. 1/5 E. Meyrick det. in Meyrick Coll.”, BMNH. Hudson 1928, p. 316, pl. xlvi fig. 14.

**acrothecta** Meyrick, 1880b, pp. 244–245 (*Glyphipteryx*) [Port Hills], Christchurch MC, E. Meyrick; LT ♀ here designated, labelled “Christchurch New Zealand 4.1.80”, “*Glyphipteryx acrothecta* Meyr. 1/6 E. Meyrick det. in Meyrick Coll.”, BMNH. Hudson 1928, p. 316, pl. xxxiv fig. 5.

**aenea** Philpott, 1917b, p. 244 (*Glyphipteryx*) The Hump [Hump Ridge] FD, A. Philpott; HT ♂ designated by Philpott, NZAC. Hudson 1928, p. 311, pl. xxxiv fig. 7.

**aerifera** Meyrick, 1912d, p. 57 (*Glyphipteryx*) Mount Ruapehu TO, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 313, pl. xxxiii fig. 22.

**astrapaea** Meyrick, 1880b, p. 245 (*Glyphipteryx*)  
Cambridge WO, E. Meyrick; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 313, not figured.

Note. The HT is scarcely distinguishable from that of *G. codonias* Meyrick (see below).

**ataracta** Meyrick, 1888e, p. 88 (*Phryganostola*)

Mount Arthur NN, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 16.1.86", "Glyphipteryx ataracta Meyr. 4/7 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 312, pl. xlvi fig. 7.

**aulogramma** Meyrick, 1907c, p. 120 (*Glyphipteryx*)

Invercargill SL, A. Philpott; LT ♂ here designated, labelled "Invercargill New Zealand AP '06", "Glyphipteryx aulogramma Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pp. 312–313, pl. xxxiii fig. 16 and 17.

**bactrias** Meyrick, 1911b, pp. 67–68 (*Glyphipteryx*)

Invercargill SL, A. Philpott; LT ♂ here designated, labelled "Invercargill New Zealand AP 10.1.09", "Glyphipteryx bactrias Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 312, pl. xxxiii fig. 9.

**barbata** Philpott, 1918, p. 130 (*Glyphipteryx*)

Waitati DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1928, p. 317, pl. xxxviii fig. 13.

**brachydelta** Meyrick, 1916b, p. 418 (*Glyphipteryx*)

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 3.15", "Glyphipteryx brachydelta Meyr. 1/6 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 315, pl. xxxviii fig. 12.

**calliactis** Meyrick, 1914a, p. 112 (*Glyphipteryx*)

Kaitoke WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 314, pl. xxxiv fig. 12 and 13.

**cionophora** Meyrick, 1888e, p. 88 (*Circica*)

[Port Hills], Christchurch MC, E. Meyrick; LT ♂ selected by J.B. Heppner, 1976, and here designated, labelled "Christchurch New Zealand 24 II 1882 Meyrick 1888 1740", "Circica cionophora Meyr. named by Meyrick", "Walsingham Collection 1910-427", BMNH.  
Hudson 1928, p. 311, pl. xxxiii fig. 10.

**codonias** Meyrick, 1909a, p. 15 (*Glyphipteryx*)

Invercargill SL, A. Philpott; HT ♂ unique, BMNH.  
Hudson 1928, p. 313, not figured, as synonym of *G. transversella*.

Note. The HTs of *G. codonias* and *G. astrapaea* are scarcely distinguishable on external characters.

**dichorda** Meyrick, 1911b, p. 76 (*Glyphipteryx*)  
[?Wellington WN], "New Zealand", G.V. Hudson; LT here

designated, labelled "New Zealand GVH /00", "Glyphipteryx dichorda Meyr. 3/3 E. Meyrick det. in Meyrick Coll.", hindwings and abdomen missing, BMNH.

Hudson 1928, p. 315; 1939, p. 458, pl. lx fig. 19.

**erastis** Meyrick, 1911b, p. 76 (*Glyphipteryx*)

Lake Wakatipu OL, E. Meyrick; LT ♂ here designated, labelled "L. Wakatipu New Zealand 15.12.82", "Glyphipteryx erastis Meyr. 3/10 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 317, pl. xxxiv fig. 20.

**euastera** Meyrick, 1880b, p. 236 (*Glyphipteryx*)

[Port Hills], Christchurch MC, E. Meyrick; LT ♀ here designated, labelled "Christchurch New Zealand 4.1.80", "Glyphipteryx euastera Meyr. 3/7 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pp. 315–316, pl. xxxiv fig. 21.

**iocheaera** Meyrick, 1880b, p. 243 (*Glyphipteryx*)

Dunedin DN, [reserved bush and forest], E. Meyrick; LT ♂ here designated, labelled "Dunedin New Zealand 6.1.80", "Glyphipteryx iocheaera Meyr. 10/15 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 314, pl. xxxiv fig. 18.

Note. The Australian *G. palaemorpha* Meyrick is very similar in shape and wing pattern.

**leptosema** Meyrick, 1888e, p. 87 (*Glyphipteryx*)

[Waitakere Range] AK, E. Meyrick; HT ♀ unique, BMNH.

Hudson 1928, pp. 314–315, pl. xxxiv fig. 9.

Note. The HT bears a label "date (month) is wrong for HT but all else agrees. J.S. Dugdale 5.9.80". Meyrick's Diary of Captures states for 17 December 1885: "Waitakere Ranges ... Glyph. leptosema, 1."; cf. his description "in January".

**metasticta** Meyrick, 1907c, pp. 19–20 (*Glyphipteryx*)

Invercargill SL, A. Philpott; LT ♂ here designated, labelled "Invercargill New Zealand AP .06", "Glyphipteryx metasticta Meyr. 1/3 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 312, pl. xxxiii fig. 15.

**morangella** Felder & Rogenhofer, 1875, pl. cxl fig. 39 (*Glyphipteryx*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, body/wings glued, BMNH.

Hudson 1928, p. 313, not figured, as synonym of *transversella*.

Note. This minute, forgotten species (wingspan 7.0 mm) is represented in NZAC by a series taken by Philpott at Golden Downs NN and specimens from a sedge area on Aorangi, Poor Knights Islands ND. It superficially resembles *G. scolias* from the Kermadec Islands.

**necopina** Philpott, 1927d, p. 88 (*Glyphipteryx*)

Golden Downs NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, pp. 457–458, pl. lx fig. 20.

► *Glyphiptericidae, Glyphiptericix*

- nephoptera** Meyrick, 1888e, pp. 87–88  
*(Glyphipteryx)*  
 [Port Hills], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 24.2.82”, “*Glyphipteryx nephoptera* Meyr. 3/13 E. Meyrick det. in Meyrick Coll.”, BMNH.  
 Hudson 1928, p. 316, pl. xxxiv fig. 19.
- octoronia** Philpott, 1924a, p. 210 (*Glyphipteryx*)  
 Gouland Downs NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
 Hudson 1928, p. 313, pl. 1 fig. 3.
- oxymachaera** Meyrick, 1880b, pp. 251–252  
*(Phryganostola)*  
 [Port Hills], Christchurch MC, E. Meyrick; LT ♀ here designated, labelled “Christchurch New Zealand 4.1.80”, “*Glyphipteryx oxymachaera* Meyr. 11/18 E. Meyrick det. in Meyrick Coll.”, BMNH.  
 Hudson 1928, pp. 313–314, pl. xxxiii fig. 18 and 19.
- rugata** Meyrick, 1915a, p. 203 (*Glyphipteryx*)  
 Tisbury, Invercargill SL, A. Philpott; HT ♀ unique, BMNH.  
 Hudson 1928, p. 312, pl. xxxiii fig. 13.
- scinteella** Walker, 1864b, p. 841 (*Glyphipteryx*)  
 Auckland AK, D. Bolton; HT ♂ unique, BMNH.  
 Omitted by Hudson 1928, 1939.
- transversella** Walker, 1864b, p. 849 (*Argyresthia*).  
 New synonymy.  
 [Nelson NN], T.R. Oxley; LT (?♂) here designated, labelled “Auckland N. Zeal. 60–73”, “LECTOTYPE *Argyresthia transversella* Wkr des. J.S. Dugdale 8.9.1980”, abdomen and hindwings missing, BMNH.  
 Hudson 1928, p. 313, pl. xxxiii fig. 20.  
 Note. The type specimens of *scinteella* and *transversella* both differ markedly from those of *codonias* and *astraea* in their possession of fields of non-metallic yellow scales on the forewing, and in the position of the silver band on the dorsum. This group (including *aulogramma*) is characteristic of *Carex* (Cyperaceae) communities.
- scintilla** Clarke, 1926, p. 420 (*Glyphipteryx*)  
 Hunter Mountains FD, [Flat Top, 4000 ft], C.E. Clarke;  
 HT ♂ designated by Clarke, AMNZ.  
 Hudson 1928, p. 315, pl. lii fig. 17.
- similis** Philpott, 1928a, pp. 369–370 (*Glyphipteryx*)  
 Mount Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
 Hudson 1939, p. 458, pl. lx fig. 22.
- triselena** Meyrick, 1880b, p. 234 (*Glyphipteryx*)  
 [Port Hills], Christchurch MC, E. Meyrick; LT ♀ here designated, labelled “Christchurch New Zealand 4.1.80”, “*Glyphipteryx triselena* Meyr. 11/14 E. Meyrick det. in Meyrick Coll.”, BMNH.  
 Hudson 1928, p. 317, pl. xxxiv fig. 6.

Note. Meyrick (1882b, pp. 188–189) gave an extensive redescription. The LT is the surviving member of the original series.

- tungella** Felder & Rogenhofer, 1875, pl. cxl fig. 40  
*(Glyphipteryx)*

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.  
 Hudson 1928, p. 315, not figured, as ?synonym of *asteronota*.

- asteronota** Meyrick, 1880b, pp. 240–241 (*Glyphipteryx*). **New synonymy.**

[The Domain], Auckland AK, E. Meyrick; LT ♀ here designated, labelled “Auckland New Zealand 20.1.80”, “*Glyphipteryx asteronota* Meyr. 5/13 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 315, pl. xxxiv fig. 11, as species.

- plagigera** Philpott, 1916, p. 423 (*Glyphipteryx*). **New synonymy.**

Bluff SL, A. Philpott; HT ♂ designated by Philpott, only the head and left forewing remaining, NZAC.  
 Hudson 1928, p. 315, not figured, as synonym of *dichorda*.

- xestobela** Meyrick, 1888e, p. 89 (*Circica*)

Arthur's Pass NC/WD, E. Meyrick; LT ♂ here designated, labelled “Arthur's Pass New Zealand 29.1.83”, “*Glyphipteryx xestobela* Meyr. 4/6 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 311, pl. xlvi fig. 22 and 37.

- zelota** Meyrick, 1888e, p. 86 (*Glyphipteryx*)

Whangarei [Heads] ND, E. Meyrick; LT ♂ here designated, labelled “Whangarei New Zealand 20.12.85”, “*Glyphipteryx zelota* Meyr. 1/13 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 316, pl. xxxiv fig. 10.

Note. There are differences in colour pattern between North Island and South Island specimens.


- **Pantosperma** Meyrick, 1888e, p. 89. Type species *Pantosperma holochalca* Meyrick, by original monotypy.

- holochalca** Meyrick, 1888e, p. 89 (*Pantosperma*)

Makatoku HB, E. Meyrick; LT ♂ selected by J.B. Heppner and here designated, labelled “Makatoku New Zealand 8.III.1883 Meyrick 1888, 1691”, “LECTOTYPE ♂ *Pantosperma holochalca* Meyr. by Heppner '78”, BMNH.  
 Hudson 1928, p. 311, pl. xxxiii fig. 14.

—◎—

**Family LYONETIIDAE**  
(in the sense of Kyrki 1984)




---

**Lyonetiidae**  
(150) *Bedellia psamminella* Meyrick

---

● **Bedellia** Stainton, 1849, p. 23. Type species *Lyonetia somnulentella* Zeller, by original monotypy.

*psamminella* Meyrick, 1889b, p. 165 (*Bedellia*)  
[Dry Bush, Port Hills], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 27.12.82", "Bedellia psamminella Meyr. 13/13 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 332; 1939, p. 463, pl. lix fig. 21.  
Note. Bradley's synonymy of *somnulentella* includes *mnesileuca* Meyrick from eastern Australia (Bradley 1961, pp. 160–161), but he did not report on *psamminella*.

*somnulentella* Zeller, 1847, p. 894 (*Lyonetia*)  
Syracuse, Sicily; LT ♀ designated by J.D. Bradley in 1960, BMNH.  
Hudson 1928, p. 332, not figured.)  
Note. Philpott's (1927h, fig. 10) drawing of ♂ genitalia of "B. Somnulentella" is not detailed enough to resolve synonymy.

● **Cateristis** Meyrick, 1889b, pp. 163–164. Type species *Cateristis eustyla* Meyrick, by original monotypy.

*eustyla* Meyrick, 1889b, p. 164 (*Cateristis*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 23/12/82", "Cateristis eustyla Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 332, not figured.  
Note. *C. eustyla* has not been recorded in New Zealand since 1882.

● **Leucoptera** Hübner, 1825, p. 426. Type species *Leucoptera spartifoliella* Hübner, according to Bradley & Carter (1982, p. 1).

*spartifoliella* Hübner, [1810 – 20 June 1813], pl. 49  
fig. 335 (*Tinea*)  
Europe; adventive to California.  
New Zealand: first recorded from Taupo TO in 1950; widespread on *Cytisus* since 1979.  
Note. This is the "Opostega sp." of Dugdale (1974, p. 138, fig. 21).


● **Stegommata** Meyrick, 1880a, p. 171. Type species *Stegommata leptomitella* Meyrick, by subsequent designation (as in Clarke 1955, p. 184).

*leptomitella* Meyrick, 1880a, p. 172 (*Stegommata*)  
Sydney N.S.W., E. Meyrick; LT ♀ here designated, labelled "Sydney N.S. Wales 30.1.79", "Lyonetia leptomitella Meyr. 8/8 E. Meyrick det. in Meyrick Coll.", BMNH.  
New Zealand: adventive, on Australian *Hakea* species (Proteaceae).

—◎—

**Superfamily GELECHIOIDEA**  
(in the sense of Common 1970)

**Family GELECHIIDAE**  
(in the sense of Common 1970)



**Gelechiidae**  
(151) *Anisoplaca ptyoptera* Meyrick

• **Anarsia** in the sense of Meyrick (1904, p. 415)

**trichodeta** Meyrick, 1904, pp. 415–416 (*Anarsia*),

in a broad sense

Mount Gambier S.A., E. Meyrick; ST ♂ labelled “Anarsia trichodeta Meyr. 1/1 E. Meyrick in Meyrick Coll.”, “Mt Gambier S. Australia 14/11/82”, BMNH.

New Zealand: adventive. On *Acacia* and *Albizzia* foliage and fruits, especially in ND and AK, since 1964.

Note. New Zealand and Toowong (Qld) specimens differ from Meyrick's syntype in having the hindwings strongly infuscate towards the apex, and no yellow scales on the frons. New Zealand specimens agree well with Queensland specimens in the series at BMNH.

• **Anisoplaca** Meyrick, §1885i, p. 590; 1886b, p. 170.

Type species *Anisoplaca ptyoptera* Meyrick, by original monotypy.

**achyrota** Meyrick, §1885i, p. 590; 1886b, p. 170  
(*Gelechia*)

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 23/12/82”, “Anisoplaca achyrota Meyr. 4/14 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, pp. 258–259, but not pl. xxviii fig. 1.

Note. The ♂ genitalia illustrated by Philpott (1927j, p. 350, fig. 3) and Hudson (1928, pl. xxviii fig. 1) refer to an undescribed species.

**acrodactyla** Meyrick, 1907c, pp. 117–118 (*Gelechia*)

Invercargill SL, A. Philpott; LT ♂ here designated, labelled “Invercargill New Zealand AP .06”, “Anisoplaca acrodactyla Meyr. 1/3 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 258, pl. xxviii fig. 3.

Note. Philpott (1927j, p. 350, fig. 4) figures ♂ genitalia of a specimen collected at the same time and place as the series he sent to Meyrick.

**cosmia** Bradley, 1956b, p. 156 (*Anisoplaca*)  
Norfolk Island, J.D. Bradley; HT ♂ designated by Bradley, BM genitalia slide no. 3787 ♂, BMNH.  
New Zealand: adventive, on *Lagunaria patersonia* ('Norfolk Island hibiscus'); larvae feed in shoots and fruits.

**fraxinea** Philpott, 1928a, p. 364 (*Anisoplaca*)  
Flora River NN, A. Philpott; HT ♂ designated by Philpott, abdomen missing, NZAC.  
Hudson 1939, p. 440, pl. lviii fig. 20.

**ptyoptera** Meyrick, §1885i, p. 591; 1886b, p. 171  
(*Anisoplaca*)  
Christchurch MC, R.W. Fereday; HT ♂ unique, BM genitalia slide no. 3796 ♂, BMNH.  
Hudson 1928, p. 259, pl. xxxviii fig. 1.  
Note. Philpott (1927j, p. 350, fig. 2) figures ♂ genitalia of a Christchurch specimen.

Also 1 undescribed species (NZAC).  
Note. Philpott (1927j, p. 350, fig. 3) erroneously regarded this species as *achyrota*. Hudson (1928, pl. xxviii fig. 1) illustrated this species as *achyrota*.

• **Athrips** of authors, but not Billberg 1820 (Sattler 1978, p. 58)

**zophochalca** Meyrick, 1918a, p. 133 (*Epithectis*)  
Auckland AK, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 253, pl. xxxviii fig. 2, as *Epithectis zophochalca*.

**transversella** Hudson, 1939, p. 438 (*Epithectis*).  
New synonymy.

Auckland Domain AK, C.E. Clarke; HT ♀ unique, AMNZ.

Hudson 1939, p. 438, pl. lxii fig. 23.  
Note. *A. zophochalca* is sexually dimorphic for wing pattern, males having the white band on the forewings indistinct. Fresh specimens have the thorax covered in coppery scales. Larvae bore the terminal shoots of *Carmichaelia* species (Fabaceae).

• **Biloba** Janse, 1954, p. 301. Type species *Gelechia (Brachmia) subsecivella* Zeller, by original designation.

**subsecivella** Zeller, 1852, p. 113 (*Gelechia (Brachmia)*)

South Africa.  
New Zealand: adventive (Philpott 1924c, p. 666).  
Hudson 1939, p. 438, as *Stomopteryx subsecivella*, with synonymy.

**columbina** Philpott, 1928a, p. 364, fig. 2 (*Stomopteryx*), replacement name for *Stomopteryx simplicella* in the sense of Philpott (1924c, p. 666) but not Walker (1864, p. 1024). Synonymised by Meyrick in Hudson (1928, footnote, pl. li).  
Nelson NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 253, pl. li fig. 21, as *Stomopteryx simplicella*.

• **Brachmia** in the sense of Meyrick (1925b, p. 248)

**Species near phryganitis** Meyrick, 1911e, p. 722  
(*Brachmia*)

Adventive. Recorded from various localities in AK since 1972.

Note. Resembles in genitalia BMNH specimens from south-eastern Australia under the name *B. phryganitis*, which was described from Maskeliya, Sri Lanka.

• **Chrysoesthia** Hübner, 1825, p. 422. Type species *Tinea zinckenella* Hübner, 1813, by subsequent designation (Meyrick 1925, p. 40); Europe.

**durella** Fabricius, 1775, p. 666 (*Tinea*)

"Habitat Hafniae".

New Zealand: adventive (Dumbleton 1964, p. 24), on Chenopodiaceae; widespread.

**hermanella** of authors but not Fabricius, 1974, p. 324 (Karsholt & Nielsen 1976b, p. 246) (*Chrysoesthia*).

• **Epiphthora** Meyrick, 1888e, p. 77. Type species *Epiphthora melanombra* Meyrick, by original monotypy.

Note. Meyrick (1923, p. 165) synonymised *Epiphthora* with the now restricted *Apatetris* Staudinger. Janse (1951, p. 232) regards *Epiphthora* as valid (Sattler 1973, p. 198).

**melanombra** Meyrick, 1888e, p. 77 (*Epiphthora*)

Christchurch MC, reared by R.W. Fereday; LT ♂ here designated, labelled "Christchurch New Zealand RWF bred /85", "Apatetris melanombra Meyr. 2/4 E. Meyrick det. in Meyrick Coll.", abdomen missing, BMNH.

Hudson 1928, p. 252, pl. xxxviii fig. 3, as *Apatetris melanombra*.

**sparsa** Philpott, 1918, p. 128 (*Gelechia*). Synonymised by Meyrick (1923, p. 165).

Dunedin DN, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 252, as synonym of *melanombra*.

**nivea** Philpott, 1930b, p. 7 (*Apatetris*) new combination

Auckland AK, C.E. Clarke; HT ♂ designated by Philpott, abdomen missing, AMNZ.

Hudson 1939, p. 438, pl. lviii fig. 1.

• **Hierodoris** of authors, but not Meyrick (1912d)

**insignis** Philpott, 1926a, p. 397 (*Hierodoris*?)

Mount Arthur Tableland NN, S. Lindsay; HT ♀ designated by Philpott, CMNZ.

Hudson 1928, p. 306, pl. lii fig. 31, in "Glyphipterygidae". Note. This species is transferred to Gelechiidae; the larva mines under the tomentum of leaves of *Celmisia* species (Asteraceae).

Also 2 undescribed species (NZAC).

• **Isochasta** Meyrick, §1885i, p. 590 (as *Tsochasta*); 1886b, p. 163. Type species *Isochasta paradesma* Meyrick, by original monotypy.

Note. A case for *Tsochasta* being an incorrect original spelling will have to be submitted to the ICZN.

**paradesma** Meyrick, §1885i, p. 590 (*Tsochasta*); 1886b, pp. 163–164 (*Isochasta*)

Invercargill SL, E. Meyrick; HT ♂ unique, BMNH. Hudson 1928, p. 253, pl. xxvii fig. 15, as *Aristotelia paradesma*.

• **Kiwaia** Philpott, 1930a, p. 248. Type species *Kiwaia jeanae* Philpott, by original designation.

**Zeempista** Povolny, 1974, p. 414. Type species *Gelechia cheradias* Meyrick, by original designation. Synonymised by Sattler (1988, p. 233).

Note. Povolny (1977, pp. 435–437) gives additional information and biogeography.

**aerobatis** Meyrick, 1924a, p. 204 (*Gelechia*)

Mount Arthur NN, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 4000' 15/1/86", "Gelechia aerobatis Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 257, pl. xlvi fig. 24.

**brontophora** Meyrick, §1885i, p. 590; 1886b, p. 168 (*Gelechia*)

[Port Hills], Christchurch MC, E. Meyrick; HT ♂ unique, BM genitalia slide no. 16620, BMNH.

Hudson 1928, p. 255, pl. xxvii fig. 11, as *Phthorimaea brontophora*.

**caerulea** Hudson, 1925, p. 221 (*Gelechia*)

Waiho Gorge WD, C.E. Clarke; HT ♂ unique, not located in NMNZ.

Hudson 1928, p. 258, pl. lii fig. 5.

**calaspidea** Clarke, 1934, p. 14 (*Gelechia*)

Flat Top Mountain, Hunter Mountains FD, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.

Hudson 1939, p. 440, pl. lvii fig. 20.

**cheradias** Meyrick, 1909a, p. 12 (*Gelechia*)

Invercargill SL, A. Philpott; LT ♂ designated by Povolny (1974, p. 421), BMNH.

Hudson 1928, p. 255, pl. xxvii fig. 9.

**contraria** Philpott, 1930b, p. 8 (*Gelechia*)

Waiho Gorge WD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 439, pl. lviii fig. 2.

» Gelechiidae, *Kiwaia*

**dividua** Philpott, 1921, p. 340 (*Gelechia*)  
Paradise, Lake Wakatipu OL, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ.  
Hudson 1928, p. 257, pl. xlvi fig. 36.

**eutrybathra** Meyrick, 1931b, p. 368 (*Gelechia*)  
Porter River MC, S. Lindsay; HT ♂ designated by Lindsay(?), CMNZ.  
Hudson 1939, p. 440, pl. lviii fig. 19.

**glaucoterna** Meyrick, 1911b, p. 63 (*Gelechia*)  
Invercargill SL, A. Philpott; LT ♂ here designated, labelled "Invercargill New Zealand AP 15.11.'08", "Phthorimea glaucoterna Meyr. 1/3 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 256, pl. xxvii fig. 19, as *Phthorimaea glaucoterna*.

Note. Povolny (1974, p. 416) synonymised this with *brontophora*, ignoring the external differences, particularly the hindwing patch of modified scales (present also in *Gelechia caerulea*, q.v., but absent from *brontophora*). *Gelechia glaucoterna* is removed from this synonymy.

**heterospora** Meyrick, 1924a, p. 204 (*Phthorimaea*)  
Mount Ruapehu TO, G.V. Hudson; LT ♂ here designated, labelled "Mt Ruapehu New Zealand GVH 4000' 1.22", "Phthorimaea heterospora Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 256, but not pl. 1 fig. 12, as *Phthorimaea heterospora*.

Note. This species superficially agrees well with *materneme* Povolny and *neglecta* Philpott; on external characters it is an obvious *Kiwaia*, and is consequently placed in this genus.

**hippeis** Meyrick, 1901, pp. 573–574 (*Gelechia*)  
Christchurch MC, R.W. Fereday; LT ♀ here designated, labelled "Gelechia hippeis Meyrick" in Meyrick's handwriting, "Christchurch New Zealand RWF/01", CMNZ.

Hudson 1928, p. 256, pl. xxvii fig. 20.

Note. The BMNH PLT ♂ has the abdomen missing.

**jeanae** Philpott, 1930a, p. 249 (*Kiwaia*)  
Birdlings Flat MC, Jean Lindsay; HT ♂ designated by Philpott, CMNZ.  
Hudson 1939, pp. 437–438, pl. lvii fig. 9 and 10.

**lapillosa** Meyrick, 1924a, p. 203 (*Gelechia*)  
Mount Ruapehu TO, G.V. Hudson; LT ♂ here designated, labelled "Mt Ruapehu New Zealand GVH 4000' 1.22", "Gelechia lapillosa Meyr. 2/4 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 258, pl. xxxi fig. 17.

**lenis** Philpott, 1929a, p. 302 (*Gelechia*)  
Lake Pukaki MK, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 439, pl. lviii fig. 3.

**lithodes** Meyrick, §1885i, p. 590; 1886b, p. 170 (*Gelechia*)  
Arthur's Pass NC/WD, E. Meyrick; HT ♂ unique, right forewing, head, and abdomen missing, BMNH.  
Hudson 1928, p. 258, pl. xxviii fig. 2 (from a North Island specimen).

**materneme** Povolny, 1974, p. 426 (*Zeempista*)  
New River SL, A. Philpott; HT ♂ designated by Povolny, slide no. Au21, NZAC.  
Note. See *Z. heterospora*, above.

**monophragma** Meyrick, 1885i, p. 590; 1886b, pp. 169–170 (*Gelechia*)  
Wellington WN, E. Meyrick; LT ♂ here designated, labelled "Wellington New Zealand 30/12/79", "Gelechia monophragma Meyr. 2/12 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 257, pl. xxviii fig. 4 and 5.

**neglecta** Philpott, 1924c, pp. 665–666 (*Gelechia*)  
Cobb Valley NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 258, pl. li fig. 14.

**parapleura** Meyrick, §1885i, p. 590; 1886b, pp. 168–169 (*Gelechia*)  
Bealey River [bed] MC, E. Meyrick; LT ♂ here designated, labelled "Bealey R. New Zealand 30/1/83", "Gelechia parapleura Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 257, pl. xxviii fig. 6.

**parvula** Philpott, 1930b, pp. 7–8 (*Gelechia*)  
Hope Arm, Lake Manapouri FD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 440, pl. lviii fig. 18.

**pharetria** Meyrick, §1885i, p. 590 (as *phaxetria*); 1886b, p. 169 (*Gelechia*)  
Arthur's Pass NC/WD, E. Meyrick; LT ♂ here designated, labelled "Arthurs Pass New Zealand 2500' 23/1/83", "Gelechia pharetria Meyr. 2/9 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 257, pl. xxviii fig. 25 and 26.

**plemochoa** Meyrick, 1916b, p. 415 (*Phthorimaea*)  
Otira River WD, G.V. Hudson; LT ♂ designated by Povolny (1974, p. 424), BMNH.  
Hudson 1928, p. 256 (but none of the type series resembles the specimen figured in pl. xxviii fig. 23 and 24).  
Note. In Povolny (1974, p. 424) for "CVH 12.13" read 'GVH 12.14'.

**pumila** Philpott, 1928c, p. 182 (*Gelechia*)  
Yaldhurst MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.  
Hudson 1939, p. 439, pl. lviii fig. 4 (from the HT).

**schematica** Meyrick, §1885i, p. 590; 1886b, p. 168  
(*Gelechia*)

Bealey River [bed] MC, E. Meyrick; LT ♂ here designated, labelled "Bealey R. New Zealand 21/1/83", "*Gelechia schematica* Meyr. 2/6 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 257, pl. xxviii fig. 7.

**thyraula** Meyrick, §1885i, p. 590; 1886b, p. 167  
(*Gelechia*) new combination

Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 5/2/83", "*Phthorimaea thyraula* Meyr. 2/6 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 255, possibly pl. xxvii fig. 10, as *Phthorimaea thyraula*.

**quieta** Philpott, 1927a, p. 706 (*Phthorimaea*). New synonymy.

Bottle Lake [Christchurch MC, S. Lindsay]; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 255, pl. lii fig. 7, as species.

**pulverea** Philpott, 1928c, p. 181 (*Phthorimaea*). Synonymised (with *quieta*) by Povolny (1974, p. 424).

Hapuka River mouth KA, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 439, pl. lviii fig. 17, as *Phthorimaea quieta*.

● **Megacraspedus** of authors, but not Zeller (1839, p. 189)

**calamogonus** Meyrick, §1885i, p. 589; 1886b, p. 163  
(*Megacraspedus*)

Christchurch MC, R.W. Fereday; LT ♂ here designated, labelled "Christchurch New Zealand RWF 1/11/68", "*Megacraspedus calamogonus* Meyr. 3/6 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pp. 252–253, pl. xxvii fig. 12 and 13.

Also 1 undescribed species (NZAC).

● **Phthorimaea** Meyrick, 1902b, pp. 103–104. Type species *Bryotropha operculella* Zeller, by original designation.

**operculella** Zeller, 1873, p. 262 (*Bryotropha*)  
Cosmopolitan.

New Zealand: adventive, throughout.

Hudson 1928, pp. 254–255, pl. xxvii fig. 18.

Note. Meyrick (1886b, p. 167) reported this in New Zealand as *Gelechia solanella* Boisduval.

**sedata** Butler, 1880, p. 560 (*Gelechia*)  
[Blenheim] MB, W. Skellon; HT unique, BMNH.

Not recorded by Hudson.

Note. This reflects the current position in BMNH (K.R. Tuck, pers. comm.).

● **Sitotroga** Heinmann, [1870], p. 287. Type species *Alucita cerealella* Olivier, by original monotypy.

**cerealella** Olivier, 1789, p. 121 (*Alucita*)  
Cosmopolitan.

New Zealand: adventive, North Island only.

Hudson 1928, p. 259, pl. xl fig. 10.

Note. Meyrick (1923, p. 164) gives the first record, received by Hudson from Levin WN.

● **Symmetrischema** Povolny, 1967, pp. 53–55. Type species *Phthorimaea plaesiosema* Turner, by original designation.

**plaesiosema** Turner, 1919, p. 126 (*Phthorimaea*)

Sydney N.S.W., Goldfinch; HT in AMSA.

Hudson 1939, p. 439, not figured.

**melanoplinta** Meyrick, 1926a, p. 279 (*Phthorimaea*). Synonymised by Hudson (1939, p. 439).

Hawkes Bay HB, Miller; LT ♂ here designated, labelled "Hawkes Bay New Zealand GVH bred 1.24", "ex stem tomato", "*Phthorimea melanoplinta* Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 256, pl. xxviii fig. 21.

Notes. Povolny (1967, pp. 53–55) described *Symmetrischema* and gave synonymies; Povolny (1977, p. 433) synonymised *Phthorimaea aquilina* Meyrick from Peru with *S. plaesiosema*. The genus and the species are of Neotropical origin.

● **Thiotricha** Meyrick, §1885i, p. 590 (as *Thistricha*); 1886b, p. 164. Type species *T. thorybodes* Meyrick, by subsequent designation of Meyrick (1915b, p. 207).

Note. A case for *Thistricha* being an incorrect original spelling will have to be submitted to the ICZN.

**lindsayi** Philpott, 1927d, p. 84 (*Thiotricha*)

Glentui NC, S. Lindsay; HT ♂ designated by Philpott, abdomen missing, CMNZ.

Hudson 1939, p. 438, not figured.

**oleariae** Hudson, 1928, p. 254 (*Thiotricha*)

Breaker Bay WN, Stella Hudson; LT ♂ here designated, labelled "712a", Hudson Collection, NMNZ.

Hudson 1928, p. 254, pl. xlvi fig. 7.

**tetraphala** Meyrick, §1885i, p. 590 (*Thistricha*); 1886b, p. 164 (*Thiotricha*)

Dunedin DN, E. Meyrick; HT ♂ unique, abdomen in gelatin capsule, BMNH.


Hudson 1928, p. 254, pl. xxvii fig. 17.

**thorybodes** Meyrick, §1885i, p. 590 (*Thistricha*); 1886b, pp. 164–165 (*Thiotricha*)

Riccarton Bush MC, E. Meyrick; LT ♂ designated by Clarke (1969, p. 447), BMNH.

Hudson 1928, p. 254, pl. xxvii fig. 16.

**Family BATRACHEDRIDAE**  
(in the sense of Nielsen & Common, in press)



---

**Batrachedridae**  
(152) *Batrachedra arenosella* (Walker)

---

• ***Batrachedra*** Stainton, in the sense of Hodges (1966, p. 594, fig. 1, 2, 37–39, 94, 106, and 110)

***agaura*** Meyrick, 1901, p. 579 (*Batrachedra*)  
[below Flora Saddle], Mount Arthur NN, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 19/1/86", "Batrachedra agaura Meyr. 14/16 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 304, pl. xxxv fig. 2 (a poor representation; Hudson 1950, pl. v fig. 6 is better, but most specimens are browner than Hudson indicates).

***arenosella*** Walker, 1864b, p. 857 (*Gracillaria*)  
"[Auckland AK] New Zealand, presented by Col. Bolton"; syntype series of 3 ("a–c") not in collection [H. Stringer, April 1932; JSD, November 1980], BMNH.  
Hudson 1928, p. 304, pl. xxxv fig. 12.

Note. None of the BMNH specimens are topotypes; the Meyrick Collection has material from Christchurch and Wellington. The New Zealand ♀♀ (BMNH genitalia slides 8081 and 19951) differ from a Queensland ♀ (genitalia slide 8084) in limen : ductus bursae length ratio.

***astricta*** Philpott, 1930b, p. 14 (*Batrachedra*)  
Opoho, Dunedin DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 442, pl. lvii fig. 22 (figured from the type series).

***eucola*** Meyrick, 1889b, pp. 180–181 (*Batrachedra*)  
Bealey River NC, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 304, pl. xxxiv fig. 8.

***filicicola*** Meyrick, 1917a, p. 247 (*Batrachedra*)  
Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 11.15", "Batrachedra filicicola Meyr. 2/5 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 304, pl. xl fig. 11 and 12.

***litterata*** Philpott, 1928, p. 370 (*Batrachedra*, as "Batrachedra")  
Greymouth WD, A.J. Turner; HT ♀ designated by Philpott, NZAC.  
Hudson 1939, p. 442, pl. lvii fig. 14.

***psathyra*** Meyrick, 1889b, p. 181 (*Batrachedra*)  
Hamilton WO, E. Meyrick; LT ♀ here designated, labelled "Hamilton New Zealand 18.1.80", "Batrachedra psathyra Meyr. 6/14 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 303, pl. xxxv fig. 13, pl. xliv fig. 8, as *B. psathyra*.


***tristictica*** Meyrick, 1901, p. 579 (*Batrachedra*)  
Makatoku HB, E. Meyrick; LT ♀ here designated, labelled "Makatoku New Zealand 8/3/83", "Batrachedra tristictica Meyr. 2/3 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 304; 1939, p. 442, pl. lvii fig. 15.

Also 1 undescribed species, from *Asplenium* sori (NZAC).

—◎—

## Family BLASTOBASIDAE

(in the sense of Hodges 1978, p. 7)



---

### Blastobasidae (153) *Blastobasis tarda* Meyrick

---

#### Subfamily BLASTOBASINAE (in the sense of Hodges 1978, p. 7)

- ***Blastobasis*** in the sense of Meyrick (1902d, pp. 168–169)

*tarda* Meyrick, 1902d, p. 170 (*Blastobasis*)

Sydney N.S.W., E. Meyrick; LT ♂ selected by I.F.B. Common and here designated, labelled "Sydney N.S. Wales 25.10.79", "Blastobasis tarda Meyr. 1/8 E. Meyrick det. in Meyrick Coll.", BMNH.

New Zealand: adventive; noted since 1973. Specimens in NZAC from ND and AK. On stored fruit and old or mummified fruit on trees.

Note. PLT 5/8 ("Sydney ... 8.8.79") is a good match with New Zealand examples.

—⊗—

#### Subfamily SYMMOCINAE (in the sense of Hodges 1978, p. 7)

- ***Oegoconia*** Stainton, 1854, p. 162. Type species *Recurvaria quadripuncta* Haworth, 1828, by monotypy.

*caradjai* Popescu-Gorj & Căpuse, 1965, p. 389 (*Oegoconia*).

New Zealand: adventive; recorded from NN (Meyrick 1911b, p. 69) as *Symmoca quadripuncta* [of authors]. Hudson 1928, p. 295, pl. xxxii fig. 12, as *Symmoca quadripuncta*. Philpott 1927j, p. 350, fig. 1, as *Oegoconia quadripuncta* [of authors].

Note. Agassiz (1982) gives a comprehensive diagnostic guide to *Oegoconia* species.

—⊗—

## Family COLEOPHORIDAE

(in the sense of Hodges 1978, pp. 7 and 9, in part)

### Subfamily COLEOPHORINAE

(in the sense of Hodges 1978, p. 7)

- ***Coleophora*** Hübner, 1822, p. 67. Type species *Tinea ornatipennella* Hübner, 1796; Europe.

*spissicornis* Haworth, 1828, p. 537 (*Porrectaria*) [Europe].

New Zealand: adventive; on clovers, throughout.

Note. New Zealand specimens agree externally and generally with material from Merton, Norfolk (the type locality). Consistent differences were seen between these and *hieronella* Zeller (southern France) and *mayrella* Zeller (Corsica), both treated as synonyms of *spissicornis* by Patzak (1974).

Hudson 1939, p. 460, pl. lviii fig. 21.

*trifolii* Curtis, 1832, folio 391 (*Tinea*) Europe.

New Zealand: adventive; on clovers, throughout.

Note. Bradley (1967, pp. 45–46) indicated – rather vaguely – the misapplication of *frischella*; Robinson & Nielsen (1983, pp. 211 and 212–213, fig. 20–23) verified this and illustrated differences. New Zealand specimens have the characters of *trifolii*.

*frischella* in the sense of authors, not of Linnaeus, 1758, p. 541 (*Phalaena Tinea*, sp. 389 [289]) [Europe].

*alcyonipenella* Kollar, 1832, p. 99 (*Ornix*). Synonymised by Patzak (1974, p. 245). [Europe].

Not mentioned by Hudson.

Note. New Zealand references used this name until 1974.

*versurella* Zeller, 1849, p. 352 (*Coleophora*)


Europe.

New Zealand: adventive, on Amaranthaceae and Chenopodiaceae; present throughout, including Antipodes Islands.

*Coleophora "sp. indet."* Dugdale, 1971b, p. 77  
Antipodes Islands, P.M. Johns; unique ♀, NZAC.

—⊗—

**Family COSMOPTERIGIDAE**  
(in the sense of Hedges 1978, pp. 11–16)



**Cosmopterigidae**  
(154) *Pyroderces apparitella* (Walker)

● ***Circoxena*** Meyrick, 1916b, p. 418. Type species  
*Circoxena ditrocha* Meyrick, by original monotypy.

***ditrocha*** Meyrick, 1916b, p. 419 (*Circoxena*)  
Wainuiomata WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, pp. 331–332, pl. xxviii fig. 19, in Plutellidae. Dugdale 1973b, p. 1020, in Cosmopterigidae.

● ***Limnaecia*** Stainton, 1851, p. 4. Type species  
*Limnaecia phragmitella* Stainton, by original monotypy.

***phragmitella*** Stainton, 1851, p. 4 (*Limnaecia*)  
[Europe].  
New Zealand: self-adventive; on *Typha*, throughout.  
Hudson 1928, p. 302, not figured.

● ***Microcolona*** Meyrick, 1897a, p. 371. Type species  
*M. limodes* Meyrick, by subsequent designation  
(Meyrick 1915b, p. 209).

***characta*** Meyrick, 1897a, p. 374 (*Microcolona*)  
Sydney N.S.W., E. Meyrick; LT ♂ selected by I.F.B.  
Common and here designated, labelled "Sydney  
N.S.W. 23/8/79", "Microcolona characta Meyr. 1/7 E.  
Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 303, pl. xxviii fig. 8.  
Note. Possibly *M. limodes* alone is present in New  
Zealand. A *characta* PLT from Nelson NN and the 8  
additional specimens placed under *characta* from  
Wellington WN (sent by Hudson) differ from the  
Australian material in their shorter pecten on the  
antennal scape and their hind femoral and tibial black  
scale patches.

***limodes*** Meyrick, 1897a, pp. 372–373 (*Microcolona*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂  
here designated, labelled "Christchurch N. Zealand  
9/3/82" "Microcolona limodes Meyr. 2/3 E. Meyrick  
det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 303, not figured.

● ***Pyroderces*** of authors, partly of Hedges (1978, pp.  
46–47)

***Syntomactis*** Meyrick, 1889b, p. 173. Type  
species *Gelechia deamatella* Walker, by original  
monotypy. Synonymised by Meyrick (1924c, p. 91).

***aellotricha*** Meyrick, 1889b, p. 175 (*Proterocosma*)  
Hamilton WO, E. Meyrick; LT ♀ here designated, labelled  
"Hamilton New Zealand 17/1/80", "Pyroderces ael-  
otricha Meyr. 1/3 E. Meyrick det. in Meyrick Coll.",  
abdomen in capsule, BMNH.

Hudson 1928, p. 301, not figured; 1939, p. 441, pl. lviii  
fig. 5.

Note. Present also in eastern Australia, and very similar  
to *P. badia* Hedges.

***anarithma*** Meyrick, 1889b, pp. 175–176  
(*Proterocosma*)

[New Plymouth] "Taranaki" TK, E. Meyrick; LT ♂ here  
designated, labelled "Taranaki New Zealand 26/2/83",  
"Pyroderces anarithma Meyr. 2/6 E. Meyrick det. in  
Meyrick Coll.", BMNH.

Hudson 1928, p. 302, pl. xxviii fig. 16.

***apparitella*** Walker, 1864b, p. 1027 (*Gelechia*)  
[Auckland AK], D. Bolton; HT ♀ unique, BMNH.  
Hudson 1928, p. 301, pl. xxviii fig. 22.

Note. The BMNH collection has *apparitella* in *Labdia*  
Walker (type species *L. deliciosa* Walker, from east-  
ern Australia), but the genitalia of the two species have  
markedly different specialisations.

***deamatella*** Walker, 1864a, p. 654 (*Gelechia*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, head damaged,  
BMNH.

Hudson 1928, p. 303, pl. xxviii fig. 18, as *Syntomactis*  
*deamatella*.

Note. Meyrick's synonymy of *Syntomactis* with *Pyro-  
derces* is contained in his discussion of *Leptozestis*  
Meyrick (Meyrick 1924c, p. 91).

Also 1 undescribed species (NZAC).

● ***Thectophila*** Meyrick, 1927a, p. 701. Type species  
*T. acmotypa* Meyrick, by original monotypy.

***acmotypa*** Meyrick, 1927a, p. 701 (*Thectophila*)  
Arthur's Pass NC/WD, G.V. Hudson; HT ♀ unique,  
BMNH.

Hudson 1928, p. 302, pl. lii fig. 18.



**Family LECITHOCERIDAE**  
(in the sense of Gozmany 1978)

• **Lecithocera** Herrich-Schaeffer, 1853, p. 45. Type species *Carcina luteicornella* Zeller, 1839, by monotypy.

**micromela** Lower, 1897, p. 55 (*Lecithocera*)  
Australia.  
New Zealand: adventive; recorded since 1926, ND-NN.  
Hudson 1939, p. 441, pl. lviii fig. 14.  
Note. Philpott (1928a, p. 364) gives early records.



**Family MOMPHIDAE**  
(in the sense of Riedl 1969, p. 639)


• **Zapryrastra** Meyrick, 1889b, p. 172. Type species *Zapryrastra calliphana* Meyrick, by original monotypy.

**calliphana** Meyrick, 1889b, p. 172 (*Zapryrastra*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 18/2/82", "Zapryrastra calliphana Meyr. 16/2 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, pp. 302–303, pl. xxviii fig. 20, in Cosmopterigidae.  
Note. In Meyrick's Diary of Captures this species is first recorded as "Hel. chrysocarpa". Dugdale (1971c, pp. 67–69) gave life-history details, and misspelled the generic name as *Zapryasta*.

**stellata** Philpott, 1931, p. 31 (*Elachista*) new combination


Pembroke [Wanaka] OL, C.E. Clarke; HT ♂ unique, AMNZ.  
Hudson 1939, p. 459, pl. ix fig. 23, as *Elachista stellata*, in Elachistidae.

Also 1 undescribed species on *Haloragis*, Haloragidaceae (NZAC).



**Momphidae**  
(155) *Zapryrastra calliphana* Meyrick

**Family OECOPHORIDAE**  
(in the sense of Hedges 1974, pp. 7–9)  
**Subfamily DEPRESSARIINAE**  
(in the sense of Hedges 1974, p. 9)



**Oecophoridae: Depressariinae**  
(156) *Proteodes carnifex* (Butler)

• **Cryptolechia** of authors, in part, but not of Zeller (1852, p. 106)

**rhodobapta** Meyrick, 1923, p. 166 (*Cryptolechia*)  
Takapuna AK, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 294, pl. xlvi fig. 25.

**semnodes** Meyrick, 1911b, p. 75 (*Cryptolechia*) new combination

Mount Arthur Tableland NN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 294, pl. xxxii fig. 22.  
Note. Philpott (1927f, p. 110, fig. 46, and p. 113) queries the generic position of *semnodes*, the only species in the group with a gnathos.

• **Eutorna** Meyrick, 1889b, p. 157. Type species *E. caryochroa* Meyrick, by subsequent designation (Meyrick 1915b, p. 220).

**Phyzanica** Turner, 1917, p. 117. Type species *P. tapinopa* Turner, 1917, p. 117 (= *Eutorna pelogenes* Meyrick, 1906). Synonymised by Meyrick (1922d, p. 182, 183).

Note. This synonymy, listed in Gaede (1939, pp. 364–365), is apocryphal (I.F.B. Common, pers. comm.).

**caryochroa** Meyrick, 1889b, p. 157 (*Eutorna*)  
["reserved bush and forest"], Dunedin DN, E. Meyrick; LT ♂ here designated, labelled "Dunedin New Zealand 6/1/80", "Eutorna caryochroa Meyr. 1/12 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 295, pl. xxxii fig. 9.

**inornata** Philpott, 1927d, p. 88 (*Eutorna*)  
Seaward Moss, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, head and abdomen missing.  
NZAC.

» Oecophoridae, *Eutorna inornata*

Hudson 1939, p. 453, pl. lix fig. 19.

Note. The HT has forewing veins *CuA1* and *CuA2* short-stalked; the AT ♀ has these veins approximated at their base on the discal cell. A series of a pale *Eutorna* species from Miranda AK shows a similar range, and Meyrick's HT ♂ *symmorphia* has the same *Cu* condition as the *inornata* AT ♀.

***phaulocosma*** Meyrick, 1906, pp. 41 (key) and 45 (*Eutorna*)

Mount Wellington Tasmania, E. Meyrick; LT ♂ selected by I.F.B. Common and here designated, labelled "Mt Wellington Tasmania 1.2.82", "Eutorna phaulocosma Meyr. 3/3 E. Meyrick det. in Meyrick Coll.", palpi missing, BM genitalia slide no. 18911, BMNH.

New Zealand: adventive; throughout, on introduced *Rubus* species.

Not mentioned by Hudson.

***symmorphia*** Meyrick, 1889b, p. 158 (*Eutorna*)

[sandhills near] Kaiapoi NC, E. Meyrick; LT ♂ here designated, labelled "Kaiapoi New Zealand 31/3/82", "Eutorna symmorphia Meyr. 4/11 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 295, pl. xxxii fig. 8.

Note. Forewing *CuA1* and *CuA2* stalked in PLT ♂ 5/11 (Hamilton WO) and 1/11 (Whangarei ND; very pallid specimen); see *inornata*, above.

• ***Heliostibes*** in the sense of Meyrick (1888e, p. 82), not of Zeller (1874)

***vibratrix*** Meyrick, 1927a, p. 702 (*Heliostibes*)

Mount Arthur NN, G.V. Hudson; HT ♀ unique, BMNH. Hudson 1928, p. 307, pl. 1 fig. 21, in *Glyptopterigidae*.

Note. The HT ♀ lacks ocelli, and superficially resembles *Phaeosaces semnodes* Meyrick.

• ***Nymphostola*** Meyrick, 1883a, pp. 424 (key) and 491–492. Type species *Cryptolechia galactina* Felder & Rogenhofer, by original monotypy.

***galactina*** Felder & Rogenhofer, 1875, pl. cxl fig. 34 (*Cryptolechia*)

[Nelson NN, T.R. Oxley]; HT ♀ designated by Felder, represented by 4 wings and meso/metathorax, BMNH. Hudson 1928, pp. 291–292, pl. xxv fig. 20.

• ***Proteodes*** Meyrick, 1883a, pp. 425 (key) and 492 (description). Type species *Cryptolechia carnifex* Butler, by original monotypy.

***carnifex*** Butler, 1877, p. 406 (*Cryptolechia*)

[Castle Hill Station MC], J.D. Enys; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 292, pl. xxxii fig. 23–25.

***rufosparsa*** Butler, 1877, p. 406 (*Cryptolechia*).

Synonymised by Meyrick (1883a, p. 493).

[Castle Hill Station MC], J.D. Enys; HT ♀ unique, abdomen glued on back to front, BMNH.

***clarkei*** Philpott, 1926a, pp. 396–397 (*Proteodes*)

Hunter Mountains FD, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 293, pl. lii fig. 27.

Note. Philpott (1931, p. 33) described the brachypterous female.

***melographa*** Meyrick, 1927a, pp. 700–701 (*Proteodes*)

Mount Arthur NN, 4000 ft, S. Woodward; HT ♂ unique, BMNH.

Hudson 1928, p. 292, pl. lii fig. 20.

***varia*** Philpott, 1928a, p. 368 (*Proteodes*). Synonymised by Meyrick (1936, p. 283)

Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 452, as synonym.

***profunda*** Meyrick, 1905, p. 236 (*Proteodes*)

Mount Holdsworth WN, 2000 ft, G.V. Hudson; HT ♂ unique, right forewing missing, BMNH.

Hudson 1928, p. 293, pl. xxv fig. 42.

***smithi*** Howes, 1946, p. 146 (*Proteodes*)

Homer FD, T.R. Smith; HT ♂ unique, designated by Howes, NMNZ.

Not mentioned by Hudson.

Also 2 undescribed species (B. Patrick Collection, Dunedin).



### Subfamily OECOPHORINAE

(in the sense of Hodges 1974, pp. 9 and 97)

**Group A:** Oecophorinae with ocelli, *Imma*-like in appearance. Philpott (1927i, pp. 346–347) describes and illustrates genitalia of 5 species.

• ***Coridomorpha*** Meyrick, 1914a, p. 111. Type species *C. stella* Meyrick, by original monotypy.


***stella*** Meyrick, 1914a, p. 111 (*Coridomorpha*)

Kauri Gully, Auckland AK, Stella Hudson; HT ♀ unique, labelled as LT by J.B. Heppner (1976), BMNH.

Hudson 1928, p. 305, pl. xxxii fig. 7 and 8.

Note. A ♀ from Karori WN also mentioned by Meyrick (1914a, p. 111) has colour pattern elements on the forewing that Meyrick would have mentioned had he seen it. I conclude that his description is based on the single specimen now in BMNH.

Also 1 undescribed species (NZAC).



**Oecophoridae: "Oecophorinae"**  
(157) *Hierodoris illita* (Felder & Rogenhofer)

- ***Hierodoris*** Meyrick, 1912d, p. 41. Type species *Hierodoris iophanes* Meyrick, by original monotypy.

***Helostibes*** in the sense of Meyrick (1888e, p. 82), not of Zeller (1874).

Note. Examination of *H. mathewi* Zeller from Chile shows that true *Helostibes* shares no genital characters with the New Zealand species, and ocelli are absent.

***Taoscelis*** Meyrick, 1938, p. 428. Type species *Taoscelis crocostoma* Meyrick, by original monotypy. **New synonymy.**

***atychioides*** Butler, 1877, pp. 405–406 (*Tachyptilia*). **New combination**

Type locality not stated, specimens from “Colls Dr Hector and JD Enys, Esq.”, i.e., Dunedin DN or Christchurch MC; HT ♀ designated by Butler, BMNH.

Hudson 1928, pp. 306–307, pl. xxxiii fig. 23, after Meyrick 1888e, p. 83; as *Helostibes atychioides*, in Glyphipterigidae.

***gregalis*** Philpott, 1928a, p. 369 (*Helostibes*).

Synonymised by Hudson (1939, p. 456). Russell ND, Mr Florance; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 456, as synonym.

***barbarica*** Philpott, 1930b, p. 12 (*Helostibes*).

Synonymised by Hudson (1939, p. 456). Whangarei ND, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 456, as synonym.

Note. The HT agrees externally with *H. atychioides* in the long antennal ciliations and the uninflated valvae.

***bilineata*** Salmon, 1948, p. 310 (*Helostibes*). **New combination**

Great Island, Three Kings Islands, E.G. Turbott; HT ♂ designated by Salmon, AMNZ.

***callispora*** Meyrick, 1912d, p. 41 (*Helostibes*). **New combination**

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 306, pl. xxxiii fig. 24, in Glyphipterigidae.

***chlorobela*** Meyrick, 1921, p. 334 (*Helostibes*). **New combination**

Mount Arthur NN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 307, pl. xl ix fig. 27 and 28; 1950, p. 111, pl. iv fig. 1; in Glyphipterigidae.

***electrica*** Meyrick, 1889b, p. 157 (*Helostibes*). **New combination**

[Mount Peel NN], E. Meyrick; HT ♀ unique, BMNH. Hudson 1928, p. 306, pl. xxxiii fig. 26, in Glyphipterigidae. Note. Meyrick's Diary of Captures and his labels on

*Borkhausenia siderota* agree in showing that *B. siderota* (as “*Hel. siderota*”) was taken from *Aciphylla* flowers on Mount Arthur NN on 16 January 1886. On 17 January 1886, on Mount Peel, Meyrick records “*Hel. electrica*, 1”, and this I interpret as the HT. In my experience *H. electrica* rarely strays from its host, *Cassinia* species (Asteraceae).

***eremita*** Philpott, 1930c, p. 438 (*Hierodoris*)

Ball Glacier MK, A. Philpott; HT ♀ designated by Philpott, CMNZ.

Hudson 1939, p. 456, pl. lx fig. 10, in Glyphipterigidae.

***frigida*** Philpott, 1923, p. 153 (*Hierodoris*)

Dun Mountain NN/MB, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, pp. 305–306, pl. xl ix fig. 16, in Glyphipterigidae.

***crocostoma*** Meyrick, 1939, p. 428 (*Taoscelis*). **New synonymy.**

Freehold Range MK, S. Lindsay; HT ♂ unique, designated by Meyrick, CMNZ.

Hudson 1939, p. 451, pl. lvii fig. 29, in Oecophoridae.

***illita*** Felder & Rogenhofer, 1875, pl. cxl fig. 32 (*Atychia*). **New combination**

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.

Hudson 1928, p. 307, pl. xxxiii fig. 25, as *Helostibes illita*, in Glyphipterigidae.

***iophanes*** Meyrick, 1912d, p. 42 (*Hierodoris*)


Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 305, pl. xxxiii fig. 3, in Glyphipterigidae.

Also 6 undescribed species (NZAC).

—◎—

**Group B: Oecophorinae without ocelli**



**Oecophoridae: Oecophorinae**  
(158) *Izatha peroneonella* (Walker)

● **Atomotricha** Meyrick, 1883a, p. 423 (key); 1883e, pp. 324–325. Type species *A. ommatias* Meyrick, by original monotypy.

**Brachysara** Meyrick, 1883a, p. 424 (key); 1883e, p. 325. Type species *Oecophora sordida* Butler, by monotypy. Synonymised by Meyrick (1914a, p. 109).

**chloronota** Meyrick, 1914a, p. 110 (*Atomotricha*)  
Invercargill SL, A. Philpott; LT ♂ here designated, labelled  
“Invercargill New Zealand P. /00”, “Atomotricha  
chloronota” Meyr. 1/3 E. Meyrick det. in Meyrick  
Coll.”, BMNH.  
Hudson 1928, p. 288, not figured.

**exsomnis** Meyrick, 1913a, pp. 26–27 (*Atomotricha*)  
Ohakune TO/RI, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 288, pl. xxv fig. 41.

**isogama** Meyrick, 1909a, pp. 13–14 (*Atomotricha*)  
Wellington WN, G.V. Hudson; LT ♂ here designated,  
labelled “Wellington New Zealand. GVH /93”, “Ato-  
motricha isogama” Meyr. 9/13 E. Meyrick det. in  
Meyrick Coll.”, BMNH.  
Hudson 1928, p. 288, pl. xxv fig. 40.

**lewi** Philpott, 1927d, p. 87, fig. 5 (*Atomotricha*)  
?Central Otago CO, J.H. Lewis; HT ♂ designated by  
Philpott, NZAC.  
Hudson 1939, p. 451, not figured.

**oeconomia** Meyrick, 1914a, p. 110 (*Atomotricha*)  
Karori, Wellington WN, G.V. Hudson; LT ♂ here des-  
ignated, labelled “Wellington New Zealand GVH  
8.12”, “Atomotricha oeconomia” Meyr. 1/3 E. Meyrick  
det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 288, pl. xxxi fig. 16, 22, and 29.

**ommatias** Meyrick, §1883c, p. 522; 1883e, p. 325,  
abbreviated description; 1884a, p. 10, full  
description (*Atomotricha*)

Christchurch MC, E. Meyrick; LT ♂ here designated,  
labelled “Christchurch New Zealand 24/9/82”, “Ato-  
motricha ommatias” Meyr. 1/6 E. Meyrick det. in  
Meyrick Coll.”, BMNH.

Hudson 1928, pp. 287–288, not figured.

**prospiciens** Meyrick, 1924b, p. 662 (*Atomotricha*)  
[Pompolona Huts, Milford Track FD], C.E. Clarke; HT  
♂ unique, BMNH.

Hudson 1928, p. 289, pl. li fig. 1 and 2.

Note. Hudson states “taken by Mr C.E. Clarke at Pom-  
polona, Te Anau–Milford Track, and at Orepuki,  
Southland”. Meyrick’s label reads “Dunedin New  
Zealand E.C.C. 12.19”, and his description states  
“Dunedin”.

**sordida** Butler, 1877, p. 405 (*Oecophora*)

Dunedin DN or Christchurch or Castle Hill MC, J. Hector  
or J.D. Enys; HT ♂ designated by Butler, BMNH.  
Hudson 1928, p. 288, pl. xxxi fig. 15.

**versuta** Meyrick, 1914a, p. 109 (*Atomotricha*)

Karori, Wellington WN, G.V. Hudson; LT ♂ here des-  
ignated, labelled “Wellington New Zealand GVH  
8.12”, “Atomotricha versuta” Meyr. 1/17 E. Meyrick  
det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 287, pl. xxxi fig. 7, 14, and 18–21.

Note. Philpott (1927f, p. 109) could not separate *A. ver-  
suta*, *A. chloronota*, and *A. sordida* on genital  
characters.

Also 1 undescribed species (NZAC).

● **Barea** Walker, 1864a, p. 819. Type species *Barea consignatella* Walker, by original monotypy.

**Phloeopola** Meyrick, 1883a, p. 423 (key); 1883e,  
pp. 347–348. Type species *Oecophora confusella*  
Walker, by subsequent designation (Meyrick 1915b,  
p. 218). Synonymised by Meyrick (1915b, p. 218).

**confusella** Walker, 1864a, pp. 682–683 (*Oecophora*)  
Sydney N.S.W., Lambert; LT ♂ selected and labelled by  
I.F.B. Common, BMNH.

New Zealand: adventive; so far restricted to AK.  
Not mentioned by Hudson.

Note. The HT has yellowish hindwings (see below), as  
have specimens in BMNH from “NSWales” (Mathew)  
and Parramatta N.S.W., and in ANIC from Bunya  
Mountains Qld to Canberra A.C.T., although the apices  
may be darkened. Specimens in BMNH from Gis-  
borne Vict. and Hobart Tasm. and in ANIC from  
Eumungerie N.S.W. have grey hindwings.

**confusella** in the sense of Philpott (1927f, pp. 108  
and 111, fig. 37) (*Barea*)

New Zealand: adventive (Meyrick 1911b, p. 69); through-  
out, including Chatham Islands.

Hudson 1928, p. 289, pl. xxxiii fig. 11.

Note. Hudson (1928) noted that New Zealand specimens  
(first recorded in 1908) were of the “Victorian form

with grey hindwings". Philpott's drawing of genitalia matches those of 'grey' specimens in both countries but not those of HT *confusella* nor of New Zealand 'yellow' specimens, which have the valval costa produced into a long, acute process with or without 1 or 2 obscure 'teeth' on the ventral margin. As New Zealand has received both morphs, it will be interesting to see whether they interact.

***consignatella*** Walker, 1864a, pp. 819–820 (*Barea*)  
Moreton Bay Qld, Diggles; HT ♂ unique, BMNH.  
New Zealand: adventive; recorded ND–AK from dead wood in urban areas.  
Not mentioned by Hudson.  
Note. New Zealand specimens agree with the HT, possessing the characteristic rose-flushed scales.

***exarcha*** Meyrick, 1883e, pp. 357–358 (*Phloeopola*)  
Mount Gambier S. Austr., E. Meyrick; HT ♂ unique, BMNH.  
New Zealand: adventive; throughout.  
Hudson 1928, pp. 289–290, pl. li fig. 20.

***planetella*** Hudson, 1923d, p. 218 *Izatha*. Synonymised by Philpott (1927d, p. 88) under *Barea exartha*.  
?Ohakune RI/TO, S. Lindsay; HT ♀ unique, CMNZ.  
Hudson 1928, p. 289, as synonym.

Also 1 undetermined species from HB-WN (NZAC).

#### ● ***Borkhausenia*** of authors

***morella*** Hudson, 1939, pp. 444–445 (*Borkhausenia*) new combination  
Onehunga AK, A.J. Hipwell; HT not located in NMNZ.  
Hudson 1939, pp. 444–445, pl. lxi fig. 31, as *Borkhausenia morella*.

Note. The large compound eyes and lack of gnathos modifications exclude this species from *Tingena*.

● ***Chersadaula*** Meyrick, 1923, p. 165. Type species  
*Chersadaula ochrogastra* Meyrick, by original monotypy.

***ochrogastra*** Meyrick, 1923, p. 165 (*Chersadaula*)  
Breaker Bay, Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH bred 11.20", "Chersadaula ochrogastra Meyr. 2/5 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 272, pl. xlvi fig. 4 and 12.  
Note. The ♂ genitalia figured by Philpott (1927f, p. 110, fig. 51) differ from those of a specimen from Stephens Island, Cook Strait (BM genitalia slide no. 3100).

Also 1 undescribed species (BMNH, NMNZ).

● ***Compsistis*** Meyrick, 1888e, p. 89. Type species  
*Gelechia bifaciella* Walker, by original monotypy.

***bifaciella*** Walker, 1864, pp. 657–658 (*Gelechia*)  
[Auckland AK], D. Bolton; LT ♀ selected by H. Durrant, labelled "Gelechia bifaciella Wkr Cat. Lep. Het. BM 29, p. 657 (1864) Type ♀", "N. Zeal 54.4" (circular label), abdomen missing, BMNH.  
Hudson 1928, p. 273, pl. xxx fig. 14.

● ***Corocosma*** Meyrick, 1927a, p. 699. Type species  
*C. memorabilis* Meyrick, by original monotypy.

***memorabilis*** Meyrick, 1927a, p. 700 (*Corocosma*)  
Shedwood Forest, Tapawera NN, Stella Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 287, pl. lii fig. 12.

● ***Endrosis*** Hübner, 1825, p. 401. Type species *Tinea betulinella* Hübner, 1818–19 (= *Tinea sarcitrella* Linnaeus), by subsequent designation (Busck 1908, p. 203).

***sarcitrella*** Linnaeus, 1758, p. 536 (*Phalaena Tinea*)  
Europe. New Zealand: adventive; human habitation and birds' nests, throughout. On The Snares and subantarctic islands only in dwellings.

Hudson 1928, p. 260, pl. xxviii fig. 12, as *Endrosis lacteela*.

***subditella*** Walker, 1864a, p. 657 (*Gelechia*). Synonymised by Meyrick (1889b, p. 160).  
"New Zealand", J. Clarke Ross; HT ♂ unique, BMNH [not seen].

● ***Eulechria*** Meyrick, 1883a, pp. 424 (key) and 508.  
Type species *Eulechria exanimis* Meyrick, 1883a, pp. 511 (key) and 519, by subsequent designation (Meyrick 1915b, p. 218).

***zophoessa*** Meyrick, 1883a, pp. 510 (key) and 515–516 (*Eulechria*)  
[Botanic Gardens], Wellington WN, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 290, pl. xxxii fig. 27.

● ***Euchersadaula*** Philpott, 1926c, p. 414. Type species *Trachypepla lathriopa* Meyrick, by original designation.

***lathriopa*** Meyrick, 1905, p. 237 (*Trachypepla*)  
Nelson NN, E. Meyrick; LT ♂ here designated, labelled "Nelson New Zealand 12/1/86", "Euchersadaula lathriopa Meyr. 9/10 E. Meyrick det. in Meyrick Coll.", abdomen in gelatin capsule, BMNH.  
Hudson 1928, p. 273, pl. xxxi fig. 13.

***tristis*** Philpott, 1926a, p. 393, fig. 9 and 10 (*Euchersadaula*)  
Nelson NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 273, not figured, and p. 286, pl. xxxi fig. 12, as *Trachypepla anastrella* (Hudson 1939, p. 446).

➤ Oecophoridae, *Euchersadaula tristis*

Note. Philpott's papers in *Transactions of the New Zealand Institute* 56 were printed in reverse order, so that the description of *Euchersadaula* is preceded by the description of *Euchersadaula tristis*. As the author's intention is clear, no corrective action is needed.

● ***Euthictis*** Meyrick, 1914b, p. 246. Type species *Compsotropha xanthodelta* Meyrick, by original designation; Australia.

***chloratma*** Meyrick, 1916b, p. 416 (*Trachypepla*)  
Table Hill SI, A. Philpott; HT ♂ unique, abdomen distorted, BMNH.

Hudson 1928, p. 291, pl. xlvi fig. 3.

Note. On genital and head characters *E. chloratma* may be better placed in genus *Tingena* (p. 99); its current placing is dictated by forewing vein *R5* ending on the termen, not the costa.

● ***Gymnobathra*** Meyrick, 1883a, p. 423 (key); 1884a, pp. 27–28. Type species *Gelechia flavidella* Walker, by subsequent designation (Meyrick 1915, p. 214).

***Leptosaces*** Meyrick, 1888e, p. 77. Types species *Leptosaces callixyla* Meyrick, by original monotypy. **New synonymy.**

Note. The ♂ genitalia of *L. callixyla* resemble those of *G. caliginosa* and *G. cenchrias* as figured by Philpott (1927b, p. 720, fig. 9 and 12). In habitus and colour pattern *L. callixyla* resembles *G. caliginosa*. *Leptosaces* is here removed from synonymy with *Barea* and synonymised with *Gymnobathra* (see Philpott 1927b, p. 720, fig. 9 and 12; 1927f, p. 108, fig. 37).

***ambigua*** Philpott, 1926a, p. 396, fig. 21 and 22 (*Barea*) **new combination**

Horseshoe Lake, Christchurch MC, W. Heighway; HT ♂ designated by Philpott, CMNZ.  
Hudson 1928, p. 289, pl. lii fig. 15, as *Barea ambigua*.

***bryaula*** Meyrick, 1905, pp. 238–239 (*Gymnobathra*)  
Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 276, pl. xxx fig. 22 and 23.

***caliginosa*** Philpott, 1927a, p. 707 (*Gymnobathra*)  
Cooper's Knob, Port Hills MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.  
Hudson 1939, p. 447, pl. lix fig. 28.

***calliploca*** Meyrick, 1883c, p. 523; 1884a, pp. 28 (key) and 30–31 (*Gymnobathra*)

Dunedin DN, A. Purdie; HT ♂ unique, worn, BMNH.  
Hudson 1928, p. 276, pl. xxx fig. 9.  
Note. The HT valva and uncus differ from those figured by Philpott (1927b, p. 719, fig. 8).

***callixyla*** Meyrick, 1888e, p. 78 (*Leptosaces*) **new combination**

Nelson NN, E. Meyrick; LT ♂ here designated, labelled “Nelson New Zealand 12/1/86”, “Cryptolechia callixyla Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 293, not figured, as *Cryptolechia callixyla*. Note. The PLT ♀ is perhaps not conspecific.

***cenchrías*** Meyrick, 1909a, p. 13 (*Borkhausenia*)

Invercargill SL, A. Philpott; HT ♂ unique, BMNH.

Hudson 1928, pp. 274–275, pl. xxix fig. 30, as *Gymnobathra cenchrías*.

Note. Philpott (1926b, p. 401) removed *cenchrías* to *Gymnobathra*.

***dinocosma*** Meyrick, §1883c, p. 522 (as *dinacosma*); 1883e, pp. 348 (key) and 349, abbreviated description; 1884a, p. 12, full description (*Phloeopola*) **new combination**

“Wellington” WN, E. Meyrick; [HT ♂ destroyed; letter from Meyrick to Hudson, 12 April 1889]. Candidate NT material in BMNH.

Hudson 1928, p. 289, pl. xxxii fig. 10, as *Barea dinocosma*.

Note. Meyrick's HT was recorded in his Diary of Captures for 1 January 1880 as “Oe. dinophorella, 1”. The specimen Meyrick was “glad to get” from Hudson (Meyrick's letter to Hudson, 12 April 1889) is also from Wellington, and is labelled “Barea dinocosma Meyr. 3/9 E. Meyrick det. in Meyrick Coll.” (BMNH). The tegumen and most of the valvae are missing. Should a NT ever be needed, specimens 3/9 and 9/9 (the latter also from Wellington, and bearing Meyrick's label “dinocosma Meyr.”) are candidates.

***thetodes*** Meyrick, 1901, p. 574 (*Gymnobathra*). **New synonymy.**

Akaroa, Banks Peninsula MC, R.W. Fereday; LT ♂ here designated, labelled “Akaroa New Zealand RWF 16/1/72”, “Gymnobathra thetodes Meyr. 1/1 E. Meyrick det. in Meyrick Coll.”, valval tips broken, BMNH.

Hudson 1928, p. 276, pl. xxx fig. 24, as species. Note. Genitalia and colour pattern agree with Wellington *dinocosma*, particularly with *dinocosma* specimen 9/9, which bears a note in Meyrick's handwriting “dinocosma Meyr.” and a label “Wellington New Zealand GVH /96”.

***flavidella*** Walker, 1864a, p. 655 (*Gelechia*)

[Auckland AK], D. Bolton; HT ♂ unique, BMNH.

Hudson 1928, p. 275, pl. xxx fig. 18 and 19.

***utuella*** Felder & Rogenhofer, 1875, p. cxl fig. 46 (*Oecophora*). **Synonymised by Meyrick (1884a, p. 31).**

[Nelson NN, T.R. Oxley]; HT ♀ unique, abdomen and hindwings glued on, BMNH.

Hudson 1928, p. 275, as synonym.

***hamatella*** Walker, 1864a, p. 700 (*Oecophora*)

[Nelson NN], T.R. Oxley; LT ♀ here designated, labelled “Oecophora hamatella Wkr Cat. Lep. Het. BM 29 p.700 (1864) Type ♂”, “Auckland N. Zeal. 60-73” (circular label), abdomen missing, BMNH.

Hudson 1928, p. 274, pl. xxx fig. 16 and 17.

**omichleuta** Meyrick, 1929, p. 489 (*Gymnobathra*). New synonymy.

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 3.29", "Gymnobathra omichleuta Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", BMNH.

Note. Except for the overall grey-brown shading, the LT ♂ and the PLT ♂ are indistinguishable from *hamatella* ♂♂ on colour pattern and external genital features, and the two taxa are here synonymised (cf. distinctions between *G. hyetodes* and *G. hamatella* - Philpott 1927b, p. 718, fig. 2 and 4).

**hyetodes** Meyrick, §1883c, pp. 523–524; 1884a, pp. 28 (key) and 32–33 (*Gymnobathra*)

Wellington WN, R.W. Fereday; LT ♂ here designated, labelled "Wellington New Zealand R.W.F. 82", "Gymnobathra hyetodes Meyr. 2/4 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 274, pl. xxx fig. 11 and 12.

**inaequata** Philpott, 1928a, p. 367 (*Gymnobathra*)

Dun Mountain NN/MB, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 447, pl. lix fig. 16.

**jubata** Philpott, 1918, p. 131 (*Dolichernis*)

Tisbury, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 325, pl. xl fig. 5, as *Dolichernis jubata*, in Plutellidae.

Note. Venation, the scaled haustellum, head characters, and external genital characters agree with *Gymnobathra* and Oecophoridae. The colour pattern is reminiscent of the BMNH ♀ specimen of *G. callixyla*.

**levigata** Philpott, 1928a, pp. 366–367 (*Gymnobathra*)

Dun Mountain NN/MB, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 447, pl. lix fig. 17.

**omphalota** Meyrick, 1888e, p. 81 (*Gymnobathra*)

Lake Wakatipu OL, E. Meyrick; LT ♀ here designated, labelled "L. Wakatipu New Zealand 16.12.82", "Gymnobathra omphalota Meyr. 4/9 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 277, pl. xxx fig. 25 and 26.

**origenes** Meyrick, 1936, p. 282 (*Gymnobathra*)

Mount St Arnaud BR/MB, S. Lindsay; HT ♀ unique, left forewing and abdomen missing, CMNZ.

Hudson 1939, p. 448, pl. lxii fig. 17.

**parca** Butler, 1877, p. 405 (*Oecophora*)

Canterbury MC or Dunedin DN, J.D. Enys or J. Hector; HT ♂ designated by Butler, BMNH.

Hudson 1928, p. 276, pl. xxx fig. 1 and 2, as *Gymnobathra parca* (see Meyrick §1883c, p. 523; 1884a, p. 29).

**limbata** Butler, 1880, p. 560 (*Oecophora*). Synonymised by Meyrick (1915b, p. 215).

Blenheim MB, W. Skellon; HT ♂ designated by Butler, BMNH.

Hudson 1928, p. 276, as synonym of *Gymnobathra parca*.

**philadelphia** Meyrick, 1883c, p. 524; 1884a, pp. 28 (key) and 33 (*Gymnobathra*)

[bush gully north of Conical Hill], Mount Hutt MC, R.W. Fereday; LT ♀ here designated, labelled "Mt Hutt New Zealand RWF. 82", "Gymnobathra philadelphia Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 274, not figured.

**habropis** Meyrick, 1888c, pp. 80–81 (*Gymnobathra*). Synonymised by Philpott (1927b, pp. 716–717).

Nelson NN, E. Meyrick; LT ♂ here designated, labelled "Nelson New Zealand 22/1/86", "Gymnobathra habropis Meyr. 3/5 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 274, pl. xxx fig. 10, as species.

**primaria** Philpott, 1928a, p. 366 (*Gymnobathra*)

Mount Arthur NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 447, pl. lix fig. 18.

**rufopunctella** Hudson, 1950, p. 107, pl. v fig. 7 (*Gymnobathra*)

Day's Bay WN, G.V. Hudson; LT ♂ here designated, labelled "1161a" ["Days Bay Nov. 22. 1929"], NMNZ.

**sarcoxantha** Meyrick, §1883c, p. 523; 1884a, pp. 28 (key) and 29 (*Gymnobathra*)

[reserved bush and forest], Dunedin DN, E. Meyrick; LT ♂ here designated, labelled "Dunedin New Zealand 6.1.80", "Gymnobathra sarcoxantha 1/6 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 275, not figured as such, but pl. xxx fig. 15 refers.

**coarctatella** not of Walker, but in the sense of Meyrick (§1883c, p. 523; 1884a, p. 28 (*Gymnobathra*))

Hudson 1928, p. 275, pl. xxx fig. 15.

Note. *Cryptolechia coarctatella* Walker is a senior synonym of *Phaeosaces liochroa* Meyrick (q.v.).

**squamata** Philpott, 1915, p. 200 (*Gymnobathra*)

Mount Cleughearn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 277, pl. xxx fig. 13.

**nigra** Philpott, 1930b, pp. 9–10 (*Gymnobathra*). Synonymised by Hudson (1939, p. 447).

Kepler Mountains FD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 447, as synonym.

**tholodella** Meyrick, §1883c, p. 523 (abstract); 1884a, pp. 28 (key) and 30 (*Gymnobathra*)

» Oecophoridae, *Gymnobathra tholodella*

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 18/2/82", "Gymnobathra tholodella Meyr. 8/12 E. Meyrick det. in Meyrick Coll.", genitalia with left valva missing, BMNH.

Hudson 1928, pp. 276–277, pl. xxx fig. 3.

• **Hofmannophila** Spuler, 1910, p. 340. Type species *Oecophora pseudospretella* Stainton, by monotypy.

**pseudospretella** Stainton, 1849, p. 14 (*Oecophora*) Europe.  
New Zealand: adventive; human habitation and birds' nests, throughout.

Hudson 1928, p. 271, pl. xxx fig. 7, as *Borkhausenia pseudospretella*.

• **Izatha** Walker, 1864a, p. 786. Type species *Izatha attactella* Walker, by original monotypy.

**Aochleta** Meyrick, 1883a, p. 425 (key); 1884a, p. 21 (description and included species). Type species *A. psychra* Meyrick, by original monotypy. New synonymy.

**Semicosma** Meyrick, 1883a, p. 424 (key); 1884a, p. 21 (description and included species). Type species *Gelechia peroneonella* Walker, by subsequent designation (Meyrick 1915b, p. 215). Synonymised by Meyrick (1905, p. 238).

**Zirosaris** Meyrick, 1910a, p. 66. Type species *Zirosaris amorbas* Meyrick, by original monotypy. Synonymised by implication (Meyrick, 1920a, p. 31).

**amorbas** Meyrick, 1910a, p. 66 (*Zirosaris*)  
Broken River MC, J.H. Lewis; HT ♂ not in BMNH.

**amorbas** Meyrick, 1911b, p. 66 (*Trachypepla*).  
Synonymised by Hudson (1928, p. 282).  
Invercargill SL, A. Philpott; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 282, pl. xxxii fig. 7, as *Izatha amorbas*. Note. Meyrick (1919, p. 353) transferred *amorbas* from *Trachypepla* to *Izatha* on the basis of more Invercargill specimens received from Philpott. NZAC also has Invercargill specimens.

**apodoxa** Meyrick, 1888e, pp. 79–80 (*Semicosma*)  
Wellington WN, A. Purdie; LT ♂ here designated, labelled "Wellington New Zealand /84", "Izatha apodoxa Meyr. 3/4 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, pp. 279–280, pl. xxxii fig. 2.

**attactella** Walker, 1864a, p. 787 (*Izatha*)  
[Auckland AK], D. Bolton; LT ♂ (?selected by H. Durrant from the 15 STs) labelled "Izatha attactella Wkr Cat. Lep. BM 29 p.787 (1864) Type ♂", "New Zeal. 54.4", "Holotype", BMNH.

Hudson 1928, p. 280, pl. xxv fig. 30 and 31. Philpott 1927f, p. 104, fig. 10.

**platyptera** Meyrick, 1888e, p. 80 (*Semicosma*).  
Synonymised by Meyrick (1915b, p. 215).

Wellington WN, G.V. Hudson; LT ♀ selected by P.A. Brown and here designated, labelled "Wellington New Zealand GVH /85", "Izatha attactella Walk. 7/8 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 280, as synonym.

**austera** Meyrick, §1883c, p. 523 (summary); 1884a, pp. 25–26 (*Semicosma*)

[Botanic Gardens and forest], Wellington WN, E. Meyrick; LT ♂ here designated, labelled "Wellington New Zealand 1/1/80", "Izatha austera Meyr. 1/9 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 282, pl. xxxii fig. 6.

**balanophora** Meyrick, 1897b, p. 389 (*Semicosma*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1939, p. 448, pl. lix fig. 8.

**milligani** Philpott, 1927d, p. 87 (*Izatha*).  
Synonymised by Hudson (1939, p. 448).

Leigh AK, D.D. Milligan; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 448, as synonym.

**caustopa** Meyrick, 1892, p. 219 (*Semicosma*)

Wellington WN, G.V. Hudson; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 280, pl. xxv fig. 25.

**churtoni** Dugdale, new species for *picarella* var. β  
Walker (*Izatha*)

**picarella** var. β Walker, 1864, p. 699 (*Oecophora*)

[?Auckland AK], J.F. Churton; HT ♀ unique, labelled "var. β", "51.136 N. Zeal." (circular label), BMNH.  
Hudson 1928, p. 279, pl. xxv fig. 39, as *Izatha picarella*. Note. Walker stated that var. β is characterised by its uniformly grey hindwings. It also has the forewing termen black and white (all white in *I. picarella*), and ♂ genitalia markedly different from those of *I. picarella* as figured by Philpott (1927f, p. 106, fig. 15).

**convulsella** Walker, 1864a, p. 656 (*Gelechia*)

[Nelson NN], T.R. Oxley; LT ♀ selected by H. Durrant as "Type", labelled "379 Gelechia convulsella" [printed], "Gelechia convulsella Wkr Cat. Lep. BM 29 p.656 (1864) Type ♀", "Auckland N. Zeal. 60-73" [circular], "Type", metathorax and abdomen glued, BMNH.

Hudson 1928, p. 282, pl. xxxi fig. 1.

**paraneura** Meyrick, 1892, p. 219 (*Semicosma*).  
Synonymised by Meyrick (1915, p. 216).

Wellington WN, G.V. Hudson; LT ♀ here designated, labelled "Wellington New Zealand GVH /90", "Izatha convulsella Meyr. 5/12 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 282, as synonym.

**copiosella** Walker, 1864b, p. 1028 (*Gelechia*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 280, pl. xxv fig. 27, but not fig. 26.  
Note. Hudson's fig. 26 on pl. xxv represents an unnamed species. See entry at end of *Izatha* species, below.

**epiphanes** Meyrick, §1883c, p. 523; 1884a, pp. 24–25 (*Semicosma*)

Wellington WN, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 281, pl. xxv fig. 24.

**florida** Philpott, 1927d, p. 86 (*Izatha*)

Mount Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 448, pl. lix fig. 10.

**griseata** Hudson, 1939, p. 448 (*Izatha*)

Mataitai, Auckland AK, C.E. Clarke; HT ♀ unique, not located in AMNZ.

Hudson 1939, p. 448, pl. lxii fig. 2.

**heroica** Philpott, 1926a, p. 396 (*Izatha*)

Flora River, Mount Arthur NN, A. Philpott; HT ♀ designated by Philpott, NZAC.

Hudson 1928, pp. 278–279, pl. xxxii fig. 32 and 33.

**toreuma** Clarke, 1926, p. 419 (*Izatha*). Synonymised by Hudson (1928, p. 278).

Arthur's Pass NC/WD, C.E. Clarke; HT ♀ unique, AMNZ.

Hudson 1928, p. 278, as synonym.

**hudsoni** Dugdale, new species for *huttoni* of authors, but not Butler (*Izatha*)

**huttoni** not of Butler (1879a) but in the sense of Meyrick (1916b, p. 416) and Hudson (1928, p. 278, pl. xxv fig. 43)

Karori, Wellington WN, G.V. Hudson; HT ♂ designated by Dugdale, genitalia on Philpott genitalia slide, NZAC. Hudson 1928, p. 278, pl. xxv fig. 43. Philpott 1927f, pp. 105 and 106, fig. 13.

**manubriata** Meyrick, 1923, pp. 165–166 (*Izatha*)

Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 279, pl. xxx fig. 8.

**mesoschista** Meyrick, 1931a, p. 96 (*Izatha*)

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH.11", "Izatha mesoschista Meyr. 4/7 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 279, pl. xxxii fig. 3, as ♂ *Izatha balanophora*; 1939, p. 448.

**metadelta** Meyrick, 1905, p. 238 (*Izatha* (*Semicosma*))

Wellington WN, G.V. Hudson; LT ♀ here designated, labelled "Wellington New Zealand GVH /03", "Iza-

tha metadelta Meyr. 6/8 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 281, pl. xxx fig. 20 and 21.

**percnitis** Meyrick, 1909a, p. 14 (*Izatha*). Synonymised by Meyrick (1912c, p. 121).

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH /00", "Izatha metadelta Meyr. 2/8 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 281, as synonym.

**mira** Philpott, 1913, p. 78 (*Izatha*)

The Hump FD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 281, pl. xxxi fig. 6.

**oleariae** Dugdale, 1971b, pp. 133–134 (*Izatha*)

Station Point, The Snares islands, P.M. Johns; HT ♂ designated by Dugdale, NZAC.

**peroneanella** Walker, 1864a, p. 658 (*Gelechia*)

[Auckland AK], D. Bolton; LT ♂ selected by H. Durrant and here designated, labelled "Gelechia peroneanella Wlk Cat. Lep. BM 29 p.658 (1864) Type ♂", "New Zeal. 54.4" (circular label), "Type HT" (red-margined circular label), BMNH.

Hudson 1928, p. 278, but pl. xxv fig. 54 portrays a colour pattern different from that of the type population.

Note. Durrant's PLT *peroneanella* (from Nelson NN!) has a different forewing pattern.

**padapertella** Walker, 1864a, p. 653 (*Gelechia*). Synonymised by Meyrick (1915, p. 215).

[Auckland AK], D. Bolton; HT (?♂) unique, lost [Durrant, Stringer]; specimen in its place is *Endrosis sarcitrella* [Durrant, Stringer]; BMNH.

Note. Walker's description agrees with a faded, weakly marked *peroneanella*.

**lichenella** Walker, 1864a, p. 769 (*Cryptolechia*).

Synonymised by Meyrick (1884a, p. 22).

Auckland AK, A. Sinclair; HT ♂ unique, BMNH.

Note. HT *lichenella* has forewing markings as in LT *peroneanella* but brown, not black.

**huttoni** Butler, 1879a, p. 511 (*Oecophora*). Synonymised by Meyrick (1884a, p. 22).

[?Dunedin DN], "Otago", F.W. Hutton; HT ♀ unique, BMNH.

Hudson 1928, pl. xxv fig. 54 (illustrates colour pattern).

**mystis** Meyrick, 1888e, p. 79 (*Semicosma*). Synonymised by Meyrick (1915, p. 215).

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♀ here designated, labelled "Christchurch New Zealand 23/12/82", "Izatha peroneanella Walk., 9/13 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pl. xxv fig. 54 (illustrates colour pattern).

Note. Specimens from south of Auckland have the forewing discal black marks either separate (typical *peroneanella*) or joined to form an oblique M (typical *huttoni* and *mystis*). South Island specimens (including Durrant's PLT *peroneanella* collected by Oxley) are all of the *huttoni* / *mystis* pattern as illustrated by Hudson.

➤ Oecophoridae, *Izatha*

**phaeoptila** Meyrick, 1905, pp. 236–237  
 (*Trachypepla*)  
 Mangatarere River WN, G.V. Hudson; HT ♂ unique,  
 BMNH.  
 Hudson 1928, p. 282, pl. xxxi fig. 27.

**picarella** Walker, 1864a, p. 699 (*Oecophora*); Meyr-  
 ick, 1884a, p. 23 (*Semiocosma*)  
 [Nelson NN], T.R. Oxley; HT ♀ unique, abdomen miss-  
 ing, BMNH.

Hudson 1928, p. 279, pl. xxv fig. 29, as *Izatha acmonias*.

**teras** Felder & Rogenhofer, 1875, pl. cxl fig. 28  
 (*Psecadia*). Synonymised by Meyrick (1884a, p. 23).  
 [Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen of ♀  
 geometrid glued on, BMNH.

Hudson 1928, p. 279, as synonym.

**acmonias** Philpott, 1921, pp. 340–341 (*Izatha*).  
 New synonymy.

[Tisbury], Invercargill SL, A. Philpott; HT ♂ designated  
 by Philpott, NZAC.

Hudson 1928, p. 279, pl. xxv fig. 29, as species.

Note. All three names refer to the South Island black and  
 white *Izatha* with the termen fringe white. Walker (1864a,  
 p. 699) distinguished on hindwing pattern “var β”, col-  
 lected by Churton (*I. churtoni*, above) which has ♂ gen-  
 italia markedly different from those of *picarella* as figured  
 by Philpott (1927f, p. 106, fig. 15), and the forewing ter-  
 men fringe partly black.

**plumbosa** Philpott, 1927d, p. 86 (*Izatha*)  
 Otira Basin, Arthur's Pass WD, A. Philpott; HT ♀ des-  
 ignated by Philpott, NZAC.  
 Hudson 1939, pp. 448–449, pl. lix fig. 11.

**prasophyta** Meyrick, §1883c, p. 523; 1884a, p. 25  
 (*Semiocosma*)  
 [Botanic Gardens and forest], Wellington WN, E. Meyr-  
 ick; LT ♂ here designated, labelled “Wellington New  
 Zealand 1/1/80”, “*Izatha prasophyta* Meyr. 1/5 E.  
 Meyrick det. in Meyrick Coll.”, BMNH.  
 Hudson 1928, p. 281, pl. xxxii fig. 4.

**psychra** Meyrick, §1883c, p. 523; 1884a, p. 21  
 (*Aochleta*) new combination  
 Porter's Pass MC, J.D. Enys; HT ♂ unique, very worn,  
 BMNH.

Hudson 1928, p. 277, not figured.

Note. The HT label gives Porter's Pass, and Meyrick's  
 description gives Castle Hill. Enys lived at Castle Hill  
 Station, but would travel over Porter's Pass to and  
 from Christchurch.

**rigescens** Meyrick, 1929, p. 490 (*Izatha*)  
 Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
 Hudson 1939, p. 448, pl. lix fig. 9.

Also 1 undescribed species – *copiosella* in the sense  
 of Hudson (1928, pl. xxv fig. 26); BMNH and

NMNZ, the latter with Hudson's *copiosella*.

● **Lathicrossa** Meyrick, §1883c, p. 523; 1884a, p. 26.  
 Type species *Lathicrossa leucocentra* Meyrick, by  
 original monotypy.

**leucocentra** Meyrick, §1883c, p. 523; 1884a, pp. 26–  
 27 (*Lathicrossa*)  
 Dunedin DN, A. Purdie; HT ♂ unique, BMNH.  
 Hudson 1928, p. 293, pl. xxxii fig. 5.

**prophética** Meyrick, 1927a, p. 701 (*Lathicrossa*)  
 Mount Arthur NN, S. Woodward; HT ♂ unique, BMNH.  
 Hudson 1928, p. 293, pl. lii fig. 8.

● **Leptocroca** in the sense of Philpott, 1926a, not of  
 Meyrick (1883a, p. 425 (key); 1886a, p. 775  
 (description). Type species *Leptocroca sanguinolenta*  
 Meyrick, 1886a, pp. 775–776, by original  
 monotypy; Australia).

**amenena** Meyrick, 1888e, pp. 78–79 (*Peltophora*)  
 new combination

Mount Arthur [Plateau] NN, E. Meyrick; LT ♂ here des-  
 ignated, labelled “Mt Arthur New Zealand 15/1/86”,  
 “*Philobota amenena* Meyr. 2/3 E. Meyrick det. in  
 Meyrick Coll.”, BMNH.

Hudson 1928, p. 291, pl. lii fig. 21.

Note. Meyrick (1915b, p. 219) placed *amenena* in *Philo-  
 bota* Meyrick, another Australian genus. On genital  
 characters it is more similar to the New Zealand spe-  
 cies currently in *Leptocroca*, and is here transferred.

**aquilonaris** Philpott, 1931, pp. 31–32 (*Leptocroca*)  
 Kauri Gully AK, C.E. Clarke; HT ♂ designated by Phil-  
 pott, AMNZ.  
 Hudson 1939, p. 445, not figured.

**asphaltis** Meyrick, 1911b, p. 65 (*Borkhausenia*)  
 [?Mount Ida CO, J.H. Lewis], “exact locality unknown”;  
 HT ♂ unique, unrecognised, possibly the ♂ labelled  
 “*Leptocroca variabilis* Philp. 3/4 E. Meyrick det. in  
 Meyrick Coll.”, “New Zealand AP .09”, head missing,  
 forewings glued to thorax, abdomen in gelatin capsule,  
 BMNH.

Hudson 1928, p. 272, pl. xxix fig. 17.

Note. The Philpott Collection (NZAC) has 1 ♂ (genitalia  
 missing) with no locality label. Philpott (1926b, p. 401)  
 removed *asphaltis* to *Leptocroca* (as *Guestia*).

**lenita** Philpott, 1931, p. 32 (*Leptocroca*)  
 Newton Flat, Buller Gorge NN/BR, G.V. Hudson; HT ♂  
 designated by Philpott, AMNZ.  
 Hudson 1939, pp. 445–446, pl. lix fig. 4.

**lindsayi** Philpott, 1930a, p. 249 (*Leptocroca*)  
 Yaldhurst MC, S. Lindsay; HT ♂ designated by Philpott,  
 abdomen missing, CMNZ.  
 Hudson 1939, p. 446, pl. lix fig. 6.

**porophora** Meyrick, 1929, p. 488 (*Borkhausenia*) new combination

Takaka NN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1939, p. 445, pl. lvii fig. 7, as *Borkhausenia porophora*.

Note. Because the HT abdomen is not modified as in *Tingena* species, and the ostiolar lamellae form a triangular spout as in New Zealand *Leptocroca* species, *porophora* is here transferred to *Leptocroca*.

**scholaea** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 35 (*Oecophora*); Philpott, 1926a, p. 393 (*Leptocroca*)

[reserved bush and forest]. Dunedin DN, E. Meyrick; LT ♂ here designated, labelled "Dunedin New Zealand 6/1/80", "Leptocroca scholaea Meyr. 5/24 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pp. 271–272, pl. xxix fig. 18.

**vacua** Philpott, 1926a, pp. 393–394, fig. 19 and 20 (*Leptocroca*). New synonymy.

Nelson NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 272, not figured.

Note. Philpott's figures (1926a, fig. 19 and 20) of PT ♂ *vacua* genitalia and those of HT ♂ do not differ from HT ♂ *scholaea*, but differ from other specimens under *scholaea* in BMNH and NZAC. Philpott's figures (1926a, fig. 13 and 14) of *scholaea* refer to one of at least 5 undescribed species, differing in genitalia and frons structure, and hitherto included in collections under *scholaea*.

**variabilis** Philpott, 1926a, p. 394, fig. 17 and 18 (*Leptocroca*)

Cobb Valley NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 272, not figured.

Note. There is little to distinguish the PT ♂ *variabilis* genitalia as figured by Philpott from those of the possible *asphaltis* HT ♂.

**vinaria** Meyrick, 1914a, p. 108 (*Trachypepla*) new combination

Otira River WD, G.V. Hudson; LT ♂ here designated, labelled "Otira R. New Zealand GVH 2.08", "Trachypepla vinaria Meyr. 6/9 E. Meyrick det. in Meyrick Coll.", abdomen missing, BMNH.

Hudson 1928, pp. 285–286, pl. xxxii fig. 26, as *Trachypepla vinaria*.

**obliqua** Philpott, 1930c, p. 437 (*Leptocroca*). Synonymised by Hudson (1939, p. 449).

Mount Cook MK, A. Philpott; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 449, as synonym of *vinaria*. Note. Because of its bifid uncus, recurved gnathos, and spinose abdomen, as well as its colour pattern, this species is referred to *Leptocroca*, where Philpott (through *obliqua*) put it.

**xyrias** Meyrick, 1931b, p. 368 (*Leptocroca*)

Blackmillar KA, S. Lindsay; HT ♂ unique, CMNZ.  
Hudson 1939, p. 446, pl. lix fig. 5.

Also 5 undescribed species (BMNH, NZAC).

• **Locheutis** Meyrick, 1883e, p. 341. Type species *Locheutis philochora* Meyrick, 1883e, p. 342, by subsequent designation (Meyrick 1922d, p. 53); Australia.

**fusca** Philpott, 1930b, p. 11 (*Locheutis*)

Whakapapa River, Tongariro National Park TO, A. Philpott; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 452, pl. lviii fig. 8.

Note. HT ♂ (with red label) and PT ♂ are double-mounted on the same polyporus strip (as at March 1983).

**pulla** Philpott, 1928g, p. 488 (*Locheutis*)

Mount Cedric BR, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 452, pl. lviii fig. 7.

**vagata** Meyrick, 1916b, p. 416 (*Locheutis*)

Tararua Range WN, 4000 ft, G.V. Hudson; LT ♂ here designated, labelled "Tararua Mtns New Zealand GVH 4000' 11.14", "Locheutis vagata Meyr. 2/2 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 290, pl. xxix fig. 7.

• **Mermerristis** Meyrick, 1915c, p. 298. Type species *M. spodiaea* Meyrick, by original monotypy.

**spodiaea** Meyrick, 1915c, p. 298 (*Mermerristis*)

George's Bay Tasm., E. Meyrick; HT ♂ unique, BM genitalia slide no. 1362 ♂, BMNH.

New Zealand: Hudson's records are from Wilton's Bush and Wainuiomata, WN.

Hudson 1939, p. 451, pl. lviii fig. 13.

Note. The *spodiaea* HT ♂ genitalia appear to be identical with those of *Trachypepla ocneropis* Meyrick (q.v.). The occurrence of *M. spodiaea* on either side of the Tasman Sea may be natural.

• **Oxythecta** Meyrick, 1883a, p. 422 (key); 1885c, p. 1059 (description). Type species *Gelechia acceptella* Walker, by subsequent designation (Meyrick 1915b, p. 218); Australia.

**austrina** Meyrick, 1914a, p. 107 (*Saropla*); 1915b, p. 218 (*Oxythecta*)

Ben Lomond OL, A. Philpott; LT ♂ here designated, labelled "Ben Lomond New Zealand AP 25.11.12", "Oxythecta australis Meyr. 2/2 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 291, pl. xxxii fig. 14.

• **Parocystola** Turner, 1896, p. 30. Type species *Parocystola leucospora* Turner, by original monotypy, Australia.

» Oecophoridae, *Parocystola*

**acroxantha** Meyrick, 1885c, pp. 1060 (key) and 1066  
(*Ocystola*)

Deioraine Tas., E. Meyrick; LT ♂ selected by I.F.B. Common, BMNH.  
New Zealand: adventive; first recorded 1886 (Hudson 1928, p. 290). On dead leaves; throughout.  
Hudson 1928, p. 290, pl. xxx fig. 27, as *Parocystola acroxantha*.

● **Phaeosaces** Meyrick, §1885i, p. 591; 1886b, pp. 171–172. Type species *Phaeosace apocrypta* Meyrick, by subsequent designation (Meyrick 1915b, p. 220).

Note. *Cryptolechia* Zeller is based on *C. straminella* Zeller from South Africa, the ♂ genitalia of which bear no resemblance to those of New Zealand species (nor Nearctic species; see figures in Hodges 1974). *Phaeosaces* is available, and is here revived, as a generic name for New Zealand species.

**apocrypta** Meyrick, §1885i, p. 591; 1886b, pp. 172–173 (*Phaeosaces*)

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 23/12/82”, “*Cryptolechia apocrypta* Meyr. 5/11 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 294, pl. xxv fig. 21.

**coarctatella** Walker, 1864a, p. 768 (*Cryptolechia*)  
new combination

[Nelson NN], T.R. Oxley; LT ♂ (as HT) here designated, labelled “*Cryptolechia coarctatella* Wkr Cat. Lep. Het. BM 29 p.768 (1864) Type ♂”, “60-73 Auckland N. Zeal.” (circular label), BMNH.

Hudson 1928, p. 294, pl. xxv fig. 23, as *Cryptolechia liochroa*.

**liochroa** Meyrick, 1891, p. 98 (*Phaeosaces*). New synonymy.

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “Wellington New Zealand GVH /90”, “*Cryptolechia liochroa* Meyr. 3/10 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 294, pl. xxv fig. 22 and 23. Philpott 1927, p. 110, fig. 49. Both as species.

**compsotypa** Meyrick, §1885i, p. 591; 1886b, p. 172  
(*Phaeosaces*)

[in dense forest], Hamilton WO, E. Meyrick; HT ♀ unique, BMNH.

Hudson 1928, pp. 294–295, pl. xxxii fig. 1.

**lindsayae** Philpott, 1928c, p. 182 (*Cryptolechia*) new combination

Blackmillar KA, Jean Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 453, pl. lviii fig. 16.

Also 2 undescribed species (NZAC).

● **Schiffermuelleria** Hübner, [1825], p. 421. Type species *Phalaena Tinea schaefferella* Linnaeus; Europe.

**orthophanes** Meyrick, 1905, p. 243 (*Compsistis*); 1915b, p. 211 (*Schiffermuelleria*)

Nelson NN, E. Meyrick; LT ♀ here designated, labelled “Nelson New Zealand 22.1.86”, “*Schiffermuelleria orthophanes* Meyr. 3/6 E. Meyrick det. in Meyrick Coll.”, very worn, BMNH.  
Hudson 1928, p. 260, pl. xxviii fig. 17.

● **Sphyrelata** Meyrick, 1883a, p. 423 (key); 1883e, p. 360 (description). Type species *Oecophora amoetella* Walker, by subsequent designation (Meyrick 1922d, p. 151); Brisbane, Qld.

**laetifica** Turner, 1917, p. 116, proposed as a new name for *indecorella* Meyrick, not Walker, 1864 (*Sphyrelata*)

**indecorella** not of Walker (1864a, p. 764) but in the sense of Meyrick (1883e, p. 362) and Turner (1917, p. 116).

New Zealand: adventive; earliest record from Henderson AK (K.A.J. Wise, 1954), subsequently recorded from AK and WI. All specimens seen have orange hindwings.

Note. Meyrick mentioned material from Sydney and Mittagong, N.S.W. Turner stated that *S. laetifica* is distinguished by its orange hindwings.

● **Thamnosara** Meyrick, §1883c, p. 523; 1884a, p. 27. Type species *Thamnosara chirista* Meyrick, by original monotypy.

**sublitella** Walker, 1864a, p. 654 (*Gelechia*)

[Auckland AK], A. Sinclair; HT ♀ unique, head missing, abdomen crushed, BMNH.

Hudson 1928, p. 273, pl. xxx fig. 28.

**chirista** Meyrick, §1883c, p. 523; 1884a, p. 27 (*Thamnosara*). Synonymised by Meyrick (1915b, p. 214).

[Dry Bush, Port Hills], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 27/12/82”, “*Thamnosara sublitella* Walker 1/17 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 273, as synonym.

● **Tinearupa** Salmon & Bradley, 1956, p. 66. Type species *Tinearupa sorenseni* Salmon & Bradley, by original designation.

**sorenseni sorenseni** Salmon & Bradley, 1956, p. 66, as species (*Tinearupa*)

Courrejolles Peninsula, Campbell Island, J.H. Sorenson; HT ♂ designated by Salmon & Bradley, NMNZ.  
Dugdale 1971, p. 135, as subspecies.

*sorenseni aucklandiae* Dugdale, 1971b, pp. 135–136

(*Tinearupa*)

Lake Turbott, Adams Island, Auckland Islands, K.A.J. Wise; HT ♂ designated by Dugdale, NZAC.

• *Tingena* Walker, 1864a, pp. 809–810. Type species *Tingena bifaciella* Walker, by original monotypy.

*Cremnogenes* Meyrick, §1883c, p. 525; 1884a, p. 45. Type species *Cremnogenes oxyina* Meyrick, by subsequent designation (Meyrick 1915b, p. 211). Note. This genus includes those species placed under *Borkhausenia* in Hudson 1928, 1939, and 1950.

*actinias* Meyrick, 1901, p. 574 (*Borkhausenia (Oecophora)*) new combination

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “New Zealand GVH /99”, “Borkhausenia armigerella Walk. 13/20 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 264, pl. xxix fig. 12, as variety of *Borkhausenia armigerella* (in the sense of Meyrick, not Walker).

Note. Philpott (1926b, p. 412, fig. 10) illustrates the ♂ genitalia (as *armigerella*).

*affinis* Philpott, 1926a, pp. 391–392, fig. 3 and 4 (*Borkhausenia*) new combination

Nelson NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 262, pl. li fig. 3, as *Borkhausenia affinis*. Note. Philpott's illustration of *affinis* ♂ genitalia is virtually identical with that of *xanthomicta* Meyrick (q.v.).

*afficta* Philpott, 1926b, pp. 401–402 and 411, fig. 9 (*Borkhausenia*) new combination

Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 267, not figured; as *Borkhausenia afficta*.

*aletis* Meyrick, 1905, pp. 235–236 (*Hypercallia*) new combination

Arthur's Pass NC/WD, [3000 ft], E. Meyrick; HT ♂ unique, BMNH.

Hudson 1928, p. 291, not figured, as *Philobota aletis*.

*amiculata* Philpott, 1926b, pp. 402 and 409, fig. 7 (*Borkhausenia*) new combination

Mount Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 265, not figured, as *Borkhausenia amiculata*.

*anaema* Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 42 (*Oecophora*) new combination

Lake Wakatipu OL, E. Meyrick; LT ♂ here designated, labelled “L. Wakatipu New Zealand 15.12.82”,

“Borkhausenia anaema Meyr. 4/4 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 263, not figured, as *Borkhausenia anaema*.

*ancogramma* Meyrick, 1919, pp. 352–353 (*Borkhausenia*) new combination

Wainuiomata WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 268, pl. xlvi fig. 6 (dubious), as *Borkhausenia ancogramma*.

Note. The HT ♂ genitalia differ from those figured by Philpott (1926, p. 409, fig. 7). One ♂ (1/5 in Meyrick's series) is not conspecific on genital characters, and like Hudson's fig. 6 on pl. xlvi has a dark patch on the dorsum just before the tornus which is lacking in the HT.

*latens* Philpott, 1928a, p. 365 (*Borkhausenia*).

Synonymised by Hudson (1939, p. 444).

Rotorua BP, A.J. Turner; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 444, as synonym.

*bellatula* Philpott, 1929a, p. 304 (*Borkhausenia*).

Synonymised by Hudson (1939, p. 444).

Lake Rotoroa BR, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 444, as synonym.

Note. The ♂ genitalia of *latens* HT and *bellatula* HT agree well with those of *ancogramma* HT, and Hudson's synonymy is upheld.

*apanthes* Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 41–42 (*Oecophora*) new combination

[swampy forest], Cambridge WO, E. Meyrick; LT ♂ here designated, labelled “Cambridge New Zealand 14.1.80”, “Borkhausenia apanthes Meyr. 4/5 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 263, pl. xxix fig. 10 (dubious – too yellow for *apanthes*), as *Borkhausenia apanthes*.

*apertella* Walker, 1864a, p. 698 (*Oecophora*) new combination

[Nelson NN], T.R. Oxley; LT ♀ designated as “Type” by H. Durrant, labelled “Oecophora apertella Wkr Cat. Lep. Het. BM 29 p.698 (1864) Type ♀”, “Auckland N. Zeal. 60-73” (circular label), BMNH.

Hudson 1928, p. 264, pl. xxix fig. 19 and 20, as *Borkhausenia apertella*.

Note. The ♂ genitalia figured by Philpott (1926b, p. 410, fig. 8) may or may not be referable to this species.

*aphrontis* Meyrick, §1883c, p. 525; 1884a, p. 46 (*Cremnogenes*) new combination

Arthur's Pass NC/WD, [4000 ft], E. Meyrick; LT ♂ here designated, labelled “Arthurs Pass New Zealand 29.1.83”, “Borkhausenia aphrontis Meyr. 5/14 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 267, pl. xxxviii fig. 22, as *Borkhausenia aphrontis*.

Note. The ♂ genitalia figured by Philpott (1926b, p. 411, fig. 9) are referable to the Mount Arthur NN population, having a longer uncus and differently proportioned valva and apical valval structures.

➤ Oecophoridae, *Tingena*

**armigerella** Walker, 1864a, p. 698 (*Oecophora*) new combination

[Auckland AK], D. Bolton; ?HT ♂ (specimen agreeing with type description, measurement, and stated locality and collector), palpi, metathorax, hindwings, and abdomen missing, BMNH.

Hudson 1928, p. 264, pl. xxix fig. 11 only, as *Borkhausenia armigerella*. Not figured by Philpott.

**bifaciella** Walker, 1864a, p. 810 (*Tingena*) new synonymy

[Auckland AK], D. Bolton; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 264, as synonym of *Borkhausenia apertella*.

Notes. *T. armigerella* is sexually dimorphic in colour pattern, ♀♀ having the head yellow-scaled (sandy-coloured in ♂♂) and only the costal margin blackened (much of the costal cell blackened in ♂♂). Walker and Durrant both regarded *armigerella* HT ♂ as lost, Walker adding a label to that effect, but it is more likely that the specimen relabelled by Durrant as “*Oecophora apertella* Wkr Paratype a” is the *armigerella* HT, misplaced after Walker's examination.

**aurata** Philpott, 1931, p. 32 (*Gymnobathra*) new combination

Opoho DN, C.E. Clarke; HT ♂ designated by Philpott, genitalia in vial, AMNZ.

Hudson 1939, p. 447, pl. lix fig. 27, as *Gymnobathra aurata*; Hudson notes resemblance to “*Borkhausenia* of the *xanthomicta* group”.

**basella** Walker, 1863c, p. 492 (*Incurvaria*) new combination

[Auckland AK], D. Bolton; HT ♂ unique, wings not spread, BMNH.

Hudson 1928, p. 265, pl. xxv fig. 26, as *Borkhausenia basella*.

**ademptella** Walker, 1864a, pp. 698–699 (*Oecophora*). Synonymised by Meyrick (1915b, p. 212).

[Auckland AK], D. Bolton; LT ♂ designated by H. Durrant as “Type”, labelled “*Oecophora ademptella* Wkr Cat. Lep. Het. BM 29: 698 (1864) Type ♂”, “N. Zeal. 54-4” (circular label), hindwings and abdomen missing, BMNH.

**berenice** Meyrick, 1929, pp. 488–489 (*Borkhausenia*) new combination

Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1939, p. 444, pl. lvii fig. 27, as *Borkhausenia berenice*.

**brachyacma** Meyrick, 1909a, p. 13 (*Borkhausenia*) new combination

Invercargill SL, A. Philpott; HT ♂ unique, BMNH.

Hudson 1928, p. 269, pl. xxix fig. 23, as *Borkhausenia brachyacma*.

**amnopis** Meyrick, 1910a, p. 65 (*Borkhausenia*) new synonymy

Invercargill SL, A. Philpott; LT ♀ here designated, labelled “Invercargill New Zealand AP '08”, “*Borkhausenia amnopis* Meyr. 2/2 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 269 in part, as species.

Note. HT ♂ *brachyacma* and LT ♀ *amnopis* have identical colour patterns. The ♂ figured by Hudson (1928, pl. xxix fig. 33) is that of another species, possibly near *T. berenice*. Meyrick (1911b, p. 65) redescribed *amnopis* as a new species.

**chloradelpha** Meyrick, 1905, pp. 239–240 (*Borkhausenia*) new combination

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “Wellington New Zealand GVH /04”, “*Borkhausenia chloradelpha* Meyr. 4/6 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 266, pl. xxx fig. 4, as *Borkhausenia chloradelpha*.

**chloritis** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 36 (*Oecophora*) new combination

Lake Wakatipu OL, E. Meyrick; HT ♂ unique, BMNH. Hudson 1928, p. 271, not figured, as *Borkhausenia chloritis*.

**chrysogramma** Meyrick, §1883c, p. 525; 1884a, pp. 34 (key) and 44–45 (*Oecophora*) new combination [low scrub], Wellington WN, E. Meyrick; HT ♀ unique, abdomen in gelatin capsule, BMNH.

Hudson 1928, p. 261, pl. xxix fig. 6, as *Borkhausenia chrysogramma*.

**clarkei** Philpott, 1928a, p. 366 (*Borkhausenia*) new combination

Kauri Gully, Birkenhead AK, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 445, pl. lix fig. 1, as *Borkhausenia clarkei*.

**collitella** Walker, 1864a, p. 655 (*Gelechia*) new combination

[Auckland AK], D. Bolton; HT ♂ unique, BMNH.

Not mentioned by Hudson. Regarded by Meyrick (1884a, p. 47) as a variety of *griseata*.

Note. Genitalia of HT *collitella* are of the *actinias* (*armigerella* of authors) form, and quite different from those of HT *griseata*.

**compsogramma** Meyrick, 1920a, p. 31 (*Borkhausenia*) new combination

Buller River NN/BR, G.V. Hudson; LT ♂ here designated, labelled “Buller R. New Zealand GVH 12.18”, “*Borkhausenia compsogramma* Meyr. 3/7 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 261, pl. xxix fig. 5, as *Borkhausenia compsogramma*.

Note. On genital and colour pattern differences, Philpott's (1926b, p. 409, fig. 7; p. 410, fig. 8) separation of *compsogramma* and *xanthodesma* Philpott is upheld (see Hudson 1928, p. 261).

***contextella*** Walker, 1864a, p. 656 (*Gelechia*) new combination

[Nelson NN], T.R. Oxley; LT ♂ designated by H. Durrant, labelled “*Gelechia contextella* Wkr Cat. Lep. BM 29 p.656 (1864) Type ♂”, “Auckland N. Zeal. 60-73” (circular label), BMNH.

Hudson 1928, p. 270, as synonym of *Borkhausenia plagiarella*.

Note. HT *crotala* Meyrick and HT *plagiarella* Walker have white-scaled heads. LT *contextella* is sandy-scaled, and is removed from synonymy with *plagiarella* (see Meyrick 1915, p. 213).

***crotala*** Meyrick, 1915b, p. 213 (*Borkhausenia*) new combination

[reserved bush and forest], Dunedin DN, E. Meyrick; LT ♂ here designated, labelled “Dunedin New Zealand 6.1.80”, “*Borkhausenia crotala* Meyr. 2/11 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 270, pl. li fig. 5 and 6, as *Borkhausenia plagiarella* [not of Walker]. Philpott 1926b, p. 410, fig. 8, as *Borkhausenia plagiarella*.

***contextella*** (not of Walker) Meyrick, 1884a, p. 37 (*Oecophora*). Synonymised by Meyrick (1915b, p. 213).

Note. Meyrick proposed the name *crotala* for his conception of Walker's *contextella*.

***decora*** Philpott, 1928a, p. 365 (*Borkhausenia*) new combination

Lake Rotoroa BR, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 442, pl. lix fig. 2, as *Borkhausenia decora*.

***enodis*** Philpott, 1927d, p. 85, fig. 3a,b (*Borkhausenia*) new combination

Cawthron Park, Nelson NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 443, not figured, as *Borkhausenia enodis*.

***epichalca*** Meyrick, 1886a, p. 793 (*Cretnogenes*) new combination

Arthur's Pass NC/WD, [?5000 ft], E. Meyrick; LT ♂ here designated, labelled “Arthurs Pass New Zealand 29.1.83”, “*Borkhausenia epichalca* Meyr. 1/5 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 267, pl. xxxviii fig. 7, as *Borkhausenia epichalca*.

Note. Philpott's (1926b, p. 410, fig. 8) figures of ♂ genitalia agree with the LT genitalia.

***epimylia*** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 36–37 (*Oecophora*) new combination

[in forest], Castle Hill MC, E. Meyrick; LT ♂ here designated, labelled “Castle Hill New Zealand 16.1.83”, “*Borkhausenia epimylia* Meyr. 8/13 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 271, pl. xxix fig. 29, as *Borkhausenia epimylia*.

Note. Philpott's (1926b, p. 409, fig. 7) illustrations of ♂ genitalia of *epimylia* differ considerably in gnathos and valval structure from LT and PLT *epimylia*.

***eriphaea*** Meyrick, 1914a, p. 107 (*Borkhausenia*) new combination

Ben Lomond OL, A. Philpott; LT ♂ here designated, labelled “Ben Lomond New Zealand 25.11.12”, “*Borkhausenia eriphaea* Meyr. 2/2 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 264, pl. xxxviii fig. 15, as *Borkhausenia eriphaea*.

Note. Philpott (1926b, p. 409, fig. 7) figures ♂ genitalia of a topotype.

***eumenopa*** Meyrick, 1926b, p. 416 (*Trachypepla*) new combination

Wainuiomata WN, G.V. Hudson; LT ♂ here designated, labelled “Wainuiomata New Zealand GVH 13/12/23”, “*Trachypepla eumenopa* Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 286, pl. xxx fig. 6, as *Trachypepla eumenopa*.

***metallifera*** Philpott, 1928a, p. 368 (*Trachypepla*). Synonymised by Hudson (1939, p. 449).

Waimarino TO, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 449, as synonym.

***falsiloqua*** Meyrick, 1932, p. 24 (*Trachypepla*) new combination

Waimarino TO, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1939, p. 450, not figured, as *Trachypepla falsiloqua*.

***fenestrata*** Philpott, 1926b, pp. 404 and 410, fig. 8 (*Borkhausenia*) new combination

Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 269, not figured, as *Borkhausenia fenestrata*.

***grata*** Philpott, 1927d, p. 85, fig. 4a,b (*Borkhausenia*) new combination

Dun Mountain NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 443, not figured, as *Borkhausenia grata*.

***griseata*** Butler, 1877, p. 405 (*Oecophora*) new combination

[?Christchurch MC or Dunedin DN], J.D. Enys or J. Hector; LT ♂ labelled as “Type” by Durrant, BMNH.

Hudson 1928, p. 268, not figured, as *Borkhausenia griseata*.

***hastata*** Philpott, 1916, p. 422 (*Borkhausenia*) new combination

Seaward Moss SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 266, pl. xxx fig. 5, as *Borkhausenia hastata*.

» Oecophoridae, *Tingena*

**hemimochla** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 38 (*Oecophora*) new combination  
[Botanic Gardens], Wellington WN, E. Meyrick; LT ♂ here designated, labelled "Wellington New Zealand 31/12/79", "Borkhausenia hemimochla Meyr. 1/11 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 270, pl. xxix fig. 32, as *Borkhausenia hemimochla*.

**homodoxa** Meyrick, §1883c, p. 525; 1884a, pp. 34 (key) and 43 (*Oecophora*) new combination  
[Ben Lomond], Lake Wakatipu OL, E. Meyrick; LT ♂ here designated, labelled "Lake Wakatipu New Zealand 17.12.82", "Borkhausenia homodoxa Meyr. 1/4 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 269, pl. xlvi fig. 14, as *Borkhausenia homodoxa*.

**honestata** Philpott, 1929a, p. 303 (*Borkhausenia*) new combination  
Lake Tekapo MK, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 445, pl. lvii fig. 6, as *Borkhausenia honestata*.  
Note. This species may prove to be better placed in *Trachypepla*.

**honorata** Philpott, 1918, p. 128 (*Borkhausenia*) new combination  
Knife and Steel FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 261, pl. xxxviii fig. 8, as *Borkhausenia honorata*.

**hoplodesma** Meyrick, §1883c, p. 525; 1884a, pp. 34 (key) and 44 (*Oecophora*) new combination  
South Rakaia MC, W.H. Gaze; HT ♂ unique, BMNH.  
Hudson 1928, pp. 261–262, pl. xxix fig. 4, as *Borkhausenia hoplodesma*.

**thranias** Meyrick, 1905, p. 240 (*Borkhausenia*).  
Synonymised by Philpott (1926c, p. 401).  
Whangarei [Heads] ND, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 261, as synonym of *Borkhausenia hoplodesma*.  
Note. HT *thranias* genitalia agree with Philpott's drawing of *hoplodesma* genitalia.

**horaea** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 40–41 (*Oecophora*) new combination  
Bealey River NC, E. Meyrick; LT ♂ here designated, labelled "Bealey River New Zealand 20.1.83", "Borkhausenia horaea Meyr. 4/10 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 263, not figured, as *Borkhausenia horaea*.  
Note. The hindwing anal fold powder puff and genitalia of LT *horaea* are similar to those of *serena*, *comosaris*, and *anaema*. Of the 9 PLTs in Meyrick's *horaea* series, only 3/10, 5/10, and 6/10 are conspecific with the LT;

the rest are referable to *T. actinias* (*armigerella* in the sense of Philpott, 1926b, p. 412, fig. 10.).

**idiogama** Meyrick, 1924b, p. 661 (*Borkhausenia*) new combination

Mount Egmont TK, [3500 ft], G.V. Hudson; LT ♂ here designated, labelled "Mt Egmont New Zealand GVH 1.23", "Borkhausenia idiogama Meyr. 1/3 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 266, pl. xxix fig. 31, as *Borkhausenia idiogama*.

**innotella** Walker, 1864a, p. 652 (*Gelechia*) new combination

[Auckland AK], D. Bolton; LT ♀ selected by H. Durrant and designated as "Type ♂", labelled "Gelechia innotella Wkr Cat. Lep. BM 29 p.652 (1864) Type ♂", "New Zeal. 54-4" (circular label), BMNH.

Hudson 1928, p. 269, pl. xxix fig. 24 (as *politis*), as *Borkhausenia innotella*.

Note. Hudson's pl. xxix fig. 21 refers to *ombrodella* (q.v.).

**monospilella** Walker, 1864a, p. 653 (*Gelechia*).

New synonymy

[Auckland AK], D. Bolton; HT ♂ unique, abdomen missing, BMNH.

Not mentioned by Hudson or Meyrick.

Note. HT *monospilella* is externally indistinguishable from PLT ♂ *innotella*, and is here synonymised.

**politis** Meyrick, 188e, p. 81 (*Oecophora*). New synonymy

Wellington WN, G.V. Hudson; HT ♀ not found in Meyrick Collection, BMNH.

Hudson 1928, p. 265, pl. xxix fig. 24, as *Borkhausenia politis*.

Note. It is doubtful – because of the emphasis in the original description on the "whitish-ochreous" ground colour – if HT *politis* and Meyrick's and Hudson's later sense of this taxon are the same thing (cf. *Planotortrix coprosmae*, Tortricidae, p. 125). Philpott's (1926b, p. 410, fig. 8) figures of the genitalia of *politis* of authors agree excellently with post-1887 material in Meyrick's series from Hudson in Wellington, collected in 1900, 1910, and 1917; these specimens are not whitish-ochreous, however, but brown.

**lassa** Philpott, 1930b, p. 9 (*Borkhausenia*) new combination

Leith DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 443, pl. lvii fig. 30, as *Borkhausenia lassa*.

**laudata** Philpott, 1930b, p. 9 (*Borkhausenia*) new combination

Bluecliff FD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, pp. 443–444, pl. lvii fig. 28, as *Borkhausenia laudata*.

**letharga** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 35–36 (*Oecophora*)

Dunedin DN, E. Meyrick; LT ♂ here designated, labelled "Dunedin New Zealand 6.1.80", "Borkhausenia leth-

arga Meyr. 1/3 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 271, pl. lii fig. 3, as *Borkhausenia letharga*.

***levicula*** Philpott, 1930b, p. 8 (*Borkhausenia*) new combination

Flat Top Mountain, Hunter Mountains FD, C.E. Clarke; HT ♀ designated by Philpott, AMNZ.  
Hudson 1939, p. 442, pl. lix fig. 2, as *Borkhausenia levicula*.

***loxotis*** Meyrick, 1905, p. 241 (*Borkhausenia*) new combination

Wellington WN, E. Meyrick; LT ♂ here designated, labelled "Wellington New Zealand 3.1.86", "Borkhausenia loxotis Meyr. 1/8 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 261, pl. xxix fig. 3 (a poor representation, with the wings too pointed and the black outlining of the yellow areas missing), as *Borkhausenia loxotis*.

Note. LT and PLT ♂ genitalia do not agree with Philpott's (1926b, p. 410, fig. 8) illustration: the uncus is longer than Philpott shows, and sinuous, as is the valval costal process.

***macarella*** Meyrick, §1883c, p. 524; 1884a, p. 43 (*Oecophora*) new combination

[steep scrubby hills, Port Lyttelton] MC, E. Meyrick; LT ♂ here designated, labelled "Christchurch New Zealand 8.1.80", "Borkhausenia macarella Meyr. 8/13 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 263, not figured, as *Borkhausenia macarella*.

***maranta*** Meyrick, 1886a, p. 791 (*Oecophora*) new combination

Invercargill SL, A. Philpott; HT ♂ unique, BMNH.  
Hudson 1928, p. 263, pl. xxix fig. 1, as *Borkhausenia maranta*.

Note. Philpott's (1926b, p. 412, fig. 10) illustration of ♂ genitalia agrees with HT.

***marcida*** Philpott, 1927a, p. 706, fig. 2 (*Borkhausenia*) new combination

Bottle Lake, Christchurch MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.  
Hudson 1928, p. 269, not figured, as *Borkhausenia marcida*.

***melanamma*** Meyrick, 1905, pp. 240–241 (*Borkhausenia*) new combination

[Ida Valley CO], J.H. Lewis; LT ♂ here designated, labelled "Dunedin New Zealand JHL -/04", "Borkhausenia melanamma Meyr. 4/8 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 262, pl. xxix fig. 2, as *Borkhausenia melanamma*.

Note. LT *melanamma* genitalia differ from those illustrated by Philpott (1926b, p. 412, fig. 10).

***sabulosa*** Philpott, 1918, pp. 128–129 (*Borkhausenia*) new synonymy

"Central Otago" CO, J.H. Lewis; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 262, as ?synonym of *Borkhausenia melanamma*.

Note. Philpott (1926b, p. 412, fig. 10) misapplied *melanamma*. HT *sabulosa* and LT *melanamma* are indistinguishable on colour pattern and genitalia, and are here synonymised.

***melinella*** Felder & Rogenhofer, 1875, pl. cxl fig. 41 (*Oecophora*) new combination

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.

Not mentioned by Hudson.

Note. Meyrick (1884a, p. 48) states: "This figure appears to me insufficient for identification".

***freta*** Philpott, 1926b, pp. 402 and 411, fig. 10 (*Borkhausenia*). New synonymy.

Nelson NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 264, pl. xlvi fig. 2, as species.

Note. The Philpott PT ♀ ♀ at BMNH are identical with HT *melinella* on external characters, as is the HT ♂ *freta* in NZAC, and *freta* is here synonymised.

***monodonta*** Meyrick, 1911b, p. 75 (*Cremnogenes*) new combination

Mount Holdsworth WN, R.M. Sunley; LT ♂ here designated, labelled "Mt Holdsworth New Zealand RMS 11.09", "Borkhausenia monodonta Meyr. 1/15 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 267, pl. xxxviii fig. 10, as *Borkhausenia monodonta*.

***nigra*** Philpott, 1914, p. 120 (*Cremnogenes*). Synonymised by Meyrick (1915b, p. 213).

Ben Lomond OL, A. Philpott; LT ♂ here designated, labelled "Ben Lomond 25/11/12", "Lectotype ♂ *Cremnogenes nigra* Philpott", NZAC.

Hudson 1928, p. 267, as synonym.

Note. The 5 ♂♂ from Ben Lomond collected on 25 November 1911 by Philpott in Philpott's collection under *Borkhausenia monodonta* are presumed to be his type series of *nigra*. His copy of *Trans. proc. N.Z. Inst.* vol. 46, p. 120, has "nigra" crossed out and "monodonta" written in pencil, in Philpott's handwriting.

***morosa*** Philpott, 1926b, pp. 403–404 and 410, fig. 8 (*Borkhausenia*) new combination

Dun Mountain NN/MB, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 270, not figured, as *Borkhausenia morosa*.

***nycteris*** Meyrick, 1890, p. 219 (*Oecophora*) new combination

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 11/88", "Borkhausenia nycteris Meyr. 8/19 E. Meyrick det. in Meyrick Coll.", BMNH.

» Oecophoridae, *Tingena nycteris*

Hudson 1928, p. 268, pl. xxix fig. 34 and 35, as *Borkhausenia nycteris*.

Note. LT *nycteris* ♂ genitalia have the costal valval process longer than is shown by Philpott (1926b, p. 412, fig. 10). Meyrick (1911b, pp. 63–64) described the ♀, and noted that his original description refers only to the ♂.

***ombrodella*** Hudson, 1950, p. 107, as variety (*Borkhausenia*) new combination

Whakapapa, Mount Ruapehu TO, G.V. Hudson; LT ♂ here designated, labelled “380DD”, [“Whakapapa 9–10.1.1941 GVH” in Hudson’s Register], NMNZ.

Hudson 1928, p. 269, pl. xxix fig. 21, as *Borkhausenia innotella*; 1950, p. 107, pl. iii fig. 10; as *Borkhausenia ombrodella*.

Note. Topotypic ♂♂ in BMNH have genitalia of the shape figured by Philpott (1926b, p. 411, fig. 9) for *innotella*.

***opaca*** Philpott, 1926b, pp. 403 and 409, fig. 7 (*Borkhausenia*) new combination

Bluff SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 265, pl. xxix fig. 15, as *Borkhausenia opaca*.

***ophiodryas*** Meyrick, 1936, p. 282 (*Borkhausenia*) new combination

Little River, Banks Peninsula MC, S. Lindsay; HT ♀ designated by Meyrick, CMNZ.

Hudson 1939, p. 443, pl. lxi fig. 14, as *Borkhausenia ophiodryas*.

***oporaea*** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 40 (*Oecophora*) new combination

Castle Hill MC, E. Meyrick; LT ♂ here designated, labelled “Castle Hill N. NEW ZEALAND 16.i.1883 Meyrick. Raynor Coll. No. 10,632”, BMNH.

Hudson 1928, p. 264, as synonym of *Borkhausenia apertella*.

Note. Only the LT ♂ and 2 PLT ♀♀ remain of Meyrick’s series. Others in the series under the name *oporaea* differ in genitalia and scale colour from LT.

***oxyina*** Meyrick, §1883c, p. 525; 1884a, pp. 45–46 (*Cremnogenes*) new combination

[?Kinloch or Paradise], Lake Wakatipu OL, E. Meyrick; LT ♂ here designated, labelled “L. Wakatipu New Zealand 16.12.82”, “Borkhausenia oxyina Meyr. 8/10 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 267, pl. xlvi fig. 6 (a very pale representation), as *Borkhausenia oxyina*.

***pallidula*** Philpott, 1924a, p. 210 (*Borkhausenia*) new combination

Gouland Downs NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 271, pl. I fig. 9, as *Borkhausenia pallidula*.

***paratrimma*** Meyrick, 1910a, p. 65 (*Borkhausenia*) new combination

Invercargill SL, A. Philpott; LT ♂ here designated, labelled “Invercargill New Zealand AP .08”, “Borkhausenia paratrimma Meyr. 2/4 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 262, pl. xxix fig. 22, as *Borkhausenia paratrimma*.

Note. Meyrick (1911b, p. 63) redescribed the two Philpott specimens, as *Borkhausenia paratrimma* n.sp.

***paula*** Philpott, 1927a, p. 707, fig. 1 (*Borkhausenia*) new combination

Pukeatua Bush, Banks Peninsula MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 262, pl. lii fig. 22, as *Borkhausenia paula*.

***pentalea*** Meyrick, 1905, p. 239 (*Borkhausenia* (*Oecophora*)) new combination

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “Wellington New Zealand GVH /00”, “Borkhausenia pentalea Meyr. 2/4 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 269, pl. xxix fig. 13, as *Borkhausenia pentalea*.

***perichlora*** Meyrick, 1907c, p. 118 (*Borkhausenia*) new combination

Invercargill SL, A. Philpott; LT ♂ here designated, labelled “Invercargill New Zealand AP .06”, “Borkhausenia perichlora Meyr. 1/4 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 264, pl. xxix fig. 14, as *Borkhausenia perichlora*.

***pharmactis*** Meyrick, 1905, p. 241 (*Borkhausenia*) new combination

Mount Arthur Tableland NN, E. Meyrick; HT ♀ unique, BMNH.

Hudson 1928, p. 264, pl. I fig. 26, as *Borkhausenia pharmactis*.

***phegophylla*** Meyrick, §1883c, p. 524; 1884a, pp. 34 (key) and 39 (*Oecophora*) new combination

[?Kinloch or Paradise], Lake Wakatipu OL, E. Meyrick; LT ♂ here designated, labelled “L. Wakatipu New Zealand 16.12.82”, “Borkhausenia phegophylla Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 265, pl. xxxviii fig. 9, as *Borkhausenia phegophylla*.

Note. LT *phegophylla* ♂ genitalia agree with Philpott’s (1926b, p. 411, fig. 9) illustration of “*basella*”.

***plagiatella*** Walker, 1863c, p. 485 (*Tinea*) new combination

[Auckland AK], D. Bolton; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 270 in part, not figured, as *Borkhausenia plagiatella*.

***pronephela*** Meyrick, 1907c, p. 119 (*Borkhausenia*)

**new combination**

Invercargill SL, A. Philpott; LT ♂ here designated, labelled “Invercargill New Zealand AP .06”, “Borkhausenia pronephela Meyr. 1/3 E. Meyrick det. in Meyrick Coll.”, palpi missing, BMNH.

Hudson 1928, p. 266, pl. xxix fig. 28, as *Borkhausenia pronephela*.

***robiginosa*** Philpott, 1915, p. 200 (*Borkhausenia*)  
**new combination**

Longwood Range SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 268, pl. xxix fig. 36.

Note. Philpott (1926b, p. 409, fig. 7) figures the ♂ genitalia.

***seclusa*** Philpott, 1921, p. 340 (*Borkhausenia*)  
**new combination**

Lake Luna OL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 270, pl. xxix fig. 16, as *Borkhausenia seclusa*.

Note. PT ♂ genitalia figured by Philpott (1926b, p. 411, fig. 9).

***serena*** Philpott, 1926b, pp. 403 and 410, fig. 8  
(*Borkhausenia*)  
**new combination**

“Sunnyside”, Waiau [River] FD/SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 269, pl. li fig. 10, as *Borkhausenia serena*.

***comosaris*** Meyrick, 1931a, p. 95 (*Borkhausenia*).  
**New synonymy.**

Gollan's Valley, Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1939, p. 445, not figured, as species.

Note. Although HT ♂ *serena* lacks an abdomen, scale pattern and hindwing anal puff scales are identical with those of HT ♂ *comosaris*. The ♂ genitalia of *serena* figured by Philpott (1926b, p. 410, fig. 8) are likewise as for HT *comosaris*.

***siderodeta*** Meyrick, §1883c, p. 525; 1884a, pp. 34  
(key) and 43–44 (*Oecophora*)  
**new combination**

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 13.1.83”, “Borkhausenia siderodeta Meyr. 7/26 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 262, pl. xxix fig. 9, as *Borkhausenia siderodeta*.

***siderota*** Meyrick, 1888e, p. 82 (*Cremnogenes*)  
**new combination**

Mount Arthur NN, [on *Aciphylla* flowers], E. Meyrick; LT ♀ selected by D.J. Carter and here designated, labelled “Mt Arthur New Zealand 16.1.86”, “Borkhausenia siderota Meyr. 1/10 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 267, pl. xxix fig. 27, as *Borkhausenia siderota*.

***sinuosa*** Philpott, 1928g, pp. 488 and 486, fig. 14  
(*Borkhausenia*)  
**new combination**

Botanical Gardens, Wellington WN, A. Philpott; HT ♂ designated by Philpott, wings damaged, abdomen on Philpott genitalia slide, NZAC.

Hudson 1939, p. 443, not figured, as *Borkhausenia sinuosa*.

***tephrophanes*** Meyrick, 1929, p. 488 (*Borkhausenia*)  
**new combination**

Flora Creek, Mount Arthur NN, G.V. Hudson; HT ♀ unique, abdomen and right hindwing missing, BMNH.

Hudson 1939, p. 444, pl. lix fig. 7, as *Borkhausenia tephrophanes*.

***terrena*** Philpott, 1926a, p. 392, fig. 5 and 6 (*Borkhausenia*)  
**new combination**

Queenstown OL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 262, as ?synonym of *Borkhausenia melanamma*.

***thalerodes*** Meyrick, 1916b, p. 416 (*Borkhausenia*)  
**new combination**

Arthur's Pass NC/WD, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 268, pl. xxxviii fig. 11, as *Borkhausenia thalerodes*.

***vestita*** Philpott, 1926a, p. 392 (*Borkhausenia*)  
**new combination**

Hunter Mountains FD, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 266, pl. li fig. 4, as *Borkhausenia vestita*.

***xanthodesma*** Philpott, 1923, pp. 151–152 (*Borkhausenia*)  
**new combination**

Tisbury SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 261, as synonym of *Borkhausenia compsogramma*.

Note. Genitalia of LT *compsogramma* do not resemble those of HT or PT ♂ *xanthodesma*.

***xanthomicta*** Meyrick, 1916b, p. 415 (*Borkhausenia*)  
**new combination**

Wellington WN, G.V. Hudson; LT ♀ here designated, labelled “Wellington New Zealand GVH .15”, “Borkhausenia xanthomicta Meyr. 4/8 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, pp. 262–263, pl. xxix fig. 8.

Note. A PLT ♂ (“8/8”) from Wellington WN has genitalia very similar to those figured for *affinis* Philpott by Philpott (1926b, p. 392, fig. 3 and 4) and agreeing with “*xanthomicta*” as figured by Philpott (1926b, p. 412, fig. 10).

Also 4 undescribed species (NZAC).

● ***Trachypepla*** Meyrick, 1883a, p. 423 (key); 1883c, pp. 367–368. Type species *Trachypepla euryleucota*

► Oecophoridae, *Trachypepla*

Meyrick, by subsequent designation (Meyrick 1915b, p. 216).

**amphileuca** Meyrick, 1914a, pp. 107–108 (*Trachypepla*)

Wainuiomata WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 284, pl. xxxi fig. 9.

**anastrella** Meyrick, §1883c, p. 523; 1883e, pp. 368 (key) and 370 (abbreviated description); 1884a, pp. 19–20 (full description) (*Trachypepla*)

[reserved bush and forest], Dunedin DN, E. Meyrick; LT ♂ here designated, labelled “Dunedin New Zealand 6/1/80”, “*Trachypepla anastrella* Meyr. 5/7 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 286; 1939, p. 449, pl. lxi fig. 6.

**angularis** Philpott, 1929a, p. 303, fig. 1 (*Borkhausenia*) new combination

Lake Rotoroa BR, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 445, pl. lvii fig. 8 (type specimen), as *Borkhausenia angularis*.

Note. The basally strongly arcuate aedeagus and the valval character of *angularis* accord well with *Trachypepla* species as figured by Philpott (1927f, pp. 106 and 108, fig. 20–28). The species is therefore transferred.

**aspidephora** Meyrick, §1883c, p. 523; 1883e, pp. 368 (key) and 370 (abbreviated description); 1884a, p. 19 (full description) (*Trachypepla*)

[Dry Bush, Port Hills], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 27/12/82”, “*Trachypepla aspidephora* Meyr. 1/10 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 285, pl. xxxi fig. 24.

**conspicuella** Walker, 1864a, p. 651 (*Gelechia*)

[Nelson NN], T.R. Oxley; LT ♂ selected by H. Durrant, labelled “*Gelechia conspicuella* Wkr Cat. Lep. BM 29: 651 (1864) Type ♂”, “Auckland N. Zeal. 60.73” (circular label), abdomen missing, BMNH.

Hudson 1928, pp. 283–284, pl. xxxi fig. 8.

**taongella** Felder & Rogenhofer, 1875, pl. cxl fig. 45 (*Gelechia*). Synonymised by Meyrick (1884a, p. 14).

[Nelson NN, T.R. Oxley]; HT ♂ unique, irrelevant abdomen glued on, BMNH.

**contritella** Walker, 1864a, p. 657 (*Gelechia*)

[Nelson NN], T.R. Oxley; HT ♀ unique, head, right forewing, and abdomen missing, BMNH.

Hudson 1928, p. 285, pl. xxxi fig. 5.

**nyctopis** Meyrick, §1883c, p. 522; 1883e, pp. 368 (key) and 369 (abbreviated description); 1884a, pp. 16–17 (full description) (*Trachypepla*). Synonymised by Meyrick (1915b, p. 217).

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ selected by P.A. Brown, labelled “Christchurch New Zealand 13/1/83”, “*Trachypepla contritella* Meyr. 1/15 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 285, as synonym.

Note. This synonymy may not stand close scrutiny; there are differences in uncus and valva structures between topotypic *contritella* and *nyctopis*.

**cypbonias** Meyrick, 1927b, p. 314 (*Trachypepla*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1939, p. 450, pl. lix fig. 14.

Note. Hudson indicated that the species was found “on the hills on the eastern side of Wellington Harbour”.

**euryleucota** Meyrick, §1883c, p. 522; 1883e, pp. 368 (key and abbreviated description); 1884a, pp. 14–15 (full description) (*Trachypepla*)

[Botanic Gardens and forest], Wellington WN, E. Meyrick; LT ♂ selected and designated by Clarke (1963, p. 461, pl. 227 fig. 1–1d); genitalia on slide JFGC 9601, right forewing and hindwing on slide, BMNH.

Hudson 1928, p. 283, pl. xxxi fig. 10.

**festiva** Philpott, 1930b, p. 10 (*Trachypepla*)

Whangarei Falls ND, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 449, pl. lix fig. 6.

**polyleuca** Meyrick, 1931a, pp. 95–96 (*Trachypepla*). Synonymised by Hudson (1939), p. 449.

Whangarei ND, S.C. Patterson; LT ♂ here designated, labelled “Whangarei New Zealand GVH .30”, “*Trachypepla polyleuca* Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1939, p. 449, as synonym.

**galaxias** Meyrick, §1883c, p. 522; 1883e, pp. 368 (key) and 369 (abbreviated description); 1884a, pp. 17–18 (full description) (*Trachypepla*)

[heath-like scrub and swamp], Hamilton WO, E. Meyrick; LT ♂ here designated, labelled “Hamilton New Zealand 17/1/80”, “*Trachypepla galaxias* Meyr. 6/8 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 284, pl. xxxi fig. 28.

**hieropis** Meyrick, 1892, p. 218 (*Trachypepla*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 284, pl. xxxi fig. 3.

**importuna** Meyrick, 1914a, p. 108 (*Trachypepla*)

Ohakune RI/TO, G.V. Hudson; LT ♂ here designated, labelled “Ohakune New Zealand GVH 1.12”, “*Trachypepla importuna* 2/4 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 286, not figured.

**indolescens** Meyrick, 1927a, p. 700 (*Trachypepla*)

Karori WN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 286, pl. 1 fig. 14.

**ingenua** Meyrick, 1911b, pp. 65–66 (*Trachypepla*)  
Otira River WD, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 284, pl. xxxi fig. 4.

**leucoplanetis** Meyrick, §1883c, p. 522; 1883e, p. 368  
(key, abbreviated description); 1884a, p. 14 (full  
description) (*Trachypepla*)  
[dense forest], Hamilton WO, E. Meyrick; LT ♂ here des-  
ignated, labelled “Hamilton New Zealand 18/1/80”,  
“Trachypepla leucoplanetis Meyr. 1/6 E. Meyrick det.  
in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 283, pl. xxxi fig. 11.

**lichenodes** Meyrick, §1883c, p. 523; 1883e, pp. 368  
(key) and 370 (abbreviated description); 1884a,  
pp. 20–21 (full description) (*Trachypepla*)  
Bealey River NC, E. Meyrick; HT ♀ unique, BMNH.  
Hudson 1928, p. 286, pl. xxxi fig. 25 and 26.

**minuta** Philpott, 1931, p. 33 (*Trachypepla*)  
Auckland Domain AK, A. Philpott; HT ♂ designated by  
Philpott, AMNZ.  
Hudson 1939, p. 450, pl. lix fig. 13.

**nimbosa** Philpott, 1930b, p. 10 (*Trachypepla*)  
Kauri Gully, Birkenhead AK, C.E. Clarke; HT ♂ desig-  
nated by Philpott, AMNZ.  
Hudson 1939, pp. 449–450, pl. lix fig. 12.

**ocneropis** Meyrick, 1936, pp. 282–283 (*Trachy-  
pepla*; as *oneropis* – printer’s error according to  
Meyrick 1937, p. 120)  
Nelson NN, E. Meyrick; LT ♂ here designated, labelled  
“Nelson New Zealand 12/1/86”, “Trachypepla ocner-  
opis Meyr. 3/5 E. Meyrick det. in Meyrick Coll.”,  
BMNH.  
Hudson 1939, p. 450, pl. lxii fig. 16, as *ocneropis*.  
Note. *T. ocneropis* LT ♂ and *Mermeristis spodiaea* (p.  
97) are virtually indistinguishable.

**photinella** Meyrick, 1883a, pp. 512 (key) and 541–  
542 (*Eulechria*)  
[Botanic Gardens], Wellington WN, E. Meyrick; HT ♂  
unique, abdomen missing, BMNH.  
Hudson 1928, p. 286, pl. xlvi fig. 8, as *Trachypepla pho-  
tinella*, after Meyrick (1918a, p. 134).

**protochlora** Meyrick, §1883c, p. 522; 1883e, pp. 368  
(key) and 370 (abbreviated description); 1884a,  
p. 18 (full description) (*Trachypepla*)  
Otira Gorge [and valley] WD, E. Meyrick; LT ♂ here des-  
ignated, labelled “Otira Gorge New Zealand  
24/1/83”, “Trachypepla protochlora Meyr. 2/9 E.  
Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 285, pl. xxxi fig. 23.


**roseata** Philpott, 1923, p. 152 (*Trachypepla*)  
Dun Mountain NN/MB, A. Philpott; HT ♂ designated  
by Philpott, NZAC.  
Hudson 1928, p. 285, pl. xlvi fig. 22.

**semilauta** Philpott, 1918, p. 129 (*Trachypepla*)  
Hunter Mountains FD, A. Philpott; HT ♀ designated by  
Philpott, NZAC.  
Hudson 1928, p. 283, pl. xxviii fig. 18.

**spartodeta** Meyrick, §1883c, p. 522; 1883e, pp. 368  
(key) and 369 (abbreviated description); 1884a,  
p. 16 (full description) (*Trachypepla*)  
[Botanic Gardens], Wellington WN, E. Meyrick; HT ♀  
unique, BMNH.  
Hudson 1928, p. 284, pl. xxxi fig. 2.

—◎—

**Subfamily STATHMOPODINAE**  
(in the sense of Hedges 1978, pp. 7–9)



**Oecophoridae: Stathmopodinae**  
(159) *Stathmopoda skelloni* Butler

• ***Stathmopoda*** Herrich-Schaeffer, [1853], p. 54, in the sense of Kasy (1973, p. 251 – full synonymy). Type species *Phalaena Tinea pedella* Linnaeus, by secondary monotypy (Herrich-Schaeffer [1853], p. 283); Palearctic.

***Boocara*** Butler, 1880, p. 562. Type species *B. skelloni* Butler, by original monotypy. Synonymised by Meyrick (1889b, p. 169 – “a grotesquely soecistic generic name”); Kasy (1973, p. 251).

***albimaculata*** Philpott, 1931, pp. 33–34 (*Stathmopoda*)  
Woodside [Mt Maungatua], Taieri DN, C.E. Clarke; HT ♀ designated by Philpott, AMNZ.  
Hudson 1939, p. 456, pl. lx fig. 6.

***aposema*** Meyrick, 1901, p. 575 (*Stathmopoda*)  
Auckland AK, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 300, pl. xxxviii fig. 6.

***aristodoxa*** Meyrick, 1926b, p. 416 (*Stathmopoda*)  
Gollan's Valley, Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 298, pl. li fig. 11.

***caminora*** Meyrick, 1890, p. 219 (*Stathmopoda*)  
Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 298, pl. xxxii fig. 18.

***campylocha*** Meyrick, 1889b, p. 168 (*Stathmopoda*)  
Dunedin DN, E. Meyrick; LT ♀ here designated, labelled  
“Dunedin New Zealand 8/2/82”, “*Stathmopoda campylocha* Meyr. 1/5 E. Meyrick det. in Meyrick Coll.”,  
BMNH.  
Hudson 1928, p. 299, not figured; 1939, p. 456, pl. lx fig. 5.

***coracodes*** Meyrick, 1923, p. 167 (*Stathmopoda*)  
Picton SD, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, pp. 299–300, pl. xlvi fig. 23.

***distincta*** Philpott, 1923, p. 152 (*Stathmopoda*)  
Dun Mountain NN/MB, A. Philpott; HT ♀ designated by Philpott, NZAC.  
Hudson 1928, p. 298, pl. xxxii fig. 19.

***endotherma*** Meyrick, 1931b, p. 368 (*Stathmopoda*)  
Little River, Banks Peninsula MC, S. Lindsay; HT ♀ designated by Meyrick, CMNZ.  
Hudson 1939, p. 456, pl. lx fig. 7.

***holochra*** Meyrick, 1889b, p. 168 (*Stathmopoda*)  
[Botanic Gardens], Wellington WN, E. Meyrick; HT ♀ unique, BMNH.  
Hudson 1928, p. 299, pl. xxxii fig. 20.

***horticola*** Dugdale, new species for *skelloni* of authors, but not Butler (*Stathmopoda*)

Tauranga BP, P. Brown; HT ♂ designated by Dugdale, labelled “New Zealand BP Tauranga Caine's Orchard 2 July 1982 P. Brown”, “reared Actinidium fruit” (yellow card), “Holotype ♂ *Stathmopoda horticola Dugdale*” (red card), NZAC.  
Hudson 1928, p. 299, pl. xxxii fig. 16, as *Stathmopoda skelloni*.

Note. Seven of the specimens in the Meyrick Collection under “*skelloni*” (in the sense of Meyrick) are of this species, which differs in ♂ and ♀ colour pattern and genitalia from *skelloni* Butler.

***melanochra*** Meyrick, 1897a, p. 321 (*Stathmopoda*)  
“Tasmania”, E. Meyrick; LT (?gender) selected by I.F.B. Common, BMNH.  
New Zealand: adventive; introduced for biological control of *Eriococcus coriaceus* (Homoptera: Eriococcidae). Not mentioned by Hudson.

***mysteriastis*** Meyrick, 1901, p. 575 (*Stathmopoda*)  
Auckland AK, E. Meyrick; HT ♂ unique, abdomen separate, BMNH.

Hudson 1928, p. 300, pl. xxxviii fig. 5.  
Note. Hudson's illustration is of a southern specimen; it differs from the HT in that the white area shown should be yellowish, and the dark area more ochreous.

***seminuda*** Philpott, 1917b, p. 244 (*Stathmopoda*). Synonymised by Meyrick (1923, p. 167).  
Broad Bay, Dunedin DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1928, p. 300, as synonym.

***plumbiflua*** Meyrick, 1911b, p. 75 (*Stathmopoda*)  
West Plains, Invercargill SL, A. Philpott; LT ♂ here designated, labelled “Invercargill New Zealand AP 21.1.09”, “*Stathmopoda plumbiflua* Meyr. 1/2 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 300, pl. xxxii fig. 21.

***skelloni*** Butler, 1880, p. 562 (*Boocara*)  
[Blenheim] MB, W. Skellon; HT ♀ designated by Butler, hindwings and hindlegs in gelatin capsule, BMNH.  
Hudson 1928, p. 299, pl. xxxii fig. 17, as *Stathmopoda phlegyra*.

***phlegyra*** Meyrick, 1889b, p. 168 (*Stathmopoda*).  
New synonymy.

“Taranaki” [New Plymouth] TK, E. Meyrick; LT ♂ here designated, labelled “Taranaki New Zealand 28/2/83”, “*Stathmopoda phlegyra* Meyr. 17/19 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 299, pl. xxxii fig. 17.

***fusilis*** Meyrick, 1914a, p. 111 (*Stathmopoda*).  
Synonymised (with *phlegyra*) by Meyrick (1921, p. 335).

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 299, as synonym of *Stathmopoda phlegyra*.

***trimolybdias*** Meyrick, 1926b, p. 416 (*Stathmopoda*)

Ashhurst, Manawatu River WI, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 299, pl. xlvi fig. 21.

Also 2 undescribed species (NZAC).

• ***Calicotis*** Meyrick, 1889b, p. 170. Type species *Calicotis crucifera* Meyrick, by original monotypy.

***crucifera*** Meyrick, 1889b, pp. 170–171 (*Calicotis*)

“Taranaki” [New Plymouth] TK, E. Meyrick; LT ♀ designated by Kasy (1973, p. 244), BMNH.

Hudson 1928, p. 297, pl. xxxii fig. 13.

• ***Thylacosceles*** Meyrick, 1889b, p. 171. Type species *Thylacosceles acridomima* Meyrick, by original monotypy.

***acridomima*** Meyrick, 1889b, p. 171 (*Thylacosceles*)  
[Botanic Gardens and forest], Wellington WN, E. Meyrick; HT ♂ unique, BM genitalia slide no. 15341 ♂, BMNH.

Hudson 1928, p. 300, pl. xxxiv fig. 4 and 5.

Note. All 4 Meyrick Collection specimens are ♂♂ from Wellington. Hudson’s collection (NMNZ) has both sexes from Wellington, with the ♀ indistinguishable from LT *epichlora*.

***epichlora*** Meyrick, 1889b, p. 169–170 (*Stathmopoda*). New synonymy.

Otira Gorge WD, E. Meyrick; LT ♀ here designated, labelled “Otira Gorge New Zealand 24/1/83”, “*Pachyrhabda epichlora* Meyr. 1/6 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 300, not figured, as *Pachyrhabda epichlora* (transferred to *Pachyrhabda* by Meyrick 1915b, p. 222).


Note. All 5 PLTs (from Auckland AK, Wellington WN, and Otira WD) are ♀, and as all agree excellently with Hudson’s illustration of a Wellington ♀ *acridomima*, the species is synonymised with *acridomima*.

***radians*** Philpott, 1918, p. 129 (*Thylacosceles*)  
Seaward Bush, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 301, pl. xxxviii fig. 16 and 17.

## Subfamily STENOMATINAE

(in the sense of Hedges 1978, pp. 7–9)



### Oecophoridae: Stenomatinae

(160) ***Agriophara colligatella*** (Walker)

• ***Agriophara*** Rosenstock, 1885, p. 439. Type species *Agriophara cinerosa* Rosenstock; Australia.

***Hypeuryntis*** Meyrick, 1879b, p. 389. Type species *Hypeuryntis coricopa* Meyrick. Synonymised by Meyrick (1915b, p. 221).

***colligatella*** Walker, 1864a, p. 768 (*Cryptolechia*) new combination, resurrected name


[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1928, p. 288, not figured, as *Atomotricha colligatella*, after Meyrick (1915b, p. 218).

***coricopa*** Meyrick, 1897b, p. 389–390 (*Hypeuryntis*), by original monotypy. New synonymy.

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 296, pl. xxv fig. 11 and 12, as *Agriophara coricopa*.



**Subfamily XYLORYCTINAE**  
(in the sense of Hodges 1978, pp. 7–9)



**Oecophoridae: Xylorictinae**  
(161) *Donacostola notabilis* (Philpott)

• ***Donacostola*** Meyrick, 1931a, p. 95. Type species *Euprionocera notabilis* Philpott, by original designation.

***notabilis*** Philpott, 1928a, pp. 368–369  
(?*Euprionocera*)

Flora River NN, A. Philpott; HT ♀ designated by Philpott, NZAC.  
Hudson 1939, p. 454, pl. lviii fig. 15, as *Donacostola notabilis*.


• ***Scieropepla*** Meyrick, §1885i, p. 590; 1886b, p. 165.  
Type species *Scieropepla typhicola* Meyrick, by original monotypy.

***typhicola*** Meyrick, §1885i, p. 590 (as *typicola*);  
1886b, p. 165 (as *typhicola*, in *Scieropepla*)  
Christchurch MC, E. Meyrick; 3 STs (1 ♂, 2 ♀ ♀) “Fere-day Collection”, CMNZ.

Hudson 1928, pp. 295–296, pl. xxxii fig. 15.  
Note. This is another instance of a printer's error in the  
*N.Z. Journal of Science* (Dunedin).

—◎—

**Family SCYTHRIDIDAE**  
(in the sense of Hodges 1978, pp. 7–9)



**Scythrididae**  
(162) *Scythris epistrota* Meyrick

• ***Scythris*** of authors, in the sense of Meyrick (1915b, p. 210)

***epistrota*** Meyrick, 1889b, p. 161 (*Butalis*).  
[Port Hills]. Christchurch MC, E. Meyrick; LT ♂ here  
designated, labelled “Christchurch New Zealand  
4/1/80”, “Scythris epistrota Meyr. 6/15 E. Meyrick det.  
in Meyrick Coll.”, BMNH.

Hudson 1928, p. 320, but not description or pl. xxviii fig.  
13, which refers to an undescribed species from Mount  
Arthur NN.

***lacustris*** Philpott, 1930a, p. 249 (*Elachista*).  
Synonymised by Meyrick (1931b, p. 369).  
Lake Pukaki MK, S. Lindsay; HT ♂ designated by Philpott, CMNZ.  
Hudson 1939, p. 459, as synonym.

***nigra*** Philpott, 1931, p. 31 (*Scythris*).  
Mount Maungatua DN, C.E. Clarke; HT ♂ designated  
by Philpott, AMNZ.  
Hudson 1939, p. 459, pl. lx fig. 3.


***niphozela*** Meyrick, 1931b, p. 369 (*Scythris*).  
Birdling's Flat MC, S. Lindsay; HT ♂ designated by  
Meyrick, CMNZ.  
Hudson 1939, p. 459, pl. lx fig. 14.

***triatma*** Meyrick, 1935, p. 304 (*Scythris*).  
Puhipuhi River KA, G.V. Hudson; LT ♂ here designated,  
labelled “Puhipuhi River New Zealand GVH  
11.11.33”, “Scythris triatma Meyr. 3/4 E. Meyrick det.  
in Meyrick Coll.”, BMNH.  
Hudson 1939, p. 459, pl. lx fig. 2.

Also 2 undescribed species (NZAC).

—◎—

**Family ELACHISTIDAE**  
 (in the sense of Traugott-Olsen & Nielsen  
 1977, pp. 9–26)




---

**Elachistidae**  
 (163) *Cosmiotes archaeonoma* (Meyrick)

---

● ***Cosmiotes*** Clemens, 1860, p. 8. Type species *Cosmiotes illectella* Clemens, 1860; North America. See Braun (1948, pp. 89–90).

***archaeonoma*** Meyrick, 1889b, p. 179, line 28  
 (*Elachista*)

Auckland [Domain] AK, E. Meyrick; LT ♂ here designated, labelled “Auckland New Zealand 12/1/80”, “*Elachista archaeonoma* Meyr. 7/18 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 319, pl. xxviii fig. 9 and 10, as *Elachista archaeonoma*.

Note. Placed in *Cosmiotes* by Dugdale (1971b, p. 79).

***exaula*** Meyrick, 1889b, pp. 178–179 (*Elachista*)

Mount Arthur NN, E. Meyrick; LT ♂ here designated, labelled “Mt Arthur New Zealand 4000 ft 15/1/85”, “*Elachista exaula* Meyr. 1/5 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 319, pl. xlvi fig. 3, pl. xxvii fig. 15, as *Elachista exaula*.

Note. Placed in *Cosmiotes* by Dugdale (1971b, p. 79).

***helonoma*** Meyrick, 1889b, p. 178 (*Elachista*)

[Port Hills], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 4/1/80”, “*Elachista helonoma* Meyr. 1/10 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 319, pl. xxviii fig. 14, as *Elachista helonoma*.

Note. Placed in *Cosmiotes* by Dugdale (1971b, p. 79).

***laqueorum*** Dugdale, 1971b, pp. 80–82, fig. 21–24  
 (*Cosmiotes*)

Sinkhole Flat, The Snares islands, P.M. Johns; HT ♂ designated by Dugdale, NZAC.

***ochroleuca*** Meyrick, 1923, p. 167 (*Elachista*)

Mount Aurum OL, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 320, pl. xl fig. 9, as *Elachista ochroleuca*.

Note. This species may be synonymous with *C. helonoma*. Placed in *Cosmiotes* by Dugdale (1971b, p. 79).

***ombrodoca*** Meyrick, 1889b, p. 179, line 6  
 (*Elachista*)

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ here designated, labelled “Christchurch New Zealand 22/3/82”, “*Elachista ombrodoca* Meyr. 1/12 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 319, pl. xxviii fig. 4.

Note. Examination of a long series suggests that *C. ombrodoca* (South Island), *C. archaeonoma* (North Island), *C. aphanta* (Turner) from Queensland, and *C. synethes* Meyrick from Sydney, as well as material from Tasmania, Lord Howe Island, and Norfolk Island, are regional populations of a single species. A specimen referable to this complex was seen from Rennell Island (Solomon Islands). Placed in *Cosmiotes* by Dugdale (1971b, p. 79).

***wattii*** Philpott, 1924a, p. 213 (*Elachista*)

Waimarino TO, M.N. Watt; HT ♂ designated by Philpott, abdomen missing, NMNZ.

Hudson 1928, p. 319, not figured, as synonym of *Elachista exaula*.

Note. Placed in *Cosmiotes* by Dugdale (1971b, p. 79).

Also 3 undescribed species (3 NZAC; 1 Watt Collection, NMNZ).

● ***Elachista*** Treitschke, 1833, p. 177, in the sense of Traugott-Olsen & Nielsen (1977, pp. 46–48). Type species *Elachista bifasciella* Treitschke, by subsequent designation (Meyrick 1915b, p. 210).

***Irenicodes*** Meyrick, 1919, p. 352. Type species *Irenicodes eurychora* Meyrick, by original monotypy. Synonymised by Traugott-Olsen & Nielsen (1977, p. 47).

***Euproteodes*** Viette, 1954, p. 19. Type species *Euproteodes galathea* Viette, by original designation. Synonymised by Traugott-Olsen & Nielsen (1977, p. 47).

Note. Examination of *E. holdgatei* Bradley from the Falkland Islands shows that the basal saccular arm articulates with the tip of the juxtal lobe (as it does in many Palearctic species), whereas in New Zealand species this arm articulates laterally with the juxtal lobe. Traugott-Olson & Nielsen (1977, pp. 17–18) recommend the terms *digitate process* for “sacculus lobes” and *juxtal lobes* for “anellar lobes” of Dugdale (1971b).

***eurychora*** Meyrick, 1919, p. 352 (*Irenicodes*)

Paekakariki WN, G.V. Hudson; HT ♂ unique, BM genitalia slide no. 15852 ♂, BMNH.

Hudson 1928, p. 259, pl. xlvi fig. 10, as *Irenicodes eurychora* in “Diplosarides”.

Note. Zimmerman (1971, p. 53) reassigned this species to Elachistidae.

***galathea*** is considered to consist of 3 subspecies, as follows:

► Elachistidae, *Elachista*

**galathea galathea** Viette, 1954, p. 21 (*Euproctodes*, as species)  
"Station L 409", Campbell Island, Lemche; HT ♂ designated by Viette, ZMKD.  
Dugdale 1971b, pp. 83–85, fig. 25–28.

**galathea antipodensis** Dugdale, 1971b, p. 85  
(*Irenicodes*)  
Antipodes Island, G. Kuschel; HT ♂ designated by Dugdale, NZAC.  
Dugdale 1971b, pp. 85–86, fig. 29–31.

1 undescribed subspecies, Auckland Islands (NZAC)

**hookeri** Dugdale, 1971b, pp. 85–86 (*Irenicodes*)  
Fairchild's Garden, Adams Island, Auckland Islands,  
K.A.J. Wise; HT ♂ designated by Dugdale, NZAC.  
Dugdale 1971b, pp. 85–87, fig. 32–36.

**napaea** Philpott, 1930c, pp. 438–439 (*Elachista*)  
Governor's Bush, Mount Cook MK, A. Philpott; HT ♂  
designated by Philpott, abdomen missing, wings  
mounted on card, CMNZ.  
Hudson 1939, pp. 458–459, pl. lx fig. 24.

**plagiaula** Meyrick, 1938, pp. 427–428 (*Thectophila*)  
new combination  
Freehold Range, Lake Ohau MK; HT ♂ designated by  
Meyrick, CMNZ.  
Hudson 1939, p. 441, pl. lxi fig. 28, as *Thectophila plagiula* in Cosmopterigidae.  
Note. Both type specimens (PLT ♂ in BMNH) lack characters of Cosmopterigidae; they are members of the  
*sagittifera / thallophora* species group in *Elachista*.

**pumila** Dugdale, 1971b, pp. 87–88 (*Irenicodes*)  
Bivouac Hill, Auckland Islands, J.L. Gressitt; HT ♂ designated by Dugdale, NZAC.  
Dugdale 1971b, pp. 87–89, fig. 37–39.

**sagittifera** Philpott, 1927d, p. 84 (*Elachista*)  
Arthur's Pass NC/WD, S. Lindsay; HT ♂ designated by  
Philpott, CMNZ.  
Hudson 1939, p. 458, pl. lx fig. 1.

**thallophora** Meyrick, 1889b, p. 178 (*Elachista*)  
[sandhills near] Kaiapoi MC, E. Meyrick; LT ♂ here designated, labelled "Kaiapoi New Zealand 31/3/82",  
"Elachista thallophora Meyr. 1/12 E. Meyrick det. in  
Meyrick Coll.", BMNH.  
Hudson 1928, p. 319, pl. xlvi fig. 12.

Also 2 undescribed species (NZAC).

• *Elachista* of authors

**gerasmia** Meyrick, 1889b, p. 177 (*Elachista*)  
[swampy forest near] Hamilton WO, E. Meyrick; LT ♂

here designated, labelled "Hamilton New Zealand 15/1/80", "Elachista gerasmia Meyr. 2/7 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pp. 319–320, pl. xxviii fig. 11.

Note. Dugdale (1971b, p. 79) lists points of difference between *E. gerasmia* and other elachistid groups in New Zealand. It has been reared from larvae mining *Juncus*; the naked pupa is held by a girdle (as in some Papilioidea).

**melanura** Meyrick, 1889b, p. 177 (*Elachista*)  
[heath-like scrub and swamp near] Hamilton WO, E.  
Meyrick; HT ♀ unique, BMNH.

Hudson 1928, p. 320, not figured.

Note. A ♀ from Sydney N.S.W. placed with the HT in BMNH differs markedly in colour pattern. *E. melanura* has not been collected since Meyrick's time in New Zealand.



Superfamily COSSOIDEA

Family COSSIDAE

● **Xyleutes** Hübner, 1820, p. 195. Type species *Xyleutes crassa* Drury, 1780, by subsequent designation (Kirby 1892, p. 874).

**boisduvali** Rothschild, 1896, p. 232 (*Xyleutes*)  
[Australia].  
New Zealand: adventive, non-establishing (Philpott 1927a, p. 708).



Superfamily SESIOIDEA

Family SESIIDAE

● **Synanthedon** Hübner, 1819, p. 129. Type species *Sphinx oestriformis* Röttemburg, as in Duckworth & Eichlin (1974, p. 25).


**tipuliformis** Clerck, 1759, pl. 9 fig. 1 (*Sphinx*)  
[Europe; adventive to Canada].  
New Zealand: adventive, widespread on *Ribes*; first reported by Fereday (1869, p. 146).

Hudson 1928, p. 250, pl. xxxiii fig. 6.  
Note. *Synanthedon tipuliformis* is conserved by ICZN opinion 1288 (1985a).

**salmachus** Linnaeus, 1758, p. 493 (*Sesia*). Name suppressed by ICZN opinion 1288 (1985a). [See Kristensen 1980, pp. 154–155.]



**Family CHOREUTIDAE**  
 (in the sense of Heppner & Duckworth  
 1981, p. 45)




---

**Choreutidae**  
 (164) *Asterivora combinatana* (Walker)

---

• ***Asterivora*** Dugdale, 1979, p. 461. Type species *Simaethis combinatana* Walker, by original designation (Dugdale 1979, p. 461).

***Asterophaga*** Horning & Greenwood, 1977, p. 295; nomen nudum.

***Simaethis*** not of Leach (1815, p. 135) but in the sense of Meyrick (1880b, p. 210).

***albifasciata*** Philpott, 1924a, p. 2113 (*Simaethis*)  
 Mt Arthur NN, 4,000 ft, A. Philpott; HT ♂ designated by Philpott, NZAC.  
 Hudson 1928, p. 309, pl. I fig. 6 and 7, as *Simaethis albifasciata*.

***analogia*** Meyrick, 1912c, p. 122 (*Simaethis*)  
 Mt Arthur NN, 4,000 ft, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 15.1.86", "*Simaethis analogia* Meyr. 3/9 E. Meyrick det. in Meyrick Coll.", BMNH.  
 Hudson 1928, p. 309, pl. xlvi fig. 1, as *Simaethis analogia*.

***antigrapha*** Meyrick, 1911b, p. 76 (*Simaethis*)  
 "Kaitoke and Karori [WN], in December and March (Hudson); three specimens" (Meyrick 1911, p. 76); LT and 2 PLTs not in BMNH; other BMNH material either wrong month (November) or wrong years (1910, 1913, 1914, 1916, 1922).  
 Hudson 1928, p. 309, pl. xxxiii fig. 28, as *Simaethis antigrapha*.

***barbigera*** Meyrick, 1915a, p. 203 (*Simaethis*)  
 Mt Cleugharn, Hunter Mountains FD, A. Philpott; HT ♀ unique, BMNH.  
 Hudson 1928, p. 310, pl. xl fig. 13, as *Simaethis barbigera*.

***chatuidea*** Clarke, 1926, p. 421 (*Simaethis*)  
 Anderson's Bay DN, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.  
 Not mentioned by Hudson.

***colpota*** Meyrick, 1911b, p. 67 (*Simaethis*)  
 West Plains, Invercargill SL, A. Philpott; HT ♀ unique, BMNH.  
 Hudson 1928, p. 308, pl. xl fig. 14, as *Simaethis colpota*.

***combinatana*** Walker, 1863c, p. 456 (*Simaethis*)  
 [Auckland AK], "New Zealand", D. Bolton; LT ♀ designated by Durrant and here published, BMNH.  
 Hudson 1928, p. 307, pl. xxxiii fig. 29, as *Simaethis combinatana*.

***abstinetella*** Walker, 1864b, p. 997 (*Simaethis*).  
 Synonymised by Meyrick (1883, p. 184).  
 [Auckland AK], "New Zealand", D. Bolton; HT ♀ unique, BMNH.  
 Hudson 1928, p. 307, as synonym.

***zomeuta*** Meyrick, 1912c, p. 121 (*Simaethis*).  
 Synonymised by Hudson (1923, p. 307).  
 Mt Arthur NN, E. Meyrick; HT ♀ unique, BMNH.  
 Hudson 1928, pp. 307–308, as synonym.

***exocha*** Meyrick, 1907c, pp. 120–121 (*Simaethis*)  
 Humboldt Range, Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, BMNH.  
 Hudson 1928, p. 307, pl. xxxiii fig. 31.

***fasciata*** Philpott, 1930b, p. 13 (*Simaethis*)  
 Arthur's Pass NC/WD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
 Hudson 1939, p. 457, pl. lx fig. 17, as *Simaethis fasciata*.

***inspoliata*** Philpott, 1930b, p. 12 (*Simaethis*)  
 Hunter Mountains FD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
 Hudson 1939, p. 457, pl. lx fig. 18, as *Simaethis inspoliata*.

***iochondra*** Meyrick, 1911b, p. 77 (*Simaethis*)  
 Mount Holdsworth, Tararua Range WN, 3,000 ft, G.V. Hudson; LT ♂ here designated, labelled "Mt Holdsworth New Zealand GVH 2.10", "*Simaethis iochondra* Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", BMNH.  
 Hudson 1928, p. 308, pl. xxxiii fig. 1 and 2, as *Simaethis iochondra*.

***marmarea*** Meyrick, 1888e, p. 85 (*Simaethis*)  
 Lake Wakatipu OL, [2,200 ft], E. Meyrick; HT ♂ unique, BMNH.  
 Hudson 1928, p. 309, not figured, as *Simaethis marmarea*.

***microlitha*** Meyrick, 1888e, p. 84 (*Simaethis*), in the restricted sense of (Meyrick 1912c, p. 122)  
 Arthur's Pass NC/WD, [3,000–5,000 ft], E. Meyrick; LT ♀ here designated, labelled "Arthur's Pass New Zealand 25.1.83", "*Simaethis microlitha* Meyr. 2/19 E. Meyrick det. in Meyrick Coll.", BMNH.  
 Hudson 1928, p. 309, pl. xl fig. 7, as *Simaethis microlitha*.  
 Note. Meyrick (1912c) restricted his concept of this species to the 2 Arthur's Pass specimens; his 1888c series included specimens from Mt Arthur NN and Castle Hill MC. I could not find the specimens from Castle Hill.

➤ Choreutidae, *Asterivora*

***ministra*** Meyrick, 1912c, p. 121 (*Simaethis*)  
Mount Holdsworth WN, G.V. Hudson; HT ♂ unique,  
BMNH.  
Hudson 1928, p. 308, pl. xxxiii fig. 30, as *Simaethis  
ministra*.

***nivescens*** Philpott, 1926a, p. 397 (*Simaethis*)  
[Gordon's Pyramid], Mt Arthur NN, A. Philpott; HT ♂  
designated by Philpott, NZAC.  
Hudson 1928, p. 310, pl. lii fig. 4, as *Simaethis nivescens*.

***oleariae*** Dugdale, 1979, p. 462 (*Asterivora*)  
The Snares islands, D.S. Horning; HT ♂ designated by  
Dugdale, NZAC.

***symbolaea*** Meyrick, 1888e, p. 85 (*Simaethis*)  
Arthur's Pass NC/WD, [3,000–5,000 ft] E. Meyrick; LT  
♂ here designated, labelled "Arthur's Pass New  
Zealand 25.1.83", "Simaethis symbolaea Meyr. 7/13  
E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 308, pl. xl fig. 21, as *Simaethis symbolaea*.

***tillyardi*** Philpott, 1924c, p. 666 (*Simaethis*)  
Mt Cook MK, A. Philpott; HT ♂ designated by Philpott,  
NZAC.  
Hudson 1928, p. 310, pl. li fig. 22, as *Simaethis tillyardi*.

***tristis*** Philpott, 1930b, p. 12 (*Simaethis*)  
[Tongariro] Mt Ruapehu TO, A. Philpott; HT ♂ designa-  
ted by Philpott, AMNZ.  
Hudson 1939, p. 457, pl. ix fig. 16, as *Simaethis tristis*.

***urbana*** Clarke, 1926, pp. 420–421 (*Simaethis*)  
Arthur's Pass NC/WD, C.E. Clarke; HT ♂ designated by  
Clarke, AMNZ.  
Hudson 1928, pp. 309–310, pl. lii fig. 16, as *Simaethis  
urbana*.

Also 3 undescribed species (NZAC).

● ***Tebenna*** Billberg, 1820, p. 90. Type species *Tinea  
bjerkandrella* Thunberg, 1784, p. 79, by subsequent  
designation (Bradley 1966, p. 220).

***micalis*** Mann, 1857, p. 181 (*Choreutis*)  
Fiume [Rijeka], Yugoslavia, Wacke; LT ♂ selected by A.  
Diakonoff, ZILR.  
Hudson 1928, p. 310, pl. xxxiii fig. 27, as *Choreutis  
bjerkandrella*.


***bradleyi*** Clarke, 1971, pp. 166–167 (*Tebenna*).  
Synonymised by Diakonoff (1986, p. 153).  
Haumoana HB, T.H. Davies; HT ♂ designated by Clarke,  
NZAC (since 1973).

***bjerkandrella*** in the sense of Meyrick (1915b, p.  
225; *Choreutis*).

—◎—

Superfamily TORTRICOIDEA

Family TORTRICIDAE



Tortricidae

(165) *Gelophaula trisulca* (Meyrick)

Subfamily CHLIDANOTINAE

(in the sense of Horak 1984, pp. 8 and 10)

Tribe Polyorthini

(in the sense of Diakonoff 1974)

● ***Lopharcha*** Diakonoff, 1941, p. 424. Type species  
*Lopharcha quinquestriata* Diakonoff, by original  
designation.

***Laciniella*** Dugdale, 1966, p. 768. Type species  
*Laciniella insolita* Dugdale, 1966, by original  
designation. New synonymy.

***insolita*** Dugdale, 1966, p. 768 (*Laciniella*) new  
combination

Rotoehu BP, R. Leggatt; HT ♂ designated by Dugdale,  
NZAC.

Note. The species *insolita* has genital characters inter-  
mediate between species groups 1 and 2 of Diakonoff  
(1974, pp. 23 and 50).

—◎—

Subfamily OLETHREUTINAE

(in the sense of Horak 1984, p. 10)

● ***Acroclita*** of authors, but not of Lederer (1859)

***discariana*** Philpott, 1930a, p. 248 (*Acroclita*)  
Porter River MC, S. Lindsay; HT ♂ designated by Phil-  
pott, CMNZ.  
Hudson 1939, p. 436, pl. lvii fig. 11 and 12.

● ***Argyroploce*** of authors, but not of Hübner (1825)

***chlorosaris*** Meyrick, 1914a, p. 106 (*Argyroploce*)  
Day's Bay WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 249, pl. lvii fig. 26.

● **Bactra** Stephens, 1834, in the sense of Diakonoff (1964). Type species *Tortrix lancealana* Hübner, 1796, by original designation.

**Noteraula** Meyrick, 1892, p. 217. Type species *Bactra straminea* Meyrick, 1892, not of Butler (1881b, p. 393); synonymised by Diakonoff (1956, p. 18).

**noteraula** Walsingham, 1907, p. 689 (*Bactra*), new name for *Noteraula straminea* Meyrick, 1885 (but not Butler, 1881b, p. 393).

**straminea** [not of Butler, 1881b] Meyrick, 1885g, p. 142 (*Noteraula*).

Wanganui WI [in swamp], E. Meyrick; LT ♂ designated by Diakonoff (1964, p. 21), BMNH.

Hudson 1928, p. 248, pl. xxvi fig. 9, as *Bactra noteraula*.

**xystrota** Meyrick, 1911b, pp. 62–63 (*Bactra*). Synonymised by Diakonoff (1964, p. 21).

Invercargill SL, A. Philpott; LT ♂ designated by Diakonoff (1964, p. 21), BMNH.

Hudson 1928, p. 249, pl. xxvi fig. 18, as species.

**optanias** Meyrick, 1911c, p. 89 (*Bactra*) [published 1 July 1911]

Hamilton WO, E. Meyrick; LT ♀ here designated, labelled “Hamilton New Zealand 13/1/80”, “*Bactra optanias* Meyr. 3/3 E. Meyrick det. in Meyrick Coll.”, BMNH. Hudson 1928, p. 248, pl. xlvi fig. 30.

Note. This is *Aphelia lanceolana* in the sense of Meyrick (1881b, pp. 651–652; 1883b, p. 59), not of Hübner. Diakonoff (1964, p. 40) chose as lectotype a ♂ from Meyrick's Australian syntype series, described by Meyrick (1911d, p. 253 [published 20 November 1911]). Meyrick described *optanias* from New Zealand and Australia separately, on 1 July and 20 November 1911 respectively, each time stating “*B. optanias* n. sp.”. Each description is based solely on material from the country in which he published (e.g., 1 July 1911 “the above description is taken solely from my New Zealand examples”). Diakonoff's lectotype designation is valid for the Australian species, but invalid for the New Zealand species. Should the two type populations be shown to be conspecific (e.g., reproductively effective), Diakonoff's actions stand, under the principle of the action of the first reviser.

● **Crocidosema** Zeller, 1847, p. 721. Type species *Crocidosema plebejana* Zeller, by original monotypy.

**plebejana** Zeller, 1847, p. 721

Cosmopolitan.

New Zealand: widespread on herbaceous Malvaceae. Hudson 1928, p. 248, pl. xlix fig. 10, as *C. plebeiana*. First reported by Philpott (1923, p. 151).

● **Cryptaspasma** Walsingham, 1900, p. 462, in the sense of Diakonoff (1959). Type species ?*Penthina*

*lugubris* Felder & Rogenhofer, by original designation; Brazil.

**querula** Meyrick, 1912c, p. 125 (*Eucosma*) Wellington WN, G.V. Hudson; LT ♂ designated by Diakonoff (1959, p. 28), BMNH. Hudson 1928, pp. 247–248, pl. xxv fig. 36 and 37, as *Eucosma querula*.

● **Cydia** Hübner, [1825], p. 375. Type species *Phalaena Tortrix pomonella* Linnaeus, by subsequent designation (Walsingham 1897, pp. 125 and 130).

**pomonella** Linnaeus, 1759, p. 538 (*Phalaena Tortrix*) European.

New Zealand: man-adventive. First recorded by Meyrick at Wellington “in town”, 31 December 1879 (Meyrick's Diary of Captures).

Hudson 1928, p. 249, pl. xxv fig. 38, as *Laspeyresia pomonella*.

Note. The binomen for the codlin moth as given here was ‘stabilised’ by Brown (1979, p. 565). Kuznetsov & Kerzhner (1984, pp. 110–113) have queried the interpretation of the evidence, and proposed conservation of *Laspeyresia* Hübner, but separate dissenting views are given by Hodges (1985), Miller (1985), and Bradley et al. (1985).

● **Grapholita** Treitschke, 1829, p. 232. Type species *Pyralis dorsana* Fabricius, as reported by Fernald (1908, p. 57).

**molesta** Busck in Quaintance & Wood, 1916, p. 373 (*Laspeyresia*)

New Zealand: man-adventive since 1973: north of 40°S (1984).

● **Hendecasticha** Meyrick, 1881b, p. 692. Type species *Hendecasticha aethaliana* Meyrick, by original monotypy.

**aethaliana** Meyrick, 1881b, p. 692 (*Hendecasticha*) Hamilton WO, E. Meyrick; LT ♂ here designated, labelled “Hamilton New Zealand 15/1/80”, “*Hendecasticha aethaliana* Meyr. 4/5 E. Meyrick det. in Meyrick Coll.”, BMNH. Hudson 1928, p. 245, pl. xlvi fig. 31.

● **Parienia** Berg, 1899, p. 78, new name for *Exoria* Meyrick, §1882c, preoccupied by *Exoria* Hübner, 1825 (Pyralidae).

**Exoria** Meyrick, §1882c, p. 278; 1883b, pp. 58 (key) and 65. Type species *Exoria mochlophorana* Meyrick, §1882c, by original monotypy.

> Tortricidae, *Parienia*

**mochlophorana** Meyrick, §1882c, p. 278; 1883b, p.

65 (*Exoria*)

South Rakaia MC, W.H. Gaze; HT ♀ unique, BMNH. Hudson 1928, p. 247, not figured.

**aphrias** Meyrick, 1901, p. 578 (*Epiblemma*). Synonymised by Philpott (1931, p. 30).

[The Elbow (near Lumsden)], Invercargill SL, E. Meyrick; LT ♂ here designated, labelled "Invercargill New Zealand 14/12/82", "Epiblemma aphrias Meyrick teste K.R. Tuck 1980", "Eucosma mochlophorana Meyr. 2/5 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 247, not figured.

**trimaculata** Philpott, 1915, pp. 198–199 (*Eurythecta*). Synonymised by Philpott (1931, p. 30).

Queenstown OL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 224, pl. xxvii fig. 6 and 7.

● **Protithona** Meyrick, §1882c, p. 278; 1883b, pp. 58 (key) and 62. Type species *Protithona fugitivana* Meyrick, by original monotypy.

**Raumatia** Philpott, 1928e, p. 473; 1928g, pp. 487–488 (nomen nudum). Type species *Eurythecta potamias* Meyrick, by original designation.

**fugitivana** Meyrick, §1882c, p. 278; 1883b, p. 62 (*Protithona*)

Lake Coleridge MC, R.W. Fereday; HT ♂ unique, BMNH. Hudson 1928, p. 247, not figured (but p. 224, pl. xlvi fig. 8, as *Eurythecta varia* Philpott); 1939, p. 437.

**varia** Philpott, 1916, pp. 421–422 (*Eurythecta*). Synonymised by Philpott (1931, p. 30).

Kaikoura Range KA, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 224, pl. xlvi fig. 8.

**potamias** Meyrick, 1909a, p. 11 (*Eurythecta*)

[Riverton], Invercargill SL, A. Philpott; LT ♂ here designated, labelled "Invercargill New Zealand AP 3.07", "Eucosma potamias Meyr. 4/4 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, pp. 224–225, pl. xxvi fig. 11 and 12; type locality given as sandhills, Riverton.

Note. This species was probably collected on salt-tolerant short turf among sandhills (B. Patrick, pers. comm.).

● **Strepsicrates** Meyrick, 1888, p. 73, new name for *Strepsicerous* Meyrick, preoccupied by *Strepsicerous* Rafinesque, 1815 (Mammalia).

**Strepsicerous** Meyrick, 1881b, p. 678. Type species *Sciaphila ejectana* Walker, 1863, by subsequent designation (Meyrick 1911, p. 87).

**charopa** Meyrick, 1888d, p. 73 (*Strepsicrates*) [Waitakere Range], Auckland AK, E. Meyrick; LT ♂ here

designated, labelled "Auckland New Zealand 17/12/85", "Strepsicrates charopa Meyr. 1/4 E. Meyrick det. in Meyrick Coll.", "Lectotype", BMNH.

Hudson 1928, p. 245, not figured; 1939, p. 436, pl. lxi fig. 11, as *Spilonota charopa*.

**dolopaea** Meyrick, 1905, pp. 232–233 (*Strepsicrates*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 245, pl. xxvii fig. 25, pl. xlvi fig. 5, as *Spilonota dolopaea*.

**ejectana** Walker, 1863c, p. 350 (*Sciaphila*)

Sydney N.S.W., Lambert; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 246, pl. xxvii fig. 1 and 2, as *Spilonota ejectana*.

**servilisana** Walker, 1863c, p. 356 (*Sciaphila*)

Synonymised by Meyrick (1881b, p. 681).

Auckland AK, D. Bolton; HT ♂ unique, BMNH.

Hudson 1928, p. 246, as synonym.

**infimana** Walker, 1863c, p. 357 [line 8] (*Sciaphila*). New synonymy.

Auckland AK, D. Bolton; LT ♂ here designated, labelled "Sciaphila infimana" (printed strip), "Sciaphila infimana Wkr Cat. Lep. Het. BM 28 p.357 (1863) Type ♂", "New Zeal. // 54.4" (circular label), "Type" (circular label), BMNH.

Note. Redescribed by Walker (1864b, p. 986) as *infimana*. Meyrick (1881b, p. 704) wrote of *infimana*: "Type very worn, unidentifiable. Meyr."

**saxana** Walker, 1863c, p. 357 [line 23] (*Sciaphila*). Synonymised by Meyrick (1881b, p. 681).

Auckland AK, D. Bolton; HT ♀ unique, BMNH.

Hudson 1928, p. 246, as synonym.

**ligniferana** Walker, 1863, p. 363 (*Conchylis*)

Synonymised by Meyrick (1881b, p. 681).

Sydney N.S.W., Diggles; HT ♀ unique, BMNH.

Hudson 1928, p. 246, as synonym.

**emplasta** Meyrick, 1901, p. 571 (*Strepsicrates*)

[West Plains] Invercargill SL, [A. Philpott] G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 246; 1939, p. 436, pl. lxi fig. 10, as *Spilonota emplasta*.

**infensa** Meyrick, 1911d, p. 228 (*Spilonota*)

Brisbane Qld, Turner; type material BMNH.

New Zealand: recorded ND–SL, on *Eucalyptus* species. Not mentioned by Hudson.

**macropetana** Meyrick, 1881b, pp. 683–684 (*Strepsicerous*)

Blackheath N.S.W., E. Meyrick; type series BMNH.

New Zealand: recorded AK–SL, on *Eucalyptus* species. Hudson 1928, p. 247, pl. xlvi fig. 8, as *Spilonota macropetana*.

Note. Meyrick (1923, p. 164) gives the first record, received by Hudson (1921) from *Eucalyptus* foliage at Auckland.

**parthenia** Meyrick, 1888d, pp. 73–74 (*Strepsicrates*)  
[Waitakere Range], Auckland AK, E. Meyrick; LT ♂ here  
designated, labelled “Auckland New Zealand  
22/12/85”, “*Strepsicrates parthenia* Meyr. 2/3 E.  
Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, pp. 245–246, pl. xxvii fig. 26, as *Spilonota*  
*parthenia*.

**sideritis** Meyrick, 1905, p. 232 (*Noteraula*) new  
combination

“New Zealand”, ?G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 249, not figured, as *Bactra sideritis*.

Note. Although the locality label reads “New Zealand  
G.V.H. /00”, Hudson (1928, p. 249) states “... is  
believed to have been taken at Wellington ... I am  
unacquainted with this insect”. Diakonoff (1964, p.  
80) notes that *sideritis* is an “apocryphal *Bactra*”; and  
as the ♀ has 2 hooked signa, it “is not a *Bactra* at  
all”.

**chaophila** Meyrick, 1909a, p. 10 (*Strepsicrates*).  
New synonymy.

Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 246, pl. xxvi fig. 35 and 36, as *Spilonota*  
*chaophila*.

**zopherana** Meyrick, 1881b, pp. 688–689  
(*Strepsicerous*)

Sydney N.S.W., E. Meyrick; LT ♂ selected by I.F.B.  
Common and here designated, labelled “*Strepsicerous*  
*zopherana* Meyr. Lectotype IFBC vii. 1954”, “Sydney  
N.S. Wales 19/1/79”, “*Strepsicerous zopherana* Meyr.  
7/7 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 246, pl. xxvii fig. 29; also pl. xxvii fig.  
27 (Hudson 1939, p. 436), as *Spilonota zopherana*.  
Note. ?Naturally common to Australia and New Zealand.

- Also 1 undescribed genus and species (NZAC).



### Subfamily TORTRICINAE

(in the sense of Horak 1984, pp. 8 and 9)

Most New Zealand Tortricinae were placed by  
Meyrick in Palearctic genera. These have since been  
redefined by modern workers on genital characters  
(largely ignored for this group by Meyrick), and  
therefore restricted. Those New Zealand species that  
lack generic placement are here listed under the  
genus names used by Hudson (1928, 1939), and are  
placed at the end of each tribal section; the reader  
is thereby warned that the classification is in need  
of revision. Placement of any two species under  
such genus names is not, therefore, an intentional  
indication of relationship. Horak (1984, pp. 3–64)  
gives a detailed account of structures used in the  
classification of this group.

### Tribe Archipini in the strict sense (Obraztsov 1942, p. 147 (in part))

Note. Only those groups lacking a costa on the ♂  
valva, and with the valva characteristically trian-  
gular and ‘plicate’, are here included in this tribe.

• **Epiphyas** Turner, 1927, p. 126. Type species *E.*  
*eucyrtta* Turner, by original designation.

**Austrotortrix** Bradley, 1956a, p. 101. Type spe-  
cies *Teras postvittana* Walker, by original desig-  
nation. Synonymised by Common (1961, p. 178).

**postvittana** Walker, 1863c, p. 297 (*Teras*)  
Sydney N.S.W., Lambert; HT ♀ unique, BMNH.

New Zealand: recorded ND-SL.  
Hudson 1928, p. 228, pl. xxiv fig. 36, as *Tortrix postvittana*.  
Note. Common (1961, p. 180) gives synonymy. A spec-  
imen in CMNZ is labelled as reared from Tasmanian  
apples in 1887; and from about 1891 the species  
became widespread (Philpott 1925, p. 366). Meyrick  
did not obtain specimens in New Zealand over the  
period December 1879 to February 1886.

• **Ericodesma** Dugdale, 1971b, p. 158. Type species  
*Tortrix melanosperma* Meyrick, 1916, by original  
designation.

**aerodana** Meyrick, 1881a, p. 520 (*Tortrix*)

Hamilton WO, E. Meyrick; LT ♂ here designated, labelled  
“Hamilton New Zealand 17.1.80”, “*Tortrix aerodana*  
Meyr. 1/6 E. Meyrick det. in Meyrick Coll.”, “Lec-  
totype *Tortrix aerodana* Meyr. J.D. Bradley 1962”,  
“BM genitalia slide no. 88960 ♂”, BMNH.

Hudson 1928, p. 227, pl. xlvi fig. 17; also p. 227, pl. xxvi  
fig. 8; as *Tortrix indigestana* [not of Meyrick (1881a,  
p. 520)].

Note. Dugdale’s (1971, p. 158) mention of *E. aerodana*  
on *Styphelia* (as *Cyathodes*) refers to an undescribed  
species. The host plants of *E. aerodana* are mat-form-  
ing *Pimelea* species.

**argentosa** Philpott, 1924a, pp. 209–210 (*Tortrix*)  
Dun Mountain NN, A. Philpott; HT ♂ designated by  
Philpott, NZAC.

Hudson 1928, p. 227, not figured, as *Tortrix argentosa*.

**subdola** Philpott, 1924a, p. 212 (*Tortrix*). New  
synonymy.

Mt Ruapehu TO, C.C. Fenwick; HT ♂ designated by  
Philpott, NMNZ.

Hudson 1928, p. 227, not figured, as *Tortrix subdola*.

**cuneata** Clarke, 1926, p. 419 (*Tortrix*)

Hope Arm, Lake Manapouri FD, C.E. Clarke; HT ♂ des-  
ignated by Clarke, AMNZ.  
Not mentioned by Hudson.

**melanosperma** Meyrick, 1916b, p. 414 (*Tortrix*)

Arthur’s Pass NC/WD, G.V. Hudson; HT ♂ unique,  
abdomen missing, BMNH.

» Tortricidae, *Ericodesma melanosperma*

Hudson 1928, p. 227, pl. xlvii fig. 4, as *Tortrix melanosperma*.

**scruposa** Philpott, 1924a, p. 212 (*Tortrix*).  
Mt Ruapehu TO, C.C. Fenwick; HT ♂ designated by  
Philpott, NMNZ.

Hudson 1928, p. 232, pl. I fig. 4, as *Tortrix scruposa*.

**maculosa** Philpott, 1927d, p. 84 (*Tortrix*).  
Synonymised by Hudson (1939, p. 433).  
Quartz Range NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 433, as synonym.

Also 2 undescribed species (NZAC).

● **Eurythecta** Meyrick, 1883b, pp. 36 (key) and 56.  
Type species *Zelotherses robusta* Butler, 1877, by original monotypy.

**robusta** Butler, 1877, p. 403 (?*Zelotherses*)  
"Canterbury Plains" MC, J.D. Enys; LT ♀ here designated, labelled "N. Zeal. 77.34", "Zelotherses robusta Btlr. P.Z.S. Lond. p. 403 + fig. 43'17 (1877)", "BM genitalia slide 10690", BMNH.  
Hudson 1928, p. 224, pl. xxvi fig. 23, as *Eurythecta robusta*.

**negligens** Butler, 1877, p. 404 (*Steganoptycha*).  
Synonymised by Meyrick (1883b, p. 56).  
"Canterbury Plains" MC, J.D. Enys; HT ♂ unique, BMNH.  
Hudson 1928, p. 224, as synonym.

**zelaea** Meyrick, 1905, pp. 233–234 (*Eurythecta*)  
[Ida Valley CO], Dunedin DN, J.H. Lewis; LT ♂ here designated, labelled "Dunedin New Zealand J.H.L. '03", "Eurythecta zelaea Meyr. 1/2 E. Meyrick det. in Meyrick Coll.", "BM genitalia slide no. 10689", BMNH.  
Hudson 1928, p. 224, pl. xxvi fig. 10.

Also 1 undescribed species (B. Patrick Collection, Dunedin).

● **Merophyas** Common, 1964, p. 298. Type species *Conchylis divulsana* Walker, 1863, p. 364; Australia.

**leucaniana** Walker, 1863c, p. 370 (*Conchylis*) new combination

Auckland AK, D. Bolton; LT ♂ selected by H. Durrant, July 1919, and here designated, labelled "Auckland New Zealand, Bolton 1854:4", "Conchylis leucaniana Wlkr ♂ Cat. Lep. Het. BM 28 p.370 sp 132 (1863) a) New Zealand, Bolton 1854:4 Type ♂ 1/5 descr.", "BM genitalia slide 8907 ♂", BMNH.

Hudson 1928, p. 226, pl. xxvi fig. 29, as *Tortrix leucaniana*.

**intactella** Walker, 1864a, p. 652 (*Gelechia*). Synonymised by Meyrick (1881a, p. 517).

Auckland AK, D. Bolton; LT ♂ selected by H. Durrant and here designated, labelled "Gelechia intactella Wlkr ♂ Cat. Lep. Het. BM 29 p.652 (1864) a) New Zealand, Bolton 54.4 Type ♂", "BM genitalia slide 12660", BMNH. Hudson 1928, p. 226, as synonym.

**pauculana** Walker, 1866b, pp. 1781–1782 (*Teras*).  
Synonymised by Meyrick (1881a, p. 517).

Auckland AK, A. Sinclair; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 226, as synonym.

Note. The HT is glued to card.

**philopoana** Meyrick, 1881a, p. 515 (*Tortrix*). New synonymy.

Hamilton WO, E. Meyrick; LT ♂ selected by J.D. Bradley and here designated, labelled "Hamilton New Zealand 15.1.80", "Tortrix philopoana Meyr. 7/7 E. Meyrick det. in Meyrick Coll.", "BM genitalia slide 2050", BMNH. Hudson 1928, p. 226, pl. xlv fig. 26, as *Tortrix philopoana*, as species.

**paraloxa** Meyrick, 1907c, p. 116 (*Eurythecta*) new combination

[Riverton] SL, A. Philpott; LT ♂ selected by J.D. Bradley and here designated, labelled "Invercargill New Zealand AP '06", "Eurythecta paraloxa Meyr. 1/5 Meyrick det. in Meyrick Coll.", "BM genitalia slide 8606 ♂", BMNH.

Hudson 1928, p. 225 (as from Riverton), pl. xxvi fig. 13, as *Eurythecta paraloxa*. Philpott's illustrations of the ♂ genitalia (1928d, p. 461, fig. 41; p. 462, fig. 47) clearly indicate that, on these structures, *paraloxa* is best placed in *Merophyas*.

Note. This species was probably collected on estuarine swards at Riverton.

Also 4 undescribed species (NZAC).

● One undescribed genus and species, Chatham Islands (NZAC).

### Tribe Archipini

(in the broad sense, e.g., of Horak 1984)

Note. The tribe Epitymbiini Common, 1958b, p. 291, might well apply to some New Zealand groups, but in the absence of a 'definitive' classification they are left in Archipini in the broad sense. All species here have oblong valvae with a strongly sclerotised costa; in most the uncus is short, usually curved or deflexed, and often expanded apically.

● **Ascerodes** Meyrick, 1905, p. 234. Type species *A. prochlora* Meyrick, 1905, by original monotypy.

**prochlora** Meyrick, 1905, p. 234 (*Ascerodes*).  
Humboldt Mountains OL, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, pp. 225–226, pl. xxiv fig. 38.

**tritochlora** Meyrick, 1912c, pp. 120–121 (*Har-mologa*). New synonymy.

[alpine areas by Lake Harris, c. 4,000 ft] OL, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, pp. 235–236, pl. xxvi fig. 3, as *Gelophaula tritochlora*.

#### • *Capua* of authors

**semiferana** Walker, 1863c, p. 306 (*Teras*)

Auckland AK, D. Bolton; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 223, pl. xxvi fig. 7.

**detrifiana** Walker, 1863c, p. 356 (*Sciaphila*). Synonymised by Meyrick (1881a, p. 453; 1883b, p. 37).

Auckland AK, D. Bolton; LT ♂ selected by J.D. Bradley and here designated, labelled “*Sciaphila detritiana* Wkr Cat. Lep. Het. BM 28, p.356 (1863) Type ♀”, “New Zeal. 54.4”, “BM genitalia slide ♂ 8559”, BMNH.

Hudson 1928, p. 223, as synonym.

**admotella** Walker, 1863c, p. 485 (*Tinea*). Synonymised by Meyrick (1882, p. 453; 1883, p. 37).

Auckland AK, D. Bolton; LT ♀ selected by J.D. Bradley and here designated, labelled “*admotella* Wkr”, “*Tinea admotella* Wkr Cat. Lep. Het. BM 28 p.485 (1863) Type ♂”, “New Zeal. 54.4”, “BM genitalia slide no. 8532 ♀”, BMNH.

Hudson 1928, p. 228, as synonym.

Note. One PLT is *Strepsicrates zopherana* Tortricidae); the other is *Trachypepla euryleucota* Meyrick (Oecophoridae).

**abnegatana** Walker, 1864b, p. 991 (*Grapholitha*). Synonymised by Meyrick (1881a, p. 453; 1883, p. 37).

Auckland AK, D. Bolton; HT ♀ unique, BM genitalia slide no. 8537 ♀, BMNH.

Hudson 1928, p. 223, as synonym.

**polias** Meyrick, 1913a, p. 26 (*Capua*). Synonymised by Hudson (1928, p. 233).

Wellington WN, G.V. Hudson; HT ♂ unique, BM genitalia slide no. 8564 ♂, BMNH.

Hudson 1928, pl. xxvi fig. 5, as synonym.

Note. *Tortrix constrictana* Walker, 1866b, p. 1785, synonymised with *C. semiferana* by Meyrick (1911c, p. 82; 1913e, p. 13), is here removed from synonymy. The type ♂ of *T. constrictana* has the entire head and palpi dark-scaled and the forewing 3× longer than broad (New Zealand ♂ specimens have the face white- or grey-scaled, and the palpi are pallid on the inner surface). The type specimen of *T. constrictana* is listed as from Australia, collected by E. Damel.

Also 2 undescribed species (NZAC).

• **Catamacta** Meyrick, 1911c, p. 81. Type species *Pandemis gavisana* Walker, 1863, by original designation.

**alopecana** Meyrick, §1885b, p. 348; 1885g, pp. 147–148 (*Cacoecia*) new combination

Bealey Valley NC, E. Meyrick; LT ♂ selected by J.D.

Bradley and here designated, labelled “Bealey River New Zealand 21.1.83”, “*Capua alopecana* Meyr. 1/8 E. Meyrick det. in Meyrick Coll.”, “BM genitalia slide no. 8510 ♂”, BMNH.

Hudson 1928, p. 230, pl. xlvi fig. 11, as *Tortrix alopecana*; 1939, p. 43, pl. lxi fig. 30, larva, as *Capua alopecana*.

Note. On wing venation, wing position at rest, labial palpi, and ♂ and ♀ genitalia this species is included in *Catamacta* (see Dugdale 1971b, p. 164, for characters distinguishing *Catamacta* from *Pyrgotis*).

**gavisana** Walker, 1863c, p. 312 (*Pandemis*)

Auckland AK, D. Bolton; HT ♀ unique, BM genitalia slide no. 280, BMNH.

Hudson 1928, pp. 220–221, pl. xxii fig. 15–18, 34, and 35, as *Catamacta gavisana*.

**innotatana** Walker, 1863c, p. 333 (*Tortrix?*). Synonymised by Meyrick (1911c, p. 81).

[Nelson NN], T.R. Oxley; HT lost, BMNH.

Hudson 1928, p. 220, as synonym.

**marginana** Walker, 1863c, p. 371 (*Conchylis*). Synonymised by Meyrick (1881a, p. 442).

Auckland AK, D. Bolton; HT ♀ unique, BM genitalia slide no. 7808 ♀, BMNH.

**porphyreana** Meyrick, 1881a, pp. 443–444 (*Pyrgotis*). Synonymised by Meyrick (1883b, p. 40), as synonym of *Teras conditana* in the sense of Meyrick, but not of Walker (1863).

Wellington WN, E. Meyrick; HT ♂ unique, BMNH.

Hudson 1928, p. 220, as synonym.

**conditana** not of Walker, but in the sense of Meyrick (1881a, p. 443, *Pyrgotis*; 1883b, p. 40, *Adoxophyes*). Synonymised by Meyrick (1911c, p. 81).

**aoristana** Meyrick, 1881a, p. 466 (*Capua*). Synonymised by Meyrick (1883b, p. 40), as synonym of *Adoxophyes conditana* in the sense of Meyrick, but not Walker.

Auckland AK, Wellington WN, E. Meyrick; ST series not found, BMNH.

Hudson 1928, p. 220, as synonym.

Note. I could not find in BMNH collections the ♂ recorded in Meyrick's Diary of Captures as “*Cap. aoristana*” collected at Wellington WN on 9 January 1880, nor the ♂ similarly recorded as collected at Auckland AK on 12 January 1880.

**lotinana** Meyrick, 1882c, p. 277; 1883b, p. 40 (*Adoxophyes*)

Christchurch MC, R.W. Fereday; LT ♂ selected by J.D. Bradley and here designated, labelled “Christchurch New Zealand RWF/82”, “*Catamacta lotinana* Meyr. 2/2 E. Meyrick det. in Meyrick Coll.”, “abdomen missing”, BMNH.

Hudson 1928, p. 220, pl. xlvi fig. 15.

**rureana** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 47 (*Rhacodia*)

» Tortricidae, *Catamacta rureana*

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.  
Hudson 1928, p. 220, pl. xxvii fig. 22 and 23.

**camelina** Meyrick, 1891, pp. 97–98 (*Adoxophyes*). Synonymised by Meyrick (1911, p. 81).  
Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1928, p. 220, as synonym.

Also 1 new species (NZAC).

• ***Cnephasia*** of authors, e.g., Meyrick (1911c, p. 86; 1912e, p. 43; 1913e, p. 44), but not Obraztsov (1955, p. 147)

**holophrna** Meyrick, 1911b, p. 74 (*Cnephasia*)  
Mount Enys MC, G.V. Hudson; HT ♂ unique, abdomen, hindwings, and right forewing missing, BMNH.  
Hudson 1928, p. 243, pl. xxvii fig. 4.  
Note. Hudson gives the type locality as Mt Olympus; both Enys and Olympus are in the Craigieburn Range. See *melanophaea*, below.

**incessana** Walker, 1863c, p. 304 (*Teras*)  
[Nelson NN, T.R. Oxley]; LT ♂ selected by K.R. Tuck and here designated, labelled “Teras incessana Wlkr Cat. Lep. Het. BM 28 p.309 (1863) Type ♂”, “Auckland N. Zeal. // 60.73”, abdomen in gelatin capsule, BMNH.  
Hudson 1928, p. 242, pl. xxv fig. 9.

**jactatana** Walker, 1863c, p. 317 (*Batodes*)  
[Nelson NN] “Auckland”, T.R. Oxley; LT ♀ selected by K.R. Tuck and here designated, labelled “Batodes jactatana Wlkr Cat. Lep. Het. 28 p.317 (1863) Type ♂”, “Auckland N. Zeal. // 60.73”, [Walker's specimen a], BMNH.  
Hudson 1928, p. 242, pl. xxv fig. 17 and 18.

**flexivittana** Walker, 1863c, p. 353 (*Sciaphila*).  
Synonymised by Meyrick (1881a, p. 488; 1883b, p. 54).  
Auckland AK, D. Bolton; HT ♀ unique, BMNH.  
Hudson 1928, p. 242, as synonym.

? **privatana** Walker, 1863c, p. 382 (*Paedisca*).  
Synonymised by Meyrick (1881a, p. 488; 1883, p. 54).  
Auckland AK, D. Bolton; HT not found (K.R. Tuck, 1976), BMNH.  
Hudson 1928, p. 242, as synonym.

**voluta** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 39 (*Paedisca*). Synonymised by Meyrick (1881a, p. 488; 1883b, p. 54).  
[Nelson NN, T.R. Oxley]; HT ♂ designated by Felder, head and abdomen missing, BMNH.  
Hudson 1928, p. 242, as synonym.

**latomana** Meyrick, §1885b, p. 348; 1885g, p. 145 (*Harmologa*)  
Arthur's Pass NC/WD, 4700 ft, E. Meyrick; HT ♀ unique, mislabelled as from Mt Arthur, BMNH.

Hudson 1928, p. 243, pl. xxvi fig. 37 and 38, as *Cnephasia latomana*.

Note. There are 2 geographic colour patterns; Hudson illustrates the Mt Arthur NN pattern, not the typical one.

**melanophaea** Meyrick, 1927a, p. 698 (*Cnephasia*)  
Mount Arthur NN, 4,200 ft, Stella Hudson; LT ♂ selected by K.R. Tuck and here designated, labelled “Mt Arthur New Zealand SH 7.1.19”, “Cnephasia melanophaea Meyr. 1/1 E. Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, p. 243, pl. lii fig. 1 and 2.

**microbathra** Meyrick, 1911b, p. 62 (*Cnephasia*)  
West Plains, Invercargill SL, A. Philpott; HT ♂ unique, BMNH.  
Hudson 1928, p. 244, pl. xxvi fig. 39.

**ochnosema** Meyrick, 1936, pp. 281–282 (*Cnephasia*)  
Jack's Pass, Hanmer MB, S. Lindsay; HT ♂ designated by Meyrick, CMNZ.

Hudson 1939, p. 436, pl. lxi fig. 7.  
Note. The HT, labelled as “Type”, was returned to Lindsay at CMNZ. It is characteristic of all Lindsay material that Meyrick holotypes were sent back to CMNZ.

**paterna** Philpott, 1926a, p. 391 (*Cnephasia*)  
Little River, Banks Peninsula MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.  
Hudson 1928, p. 244, pl. lii fig. 26.

Also 17 undescribed species (NZAC).

• ***Ctenopseustis*** Meyrick, §1885b, p. 348; 1885g, p. 146. Type species *Teras obliquana* Walker, 1863, by original monotypy.

**fraterna** Philpott, 1930b, p. 7 (*Ctenopseustis*)  
Whangarei ND, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1928, pl. xxv fig. 3 (according to Hudson 1939, p. 433).  
Note. *Ctenopseustis fraterna* was synonymised with *C. obliquana* by Hudson (1939, p. 433), but shown to be a valid species by Green & Dugdale (1983).

**obliquana** Walker, 1863c, p. 302 (*Teras*) in the sense of Green & Dugdale (1983, p. 430)  
Auckland AK, D. Bolton; HT ♀ with abdomen missing, BMNH.  
Hudson 1928, p. 234, pl. xxv fig. 4–6 and 8–10.

**spurcatana** Walker, 1863c, p. 305 (*Teras*).  
Synonymised by Meyrick (1883b, p. 60).  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1928, p. 234, as synonym, after Meyrick (1883, p. 60).

**transtrigana** Walker, 1863c, p. 354 (*Sciaphila*).  
Synonymised by Meyrick (1881a, p. 487, with *spurcatana*; 1883b, p. 60, with *obliquana*).

[Nelson NN], T.R. Oxley; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 234, as synonym.

**turbulentana** Walker, 1863c, p. 355 (*Sciaphila*).  
Synonymised by Meyrick (1881a, p. 487, with *spurcatana*; 1883b, p. 60, with *obliquana*).  
[Nelson NN], T.R. Oxley; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 234, as synonym.

**ropeana** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 46 (*Teras*). Synonymised by Meyrick (1881a, p. 487, with *spurcatana*; 1883b, p. 60, with *obliquana*).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1928, p. 234, as synonym.

Note. The *ropeana* HT falls within the range of morphological variation shown by topotypic *inana*. Cross-attraction between populations from Nelson and Christchurch has yet to be investigated.

**herana** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 52 (*Teras*). Synonymised by Meyrick (1883b, p. 60).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 234, as synonym.

Note. As for *ropeana*, HT *herana* may yet prove to be conspecific with *Ctenopseustis inana*.

**charactana** Meyrick, 1881, p. 492 (*Cacoecia*). New synonymy.  
Auckland AK, E. Meyrick; HT ♀ unique, labelled "Auckland New Zealand 12.1.80", BMNH.  
Hudson 1928, p. 227, is irrelevant.

Note. The HT was found in a "supplementary" series of *C. obliquana* at BMNH. Meyrick had removed it from his series of *charactana*, all of which are of a *Coprosma*-eating tortricid superficially like *C. obliquana* which now lacks a name. See under *Planotortrix coprosmae* (p. 125).

**inana** Butler, 1877, p. 403, pl. xlili fig. 13 (*Cacoecia*) revised status

[?Christchurch MC or ?Dunedin DN], J. Hector; HT ♀ unique, BM genitalia slide no. 1069 ♀ (abdomen glued), BMNH.

Hudson 1928, p. 234, as synonym of *obliquana*.  
Note. *Cacoecia inana* was wrongly placed in synonymy with *Planotortrix excessana* (Walker) by Dugdale (1966, p. 396) and synonymised with *obliquana* by Green & Dugdale (1983, p. 430). The exact type locality is in doubt, but the material sent by Dr Hector was largely collected by J.D. Enys in mid Canterbury. Populations from this area are pheromonally distinctive, showing no cross-attraction with *Ctenopseustis obliquana* (Foster & Roelofs 1987).

**servana** Walker, 1863c, p. 306 (*Teras*)  
Auckland AK, D. Bolton; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, pl. xxv fig. 7, as *Ctenopseustis obliquana*.

**priscana** Walker, 1863c, p. 307 (*Teras*). Synonymised by Green & Dugdale (1983, p. 431).

[Auckland AK], D. Bolton; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 234, as synonym of *obliquana*.

**congestana** Walker, 1863c, p. 308 (*Teras*). Synonymised by Green & Dugdale (1983, p. 431).  
[Auckland AK], D. Bolton; HT ♂ unique, BM genitalia slide no. 20925 ♂, BMNH.  
Hudson 1928, p. 234, as synonym of *obliquana*.

**cuneiferana** Walker, 1866b, p. 1780 (*Teras*).  
Synonymised by Green & Dugdale (1983, p. 431).  
[Auckland AK], D. Bolton; LT ♂ designated by Green & Dugdale, BMNH.  
Hudson 1928, p. 234, as synonym of *obliquana*.

**abjectana** Walker, 1866b, p. 1781 (*Teras*?).  
Synonymised by Green & Dugdale (1983, p. 431).  
[Auckland AK], D. Bolton; HT ♂ unique, BM genitalia slide no. 3185 ♂, BMNH.  
Hudson 1928, p. 234, as synonym of *obliquana*.

**contractana** Walker, 1866b, p. 1782 (*Teras*).  
Synonymised by Green & Dugdale (1983, p. 431).  
[Auckland AK], A. Sinclair; HT ♂ unique, BM genitalia slide no. 20924 ♂, BMNH.  
Hudson 1928, p. 234, as synonym of *obliquana*.

Also 1 undescribed species (NZAC).

• **Curvisacculus** Dugdale, 1966b, p. 772. Type species *Tortrix encausta* Philpott, 1930, by original designation.

**philpotti** Dugdale, 1978, p. 443 (*Curvisacculus*). New name for *Tortrix encausta* Philpott, 1930, preoccupied by *Tortrix encausta* Meyrick (1907a, p. 735; Sri Lanka).

**encausta** Philpott, 1930b, pp. 6–7 (*Tortrix*)  
Kao ND, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 433, not figured.

**triorhota** Meyrick, 1927a, p. 698 (*Epichorista*)  
Wainuiomata WN, G.V. Hudson; HT ♂ unique, BM genitalia slide no. 10573, BMNH.  
Hudson 1928, p. 239, pl. lii fig. 28, as *Epichorista triorhota*.

Also 1 new species (NZAC).

• **Ecclitica** Meyrick, 1923, p. 164. Type species *Dipterina hemiclista* Meyrick, 1905, by original designation.

**hemiclista** Meyrick, 1905, p. 233 (*Dipterina*)  
Wellington WN, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 242, pl. xxvi fig. 14.

**torogramma** Meyrick, 1897b, pp. 388–389  
(*Cacoecia*)

» Tortricidae, *Ecclitica torogramma*

"New Zealand", G.V. Hudson; LT ♂ here designated, labelled "New Zealand GVH .95", "Tortrix torogramma Meyr. 4/5 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 228, pl. xxvi fig. 27, as *Tortrix torogramma*.

Note. Philpott (1928d, p. 447) transferred this species to *Ecclitica*.

● ***Epalxiphora*** Meyrick, 1881b, p. 648. Type species *E. axenana* Meyrick, 1881, by original monotypy.

*axenana* Meyrick, 1881b, p. 648 (*Epalxiphora*)

Wellington WN, E. Meyrick; HT ♀ unique, BMNH. Hudson 1928, pp. 233–234, pl. xxv fig. 44–53.

Note. The HT ♀ has the metathorax, hindwings, and abdomen of a ♂ glued expertly to it.

● ***Epichorista*** Meyrick, 1911c, p. 83. Type species *Proselena hemionana* Meyrick, 1882 (1883), by original designation. True *Epichorista*.

*aspistana* Meyrick, §1882c, p. 277; 1883b, p. 42 (*Proselena*)

[Porter's Pass, part of] Castle Hill MC, J.D. Enys; LT ♂ here designated, labelled "Castle Hill New Zealand JDE /82", "Epichorista aspistana Meyr. 4/6 E. Meyrick det. in Meyrick Coll.", BMNH. Hudson 1928, p. 237, pl. xxvi fig. 32.

*hemionana* Meyrick, §1882c, p. 277; 1883b, pp. 42 (key) and 43 (*Proselena*)

Lake Guyon MB/BR, R.W. Fereday; LT ♂ selected by J.D. Bradley and here designated, labelled "L. Guyon New Zealand RWF 3/82", "Epichorista hemionana Meyr. ♂ 1/3 E. Meyrick det. in Meyrick Coll.", "abdomen missing", BMNH. Hudson 1928, p. 235, pl. xxvi fig. 33.

● ***Epichorista*** of authors. Species with a costal fold on the ♂ forewing.

*allogama* Meyrick, 1914a, p. 105 (*Harmologa*)

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH 24.12.12", "Epichorista allogama Meyr. 1/12 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 238, pl. xxvi fig. 20 and 21.

*emphanes* Meyrick, 1901, p. 571 (*Proselena*)

Mt Peel NN, [resting on snow, 5,400 ft], G.V. Hudson; HT ♀ unique, not in BMNH.

Hudson 1928, p. 238, pl. xxvi fig. 15, 30, and 31; 1939, p. 434.

Note. Meyrick (1911b, p. 73) gave a description of the ♂.

***achrosta*** Meyrick, 1901, p. 572 (*Harmologa*). Synonymised by Meyrick (1923, p. 163).

Mt Arthur NN, 2,800 ft, G.V. Hudson; HT ♂ not in BMNH.

Hudson 1928, p. 238, as synonym.

***epicura*** Meyrick, 1911c, p. 86 (*Harmologa*). Synonymised by Meyrick (1911b, p. 74).

Castle Hill MC, 3,000 ft, E. Meyrick; HT ♂ unique, BMNH.

***theatralis*** Philpott, 1918, p. 128 (*Epichorista*). Synonymised by Philpott (1923, p. 150).

Mt Cleughearn FD, A. Philpott; ?HT ♀ unique, NZAC. Hudson 1928, p. 238, as synonym.

***candida*** Clarke, 1926, p. 419 (*Epichorista*). Synonymised by Hudson (1928, p. 238).

Hope Arm, Lake Manapouri FD, C.E. Clarke; HT ♀ unique, AMNZ.

Also 1 undescribed species (NZAC).

● ***Epichorista*** of authors. Species lacking a costal fold on the ♂ forewing.

***abdita*** Philpott, 1924c, pp. 664–665, fig. 2D,E (*Epichorista*)

Mt Arthur NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 434, as synonym of *E. emphanes*.

***crypsidora*** Meyrick, 1909a, p. 11 (*Dipterina*)

Invercargill SL, A. Philpott; HT ♂ unique, BMNH.

Hudson 1928, p. 239, pl. xxvi fig. 24 and 25, as *Epichorista crypsidora*.

***carcharodes*** Meyrick, 1914a, p. 104 (*Epichorista*). Synonymised by Hudson (1928, p. 239).

Haeo ND, G. V. Hudson; HT ♂ unique, BMNH.

***elephantina*** Meyrick, §1885b, p. 348; 1885g, p. 143 (*Proselena*)

Arthur's Pass NC/WD, 4,700 ft, E. Meyrick; HT ♂ unique, BMNH.

Hudson 1928, p. 237, pl. 1 fig. 16, as *Epichorista elephantina*.

***eribola*** Meyrick, 1889b, p. 156 (*Proselena*)

Otira River WD, 1,500 ft, E. Meyrick; LT ♂ here designated, labelled "Otira River New Zealand 1500 ft 24.1.83", "Epichorista eribola Meyr. 2/5 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 238, pl. xlvi fig. 13, as *Epichorista eribola*.

***fraudulenta*** Philpott, 1928, p. 363 (*Eurythecta*)

Mt Arthur NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 434, as synonym of *Epichorista emphanes*.

***lindsayi*** Philpott, 1928c, p. 181 (*Epichorista*)

Little River MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 435, pl. lvii fig. 26.

**mimica** Philpott, 1930b, p. 5 (*Epichorista*)  
Mt Ida CO, C.E. Clarke; HT ♂ designated by Philpott,  
AMNZ.  
Hudson 1939, p. 435, pl. lvii fig. 5.

**persecta** Meyrick, 1914a, p. 104 (*Epichorista*)  
Tisbury [Invercargill], SL, A. Philpott; LT ♂ selected by  
J.D. Bradley and here designated, labelled "Tisbury  
New Zealand AP 24.11.10", "Epichorista persecta  
Meyr. 3/1 Meyrick det. in Meyrick Coll.", "BM gen-  
italia slide no. 10663 ♂", BMNH.  
Hudson 1928, p. 237, pl. xlvi fig. 20 and 21.

**semicocata** Meyrick, 1914a, p. 104 (*Epichorista*),  
as variety  
Tisbury [Invercargill] SL, A. Philpott; HT ♂ labelled  
"Tisbury New Zealand AP 3.11.11", "Epichorista per-  
secta Meyr. 3/3 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 237, pl. xxvi fig. 9, as variety.

**siriana** Meyrick, 1881a, pp. 521–522 (*Tortrix*)  
Hamilton WO, E. Meyrick; LT ♂ selected by D.J. Carter  
and here designated, labelled "Hamilton New Zealand  
18.1.80", "Epichorista siriana Meyr. 1/5 E. Meyrick  
det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 23, pl. xxvi fig. 34, as *Epichorista siriana*,  
following Meyrick (1911c, p. 83).  
Note. This species and "*Eurythecta*" *eremana* Meyrick  
(see below) are probably North Island and South Island  
forms respectively of the one species, as Hudson (1928,  
p. 237) suspected.

**speciosa** Philpott, 1927d, pp. 83–84 (*Epichorista*)  
Arthur's Pass NC/WD, S. Lindsay; HT ♂ designated by  
Philpott, CMNZ.  
Hudson 1939, p. 434, pl. lvii fig. 17.

**tenebrosa** Philpott, 1917b, pp. 243–244 (*Epichorista*)  
Ben Lomond OL, C.E. Clarke; HT ♂ designated by Phil-  
pott, AMNZ.  
Hudson 1928, p. 237, pl. xlvi fig. 10 and 11.

**zatrophana** Meyrick, §1882c, p. 277; 1883b, pp. 44  
(key) and 46 (*Harmologa*)  
Christchurch MC, E. Meyrick; HT ♀ unique, not in  
BMNH.  
Hudson 1928, p. 238, pl. xlvi fig. 16 (Arthur's Pass NC/WD  
specimen), as *Epichorista zatrophana*.

Note. The identity of the true *zatrophana* Meyrick is in  
doubt. The HT ♀ was collected at Christchurch on  
19 March 1882 to light during a "fresh NW wind".  
Meyrick's subsequent redescription (1883b, p. 44),  
based on males from Arthur's Pass NC/WD, may relate  
to a different species, particularly as these were col-  
lected on 25 and 29 January 1883, at altitudes over  
3,000 ft. [Information from Meyrick's Diary of  
Captures.]

• "*Eurythecta*" of authors, but not Meyrick (1883,  
p. 56, q.v.)

Species placed in this genus are small, usually slender tortricids characteristic of dense, damp swards.

**curva** Philpott, 1918, p. 127 (*Eurythecta*)  
Hunter Mountains FD, 3500 ft, A. Philpott; HT ♂ des-  
ignated by Philpott, NZAC.  
Hudson 1928, p. 225, pl. xlvi fig. 14.

**eremana** Meyrick, §1885b, p. 348; 1885g, p. 144  
(*Proselena*)  
Castle Hill MC, 2,000 ft, E. Meyrick; LT ♂ selected by  
K.R. Tuck and here designated, labelled "Castle Hill  
New Zealand 2000 ft 18/1/83", "Eurythecta eremana  
Meyr. 7/13 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 225, as *Eurythecta eremana*; notes simi-  
larity to *Epichorista siriana* (see above).

**leucothrinca** Meyrick, 1931b, p. 367 (*Eurythecta*)  
Black Hill [Rakaia Valley] MC, S. Lindsay; HT ♂ unique,  
CMNZ.  
Hudson 1939, p. 432, pl. lvi fig. 18.

**loxias** Meyrick, 1888d, p. 74 (*Proselena*)  
Mt Arthur NN, E. Meyrick; LT ♂ here designated, labelled  
"Mt Arthur New Zealand 4000 ft 15.1.86", "Eury-  
thecta loxias Meyr. 2/3 E. Meyrick det. in Meyrick  
Coll.", BMNH.  
Hudson 1928, p. 225, pl. xxvi fig. 22, as *Eurythecta loxias*.

**phaeoxyla** Meyrick, 1938, p. 427 (*Eurythecta*)  
Mount Torlesse MC, S. Lindsay; HT ♂ designated by  
Meyrick, CMNZ.  
Hudson 1939, p. 432, pl. lvi fig. 8.

• **Gelophaula** Meyrick, 1923, p. 163. Type species  
*Harmologa trisulca* Meyrick, 1916, by original  
designation.

**aenea** Butler, 1877, p. 402 (*Teras*)  
[Mt Torlesse MC], J.D. Enys; HT ♂ unique, BMNH.  
Hudson 1928, p. 235, not figured; p. 236, pl. xxv fig. 13  
and 14, as *Gelophaula siraea*.

**aridella** Clarke, 1934, p. 13 (*Gelophaula*)  
Flat Top, Hunter Mountains FD, C.E. Clarke; HT ♂ des-  
ignated by Clarke, AMNZ.  
Hudson 1939, p. 434, pl. lvi fig. 23.

**lychnophanes** Meyrick, 1916b, p. 415 (*Harmologa*)  
Mt Arthur NN, 4,500 ft, G.V. Hudson; HT ♂ unique,  
BMNH.  
Hudson 1928, p. 236, pl. xlvi fig. 13, as *Gelophaula lychnophanes*.

**palliata** Philpott, 1914, p. 120 (*Harmologa*)  
Obelisk, Old Man Range CO, W.G. Howes; HT ♂ des-  
ignated by Philpott, NZAC.  
Hudson 1928, p. 235, pl. xxv fig. 35, pl. xlvi fig. 25, as  
*Gelophaula palliata*.

➤ Tortricidae, *Gelophaula*

***praecipitalis*** Meyrick, 1934, p. 152 (*Gelophaula*)  
Mt Peel NN, Lawford White; LT ♂ here designated,  
labelled "Mt Peel 26/3/32", "Gelophaula praecipitalis  
Meyr." (in S. Lindsay's handwriting), CMNZ.  
Hudson 1939, p. 434, pl. lxi fig. 16.

***siraea*** Meyrick, §1885b, p. 348; 1885g, pp. 145–146  
(*Harmologa*)

Arthur's Pass NC/WD, E. Meyrick; LT ♂ here designated, labelled "Arthurs Pass New Zealand 29.i.1883  
1676", "Harmologa siriana Meyr. named by Meyrick", "Walsingham Collection 1910–427", BMNH.

Hudson 1928, p. 236, pl. xlvi fig. 29, as *Gelophaula brevicula*.

Note. Meyrick appears to have given some of his 1883 material to Walsingham, and as the specimen labels agree with Meyrick's Diary of Captures I have no hesitation in making the Walsingham ♂ the LT.

***brevicula*** Meyrick, 1921, p. 334 (*Harmologa*). New synonymy.

Arthur's Pass NC/WD, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 236, pl. xlvi fig. 30, as species.

***tributaria*** Philpott, 1913, pp. 77–78 (*Harmologa*)  
Obelisk, Old Man Range CO, W.G. Howes; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 236, pl. xxv fig. 34, as *Gelophaula tributaria*.

***trisulca*** Meyrick, 1916b, p. 414 (*Harmologa*)

Arthur's Pass NC/WD, G.V. Hudson; LT ♂ here designated, labelled "Arthurs Pass New Zealand GVH 3500'  
12.14", "Gelophaula trisulca Meyr. 1/4 E. Meyrick det.  
in Meyrick Coll.", BMNH.

Hudson 1928, p. 235, pl. xxv fig. 15 and 16.

***vana*** Philpott, 1928g, p. 487 (*Gelophaula*)

Mt Cleugharn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 434, not figured.

Also 5 undescribed species (NZAC).

● ***Harmologa*** Meyrick, §1882c, p. 277; 1883b, pp. 36 (key) and 44. Type species *Teras oblongana* Walker, 1863, by subsequent designation (Fernald 1908, p. 44).

***Trachybathra*** Meyrick, 1907c, p. 113. Type species *Trachybathra scoliastis* Meyrick, 1907. Synonymised by Meyrick (1911c, p. 85).

***amplexana*** Zeller, 1875, p. 222 (*Idiographis*?)

[exact provenance unknown], New Zealand; LT ♂ selected by D.J. Carter and here designated, labelled "Tortrix Idiographis amplexana Z., N. Seeland l.w", "TORTRIX (IDIOPHRASIS) AMPLEXANA Zeller Ver. Z-B. Ges. Wien xxv: 222–3, pl. viii: 2 Named by Zeller, Type ♂", BMNH.

Hudson 1928, p. 239, pl. xxiv fig. 2 and 3, as *Harmologa amplexana*.

***vilis*** Butler, 1877, p. 42, pl. 45 fig. 15 (*Cacoecia*).  
Synonymised by Meyrick (1881a, p. 494).  
[mid Canterbury MC], J.D. Enys; HT ♂ designated by Butler, BMNH.  
Hudson 1928, p. 239, as synonym.

***columella*** Meyrick, 1927a, p. 699 (*Harmologa*)  
Arthur's Pass NC/WD, Stella Hudson; HT ♂ unique,  
abdomen missing, BMNH.  
Hudson 1928, p. 241, pl. lii fig. 4.

***festiva*** Philpott, 1915, p. 199 (*Harmologa*)  
Mt Cleugharn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 240, pl. xxvii fig. 21.

***oblongana*** Walker, 1863c, p. 303 (*Teras*)  
[Nelson NN], T.R. Oxley; HT ♀ unique, BM genitalia slide no. 5753, BMNH.  
Hudson 1928, p. 239, pl. xxvi fig. 16, as *Harmologa oblongana*.

***inaptana*** Walker, 1863c, p. 304 (*Teras*).  
Synonymised by Meyrick (1881a, p. 489).  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1928, p. 239, as synonym.

***cuneigera*** Butler, 1880, p. 559 (*Teras*).  
Synonymised by Meyrick (1883b, p. 45).  
[Blenheim MB], W. Skellon; HT ♀ unique, BM genitalia slide no. 10599, BMNH.  
Hudson 1928, p. 239, as synonym.

***indomita*** Philpott, 1930b, p. 6 (*Tortrix*). New synonymy.  
Dunedin DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 432, pl. lvii fig. 1, as species.  
Note. Examination of the genitalia showed no differences between *indomita* and *oblongana*; the ♂ genitalia are characteristic (see Philpott 1928, pp. 446 and 456, fig. 12).

***petriana*** Meyrick, 1901, p. 572 (*Harmologa*)  
Invercargill SL, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 243, pl. xxvii fig. 5, as *Cnephasia petriana*.  
Note. Philpott (1928d, p. 446) transferred *petriana* back to *Harmologa* on ♂ genital characters.

***pontifica*** Meyrick, 1911b, p. 74 (*Harmologa*)  
Mt Arthur NN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 240, pl. xxv fig. 32.

***reticularis*** Philpott, 1915, pp. 199–200 (*Harmologa*)  
Longwood Range SL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 241, pl. xxvii fig. 8.

***sanguinea*** Philpott, 1915, p. 199 (*Harmologa*)  
Mt Cleugharn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, pp. 241–242, pl. xxvii fig. 14.

**scoliastis** Meyrick, 1907c, p. 113 (*Trachybathra*)  
Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, head  
missing, BMNH.  
Hudson 1928, p. 240, pl. xxiv fig. 4.

**sisyrana** Meyrick, §1882c, p. 277; 1883b, p. 44  
(*Harmologa*)

[Kaiapoi MC, sandhills], E. Meyrick; LT ♂ selected by  
D.J. Carter and here designated, labelled "Christ-  
church New Zealand 31.3.82", "Harmologa sisyrana  
Meyr. 3/1 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 240, pl. xxvi fig. 17.

**antitypa** Meyrick, 1914a, p. 105 (*Harmologa*)  
Wellington WN, G.V. Hudson; HT ♂ unique, abdomen  
missing, BMNH.

Hudson 1928, p. 240, as synonym of *sisyrana*.

**speciosa** Philpott, 1927d, pp. 83–84 (*Epichorista*)  
new combination

Arthur's Pass NC/WD, S. Lindsay; HT ♂ designated by  
Philpott, CMNZ.  
Hudson 1939, p. 434, pl. lvii fig. 17.

**torotermia** Hudson, 1925, p. 221 (*Harmologa*)  
Mount Ida CO, C.E. Clarke; HT ♂ unique, NMNZ.  
Hudson 1928, p. 241, pl. lii fig. 23.

Also 2 undescribed species (NZAC).

● **Ochetarcha** Meyrick, 1924b, p. 661. Type species  
*Olindia miraculosa* Meyrick, 1917a, by original  
designation.

**miraculosa** Meyrick, 1917a, p. 246 (*Olindia*)  
Wainuiomata WN, Stella Hudson; HT ♀ unique, geni-  
talia mounted with specimen, BMNH.

Hudson 1928, p. 244, pl. xxv fig. 33, pl. xlvi fig. 9.  
Note. *Paedisca mahiana* Felder, 1875, pl. cxxxvii fig. 40,  
thought by Meyrick to be a possible synonym, was  
examined by me and is a senior synonym of *Episimus*  
*encaustica* Meyrick, 1922a, p. 518 (Olethreutinae;  
Brazil). Dugdale (1966b, p. 766) wrongly placed *Och-  
etarcha* in Cnephasiini; on genitalia and external char-  
acters it closely resembles *Terricula* Falkovitsh, 1965,  
p. 418, from Japan and Ussuri.

● **Philocryptica** Meyrick, 1923, p. 164. Type species  
*Harmologa polypodii* Watt, 1921b, by monotypy.

**polypodii** Watt, 1921b, pp. 257–258 (*Harmologa*)  
Wellington WN, M.N. Watt; LT ♂ here designated,  
labelled "Wellington 11/11/19," NMNZ.  
Hudson 1928, p. 241, pl. xlvi fig. 33.

● **Planotortrix** Dugdale, 1966, p. 392. Type species  
*Teras excessana* Walker, 1863, by original  
designation.

**clarkei** Philpott, 1930b, pp. 5–6 (*Tortrix*)  
Waimarino TO, C.E. Clarke; HT ♂ designated by Phil-  
pott, AMNZ.  
Hudson 1939, pp. 432–433, pl. lvii fig. 2.

**conditana** Walker, 1863c, p. 306 (*Teras*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BM genitalia  
slide no. 3186 ♂, BMNH.

Hudson 1928, p. 229, pl. xxiv fig. 42–44.

**astrologana** Meyrick, 1889b, p. 156 (*Cacoecia*).  
Synonymised by Meyrick (1911, p. 84).  
Wellington WN, G.V. Hudson; LT ♂ selected by J.D.  
Bradley, 1963, and designated by Dugdale (1966, p. 396),  
BM genitalia slide no. 2328 ♂, BMNH.  
Hudson 1928, p. 229, as synonym.

Note. Examination of the type material, particularly ♂  
genitalia, leads me to restrict the *conditana* synonymy as  
above.

**coprosmae** Dugdale, new species (proposed here) for  
*charactana* of authors, but not Meyrick (1881,  
p. 492) (*Tortrix*)

[Riccarton Bush] MC, E. Meyrick; HT ♂ selected as NT  
by J.D. Bradley, 1962, labelled "Christchurch New  
Zealand 18/4/82", "Tortrix charactana Meyr. 1/12 E.  
Meyrick det. in Meyrick Coll.", BM genitalia slide no.  
8914 ♂, BMNH.

Hudson 1928, p. 227, pl. xxiv fig. 33–35, as *Tortrix*  
*charactana*.

Note. The long-mislaid HT of *Cacoecia charactana*  
Meyrick has been found, and is identical with *Cten-  
opseustis obliquana* (p. 121). The *Coprosma*-feeding  
species is well represented in most collections (invari-  
ably under the name *charactana*), and has been  
described (Meyrick 1883, p. 50; Hudson 1928, p. 227;  
Philpott 1929, pp. 450–451, fig. 60 (p. 465) ♂ geni-  
talia). I have honoured Dr J.D. Bradley's hard work  
on my behalf, and the hospitality of BMNH, by using  
his NT as the HT for *coprosmae*. Although this specie-  
s lacks a costal fold on the ♂ forewing, it is placed  
in *Planotortrix* on genital and habitus characters.

**excessana** Walker, 1863c, p. 303 (*Teras*)  
[Nelson NN], T.R. Oxley; LT ♂ selected by J.D. Bradley,  
1962, and designated by Dugdale (1966a, p. 396),  
BMNH.

Hudson 1928, p. 230, pl. xxiv fig. 5, 6, and 27–30.

**biguttana** Walker, 1863c, p. 305 (*Teras*). Syn-  
onymised by Meyrick (1883b, p. 48).  
[Nelson NN], T.R. Oxley; LT ♂ selected by J.D. Bradley,  
1962, and designated by Dugdale (1966a, p. 396), BMNH.

Hudson 1928, p. 230, as synonym.

Note. Studies of cross-attraction show that *P. excessana*  
of authors embraces two species (Galbreath 1985, Foster  
*et al.* 1986). Both LT *excessana* and LT *biguttana* fall  
within the range of costal fold / forewing length ratios  
characteristic of laboratory colonies of "P. excessana Type  
B", but not "Type A", of Foster *et al.* (1986, p. 156, table).

**fastigata** Philpott, 1916, p. 422 (*Tortrix*) new  
combination

> Tortricidae, *Planotortrix fastigata*

Mt Cleughearn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 231, pl. xliv fig. 26 and 27.

**ascomorpha** Meyrick, 1934, p. 153 (*Tortrix*). New synonymy.

Arthur's Pass NC/WD, S. Lindsay; LT ♀ designated by Dugdale (1966a, p. 396), CMNZ.

Hudson 1939, p. 433, pl. lvii fig. 19, as species.

Note. Material collected by Mr B. Patrick from Longwood Range SL clearly necessitates the above synonymy.

**flammea** Salmon, 1956, p. 575 (*Bactra*)

Homer Forks FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.

Salmon 1956, pl. 22 fig. 2 and 3.

**flavescens** Butler, 1877, p. 402 (*Teras*). New combination.

?Dunedin DN, J. Hector; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 231, pl. xxv fig. 1 and 2, as *Tortrix flavescens*.

**acrocausta** Meyrick, 1907c, p. 115 (*Cacoecia*).

Synonymised by Hudson (1928, p. 231).

[Riccarton Bush], Christchurch MC, E. Meyrick; LT ♂ selected by J.D. Bradley and here designated, labelled "Christchurch New Zealand 22/9/82", "BM genitalia slide no. 8604 ♂", BMNH.

**inusitata** Philpott, 1919, p. 225 (*Tortrix*). New synonymy.

Waitati DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1928, pp. 232–233, pl. xlvi fig. 28.

Note. Philpott (1928d, p. 451) noted that the ♂ genitalia of *inusitata* and *flavescens* agree "almost exactly", but because of the "small number of [*inusitata*] ... it is not thought advisable to unite the species at this juncture". Examination of large series reared by the Forest Biology Survey, Forest Research Institute, Rotorua, showed the synonymy to be valid.

**notophaea** Turner, 1926, p. 135 (*Tortrix*)

Epping N.S.W., Australia; HT ♂ unique, transferred from ANIC to NZAC.

Hudson 1928, pl. xxiv fig. 6, as *Tortrix excessana*.

**distincta** Salmon, 1948, p. 310 (*Ctenopseustis obliquana* subspecies). Synonymised by Dugdale (1966, p. 397).

Great Island, Three Kings Islands; HT ♂ designated by Salmon, AMNZ.

**orthocopa** Meyrick, 1924b, p. 661 (*Tortrix*)

Wellington WN, G.V. Hudson; LT ♂ selected by J.D. Bradley, 1963, and designated by Dugdale (1966, p. 397), abdomen missing, BMNH.

Hudson 1928, p. 229, pl. xxiv fig. 1.

**orthropis** Meyrick, 1901, p. 573 (*Cacoecia*)

Dunedin DN, [reserved bush and forest], E. Meyrick; LT

♂ selected by J.D. Bradley, 1962, and designated by Dugdale (1966a, p. 397), BM genitalia slide no. 8924 ♂, BMNH.

Hudson 1928, p. 228, pl. xxiv fig. 11, 31, and 32.

**pictoriana** Felder & Rogenhofer, 1875, pl. cxxxvi fig. 55 (*Grapholitha*)

[Nelson NN, T.R. Oxley, 1864]; HT ♂ unique, BM genitalia slide no. 8913 ♂, BMNH.

Hudson 1928, p. 226, pl. xxiv fig. 47 and 48, as *Tortrix pictoriana*.

Note. Meyrick (1913e, p. 39) lists this species under *Eulia*. Horak (1984) treated it as a *Planotortrix*.

**spatiosa** Philpott, 1923, pp. 150–151 (*Tortrix*)

Dun Mountain NN/MB, A. Philpott; HT ♀ designated by Philpott, NZAC.

Hudson 1928, p. 229, pl. xl ix fig. 5.

Note. Dugdale (1966, p. 397) erroneously reversed the HT and AT designations of Philpott.

**syntona syntona** Meyrick, 1909b, pp. 73–74 (*Cacoecia*, as species)

[Port Ross], Auckland Islands, J.S. Tennant; HT ♂ unique, BM genitalia slide no. 3104 ♂, BMNH.

Hudson 1928, p. 229, pl. xxiv fig. 46 (♂, not ♀), as species.

**syntona laqueorum** Dugdale, 1971b, p. 162 (*Planotortrix*, as subspecies). Amended name.

The Snares islands, P.M. Johns; HT ♂ designated by Dugdale, NZAC.

Note. The epithet "laqueorum" is inadmissible, and is here amended to the correct spelling, 'laqueorum'.

**taipana** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 46 (*Tortrix*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, BM genitalia slide no. 10662 ♂, BMNH.

Not mentioned by Hudson.

**enopla** Meyrick, §1882c, p. 277; 1883b, pp. 44 (key) and 49–60 *Cacoecia*. Synonymised by Meyrick (1911, p. 84).

Wellington WN, R.W. Fereday; HT ♂ unique, BM genitalia slide no. 8927 ♂, BMNH.

Hudson 1928, p. 229, as synonym of *Tortrix conditana*.

Note. Male genital characters separate these 2 type specimens from those of *conditana* and *astrologana*.

**tigris** Philpott, 1914, p. 120 (*Tortrix*)

Tisbury, [Invercargill] SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 229, pl. xl v fig. 9.

Also 4 undescribed species (NZAC).

• **Pyrgotis** Meyrick, 1881a, p. 439, in the sense of Dugdale (1971b, p. 164). Type species *Conchylis plagiata* Walker, by subsequent designation (Fernald 1908, p. 42); not *Pyrgotis insignana* Meyrick, 1881a, p. 440 (Australia), designated by Meyrick (1913e, p. 8).

- arcuata*** Philpott, 1915, p. 198 (*Capua*)  
Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 223, pl. xlvi fig. 27, as *Capua arcuata*;  
Dugdale 1971b, p. 165, as *Pyrgotis arcuata*.
- calligypsa*** Meyrick, 1926b, p. 415 (*Catamacta*)  
Wellington WN, G.V. Hudson; HT ♀ unique, abdomen missing, BMNH.  
Hudson 1928, p. 221, pl. lii fig. 25, as *Catamacta calligypsa*.
- variegata*** Philpott, 1930b, pp. 4–5 (*Capua*). New synonymy.  
Wairakei TO, C.E. Clarke; HT ♂ designated by Philpott, left forewing missing, AMNZ.  
Hudson 1939, p. 431, pl. lvii fig. 3, as *Capua variegata*. Note. Reared material yielded males and females indistinguishable from the 2 type specimens.
- chrysomela*** Meyrick, 1914a, p. 103 (*Catamacta*)  
Kaeo ND, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 221, pl. xxvii fig. 24, as *Catamacta chrysomela*; Dugdale 1971b, p. 165, as *Pyrgotis chrysomela*.
- consentiens*** Philpott, 1916, p. 421 (*Pyrgotis*)  
Cleughearn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, pp. 219–220, pl. xlvi fig. 10.  
Note. *Pyrgotis consentiens* and *P. humilis* (see below) may well be allopatric populations of a single species. They have the same host plant, and the genitalia are very similar.
- eudorana*** Meyrick, §1885b, p. 348; 1885g, p. 143 (*Pyrgotis*)  
[New Plymouth], “Taranaki” TK, E. Meyrick; HT ♀ unique, BM genitalia slide no. 8527, BMNH.  
Hudson 1928, p. 219, pl. xxii fig. 31.  
Note. Meyrick (1911b, p. 62) described the ♂.
- humilis*** Philpott, 1930b, p. 4 (*Pyrgotis*)  
Mount Maungatua DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, p. 431, not figured.
- plagiatana*** Walker, 1863c, p. 370 (*Conchyliis*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 222, pl. xxiv fig. 13–19 and 45, as *Capua plagiatana*; Dugdale 1971b, p. 166, as *Pyrgotis plagiatana*.
- recusana*** Walker, 1863c, p. 371 (*Conchyliis*). Synonymised by Meyrick (1881a, p. 441).  
[Auckland AK], D. Bolton; HT ♂ unique, BM genitalia slide no. 8516, BMNH.  
Hudson 1928, p. 222, as synonym.
- luciplagana*** Walker, 1863c, p. 381 (*Paedisca*). Synonymised by Meyrick (1881a, p. 441).
- [Nelson NN], T.R. Oxley; HT ♀ unique, BM genitalia slide no. 7809, BMNH.  
Hudson 1928, p. 222, as synonym.
- punana*** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 43 (*Grapholitha*). Synonymised by Meyrick (1881a, p. 441).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 222, as synonym.
- xylinana*** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 44 (*Grapholitha*). Synonymised by Meyrick (1881a, p. 441).  
[Nelson NN, T.R. Oxley]; HT ♀ unique, abdomen missing, remains of an abdomen glued on, BMNH.  
Hudson 1928, p. 222, as synonym.
- trichroa*** Meyrick, 1901, p. 578 (*Catamacta*).  
Synonymised by Dugdale (1971, p. 166).  
Whangarei [Heads] ND, E. Meyrick; HT ♀ unique, BM genitalia slide no. 10553, BMNH.  
Hudson 1928, p. 222, as synonym.
- tornota*** Meyrick, 1907c, p. 114 (*Pyrgotis*).  
Synonymised by Meyrick (1914a, p. 104).  
Invercargill SL, A. Philpott; HT ♀ unique, BM genitalia slide no. 8528, BMNH.
- parallela*** Salmon & Bradley, 1956, p. 72 (*Epagoge*).  
Synonymised by Dugdale (1971, p. 166).  
?Auckland Island, J.H. Sorensen; HT ♀ designated by Salmon & Bradley, NMNZ.  
Note. There was room for error in the labelling of Cape Expedition material (R.G. Ordish, pers. comm.); see *Scoaria albafascicula* and *Xanthorhoe subantarctica* for other examples.
- plinthoglypta*** Meyrick, 1892, p. 218 (*Pyrgotis*)  
Wellington WN, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.  
Hudson 1928, p. 223, pl. xxvi fig. 4, as *Capua plinthoglypta*.
- pyramidias*** Meyrick, 1901, p. 571 (*Pyrgotis*)  
Invercargill SL, A. Philpott; HT ♂ unique, BM genitalia slide no. 8514, BMNH.  
Hudson 1928, p. 219, pl. xxiv fig. 12.  
Note. The HT is small and drab, and may represent the entity browsing on *Cyatethodes* (*Styphelia*) rather than that on *Nothofagus*. The degree of interaction between these ‘forms’ is not known.
- transfixa*** Meyrick, 1924a, p. 203 (*Catamacta*)  
Wellington WN, G.V. Hudson; HT ♂ unique, BM genitalia slide no. 8520, BMNH.  
Hudson 1928, p. 221, pl. l fig. 5, as *Catamacta transfixa*; Dugdale 1971b, p. 165, as *Pyrgotis transfixa*.
- zygiana*** Meyrick, §1882c, p. 277; 1883b, p. 39 (*Pyrgotis*)  
[Riccarton Bush], Christchurch MC, E. Meyrick; HT ♂ unique, BM genitalia slide no. 8521 ♂, BMNH.  
Hudson 1928, p. 222, not figured, as *Capua zygiana*; Dugdale 1971b, p. 165, as *Pyrgotis zygiana*.

> Tortricidae, *Pyrgotis*

Also 2 undescribed species (NZAC; B. Patrick Collection, Dunedin).

- **Sorensenata** Salmon & Bradley, 1956, p. 73. Type species *Sorensenata agilitata* Salmon & Bradley, by original designation.

**agilitata** Salmon & Bradley, 1956, p. 73  
(*Sorensenata*)  
Campbell Island, J.H. Sorenson; HT ♂ designated by Salmon and Bradley, NMNZ.

- **Sperchia** Walker, 1869, p. 83. Type species *Sperchia intractana* Walker, by original monotypy.

**intractana** Walker, 1869, p. 83 (*Sperchia*)  
Melbourne Vic., Australia; type specimen in NMVA.  
New Zealand: adventive, first recorded by Philpott (1924c, p. 664) in 1923.  
Hudson 1928, pp. 223–224, pl. li fig. 24, as *Capua intractana*.

- **Tortrix** of authors, e.g., Meyrick (1912e, 1913e), but not in the sense of Obraztsov (1955, p. 181).

**antichroa** Meyrick, 1919, p. 351 (*Tortrix*)  
Mount Egmont TK, G.V. Hudson; HT ♂ unique, BM genitalia slide no. 2320 ♂, BMNH.  
Hudson 1928, p. 232, pl. xlvi fig. 1.

**demiana** Meyrick, §1882c, p. 277; 1883b, pp. 50 (key) and 51 (*Tortrix*)  
Rakaia MC, W.H. Gaze; HT ♂ unique, BM genitalia slide no. 8611 ♂, BMNH.  
Hudson 1928, pp. 226–227, pl. xxiv fig. 37.

**fervida** Meyrick, 1901, pp. 572—573 (*Cacoecia*)  
Kaitoke WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 231, pl. xxvi fig. 1 and 2, as *Tortrix fervida*.

**incendiaria** Meyrick, 1923, p. 164 (*Ecclitica*)  
Mt Egmont TK, 4,000 ft, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 242, pl. xl ix fig. 32, as *Ecclitica incendiaria*.  
Note. Philpott (1928, p. 447) removed this species to *Tortrix*.

**molybditis** Meyrick, 1907c, pp. 115–116 (*Tortrix*)  
Wellington WN, G.V. Hudson; HT ♂ unique, BM genitalia slide no. 8579 ♂, BMNH.  
Hudson 1928, pp. 231–232, pl. xxvi fig. 3.

**sphenias** Meyrick, 1909, p. 11 (*Cacoecia*)  
Invercargill SL, A. Philpott; HT ♂ unique, BMNH.  
Hudson 1928, p. 231, pl. xl v fig. 28, as *Tortrix sphenias*.

**zestodes** Meyrick, 1924a, p. 203 (*Tortrix*)  
Flora [Hut], Mount Arthur NN, Stella Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 232, pl. i fig. 20.

**Tribe Schoenotenini**  
(in the sense of Common 1965b, p. 656)

- **Dipterina** Meyrick, 1881, p. 527. Type species *Dipterina imbriferana* Meyrick, by subsequent designation (Walsingham 1907, p. 697).

**imbriferana** Meyrick, 1881a, pp. 527–528  
(*Dipterina*)  
Wellington WN, E. Meyrick; LT ♂ selected by J.D. Bradley, labelled "Wellington New Zealand 1.1.80", "BM genitalia slide no. 10572 ♂", "Cnephasia imbriferana Meyr. 7/7 E. Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, pp. 243–244, pl. iii fig. 2 (larva), pl. xxvi fig. 40–42, as *Cnephasia imbriferana*.

- **Maoritenes** Dugdale, 1966b, p. 761. Type species *Epagoge cyclobathra* Meyrick, 1907, by original designation.

**cyclobathra** Meyrick, 1907c, p. 114 (*Epagoge*)  
Invercargill SL, A. Philpott; LT ♂ selected by J.D. Bradley, labelled "Invercargill New Zealand AP .06", "Capua cyclobathra Meyr. 2/4 E. Meyrick det. in Meyrick Coll.", "BM genitalia slide no. 8572 ♂", BMNH.  
Hudson 1928, p. 222, pl. xxvi fig. 28, as *Capua cyclobathra*.

**modesta** Philpott, 1930b, p. 4 (*Pyrgotis*)  
Waiho Gorge WD, C.E. Clarke; HT ♂ unique, AMNZ.  
Hudson 1939, p. 431, pl. lvii fig. 4, as *Pyrgotis modesta*.

- **Prothelymna** Meyrick, §1882c, pp. 277–278; 1883b, pp. 36 (key) and 57. Type species *P. nephelotana* Meyrick, 1882 (= *Teras antiquana* Walker), by original designation.

**antiquana** Walker, 1863c, p. 397 (*Teras*)  
[Auckland AK], D. Bolton; LT ♀ selected by J.D. Bradley, labelled "Teras antiquana Wkr Cat. Lep. Het. BM 28 p.307 (1863) Type ♀", "New Zeal. // 54.4", "abdomen missing", BMNH.  
Hudson 1928, p. 21, pl. xl v fig. 29, as *Proselena antiquana*.

**maoriana** Walker, 1863c, p. 308 (*Teras*). Synonymised by Meyrick (1911c, p. 80).  
[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1928, p. 219, as synonym.  
Note. HT represented by right forewing, hindwing, and a scrap of thorax.

**fusiferana** Walker, 1863c, p. 355 (*Sciaphila*).  
Synonymised by Meyrick (1911c, p. 80).

[Auckland AK], D. Bolton; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 219, as synonym.

***spoliatana*** Walker, 1863c, p. 356 (*Sciaphila*).  
Synonymised by Meyrick (1911c, p. 80).

[Auckland AK], D. Bolton; LT ♂ here designated, labelled "New Zeal. // 54.4", "BM genitalia slide no. 10552 ♂", BMNH.

Hudson 1928, p. 219, as synonym.

***vetustana*** Walker, 1863c, p. 358 (*Olindia*).  
Synonymised by Meyrick (1911c, p. 80).

[Auckland AK], D. Bolton; HT ♀ unique, abdomen and genitalia on BM genitalia slide no. 10551, specimen unspread, BMNH.

Hudson 1928, p. 218, as synonym.

***morosana*** Walker, 1863c, p. 382 (*Paedisca*).  
Synonymised by Meyrick (1911c, p. 80).

[Auckland AK], D. Bolton; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 219, as synonym.

***accensa*** Walker, 1864b, p. 983 (*Teras*).  
Synonymised by Meyrick (1911c, p. 80).

[Auckland AK], D. Bolton; HT ♂ unique, abdomen and genitalia on BM genitalia slide no. 10576, BMNH.

Hudson 1928, p. 219, as synonym.

***nephelotana*** Meyrick, §1882c, p. 278; 1883b,  
p. 57 (*Prothelymna*).  
Synonymised by Meyrick (1911c, p. 80).

[Lyttelton], Christchurch MC, E. Meyrick; HT ♂ unique,  
abdomen and genitalia on BM genitalia slide no. 3326,  
BMNH.

Hudson 1928, p. 219, as synonym.

***niphostrota*** Meyrick, 1907c, pp. 112–117  
(*Proselena*)

Invercargill SL, A. Philpott; HT ♂ unique, abdomen and  
genitalia on BM genitalia slide no. 10578, BMNH.

Hudson 1928, p. 219, pl. xxxiii fig. 14, as *Proselena*  
*niphostrota*.

### Tribe Tortricini

(in the sense of Obraztsov 1955, p. 178)

● ***Acleris*** Hübner, 1825, p. 384. Type species [*Tortrix*] *aspersana* Hübner, 1817, by subsequent designation (Fernald 1908, pp. 12 and 55).

***comariana*** Lienig & Zeller, 1846, p. 263 (*Acleris*)  
Palaearctic.


New Zealand: various localities in BR, swampy paddocks  
in high-rainfall districts (NZAC).

—◎—

## Superfamily COPROMORPHOIDEA

(in the sense of Minet 1983)

### Family COPROMORPHIDAE



#### Copromorphidae

(166) *Isonomeutis amauropa* Meyrick

Note. Philpott (1928b, pp. 371–374) describes and illustrates morphological characters of the two genera.

● ***Isonomeutis*** Meyrick, 1888d, p. 75. Type species *Isonomeutis amauropa* Meyrick, by original monotypy.

***amauropa*** Meyrick, 1888d, pp. 75–76 (*Isonomeutis*)  
[Mount Manaia], Whangarei ND, E. Meyrick; HT ♂  
unique (?; see note), BMNH.

Hudson 1928, p. 297, pl. xlvi fig. 22 and 23; 1939, pl. lix  
fig. 24, as *I. restincta*.

Note. In his Diary of Captures for 21 December 1885  
Meyrick notes "Ison. amauropa, 4 from tree trunk".

***restincta*** Meyrick, 1923, p. 166 (*Isonomeutis*)  
[Kaeo ND], "Auckland", G.V. Hudson; HT ♂ unique,  
abdomen in gelatin capsule, BMNH.  
Hudson 1928, p. 297, pl. xlvi fig. 13.

● ***Phycomorpha*** Meyrick, 1914a, p. 106. Type species *Phycomorpha metachrysa* Meyrick, 1914, by original designation.


***metachrysa*** Meyrick, 1914a, pp. 106–107  
(*Phycomorpha*)

Dunedin DN, [W.G. Howes] A. Philpott; LT ♂ here designated, labelled "Dunedin AP 1.11.09", "Phycomorpha metachrysa Meyr. 2/2 E. Meyrick det. in Meyrick Coll.", abdomen missing, BMNH.

Hudson 1928, p. 296, pl. xxvi fig. 43.

Note. No specimens in collections have any green colour remaining, in contrast to *P. bryophylla* Meyrick from Samoa.

## Family CARPOSINIDAE



**Carposinidae**  
(167) *Heterocrossa eriphylla* (Meyrick)

● **Campbellana** Salmon & Bradley, 1956, p. 68. Type species *Campbellana attenuata* Salmon & Bradley, by original designation.

Note. Salmon & Bradley originally placed this genus in Yponomeutidae, but it was later shown to be a carposinid (Dugdale 1971b, pp. 73–75).

**attenuata** Salmon & Bradley, 1956, p. 69  
(*Campbellana*)

Campbell Island, J.H. Sorensen; HT ♂ designated by Salmon & Bradley, NMNZ.

● **Glaphyrarcha** Meyrick, 1938, pp. 428–429. Type species *Glaphyrarcha euthrepta* Meyrick, by original monotypy.

**euthrepta** Meyrick, 1938, p. 429 (*Glaphyrarcha*)  
Arthur's Pass NC/WD, "Mr Scott"; HT ♀ unique, CMNZ.  
Hudson 1939, p. 455, pl. lxi fig. 19.

● **Heterocrossa** Meyrick, 1882b, p. 178. Type species *Heterocrossa rubophaga* Dugdale (q.v.).

Note. Zimmerman (1978, p. 797) removed *Heterocrossa* from synonymy with *Carposina* Herrich-Schaeffer.

**adreptella** Walker, 1864a, p. 654 (*Gelechia*)  
[Auckland AK], D. Bolton; HT ♂ unique, BM genitalia slide no. 1852 ♂, head missing, BMNH.

Hudson 1928, p. 217, as *Carposina adreptella*.

**charaxias** Meyrick, 1981, p. 98 (*Heterocrossa*).  
New synonymy.

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Wellington New Zealand GVH /90", "Carposina charaxias Meyr. 3/11 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 217, pl. xxii fig. 20, as *Carposina charaxias*.

Note. Meyrick misinterpreted Walker's type material (see *H. rubophaga*, below). The type specimens of *adreptella* and *charaxias* are the only ones with broad, yellowish scales on the hindwing anal area, and the ♂ genitalia of *charaxias* – figured by Philpott (1928f, p. 478, fig. 4) from a Wellington WN specimen sent by Hudson (no. 114p) – agree well with the BMNH genitalia preparation from the *adreptella* HT (cf. *adreptella* of authors – Philpott 1928f, p. 480, fig. 9).

**canescens** Philpott, 1930c, p. 437 (*Carposina*)  
[Governor's Bush], Mount Cook MK, A. Philpott; HT ♂  
designated by Philpott, CMNZ.  
Hudson 1939, p. 455, pl. lx fig. 4.

**contactella** Walker, 1866b, pp. 1813–1814 (*Tinea*)  
[Nelson NN], T.R. Oxley, 1860; LT ♀ [HT] here designated,  
labelled "Tinea contactella" (cut-out strip from  
Walker's publication), "Tinea contactella Wkr TYPE  
♂", "Auckland N. Zeal. 60.73" (circular), BMNH.  
Hudson 1928, p. 215, possibly pl. xxii fig. 21.

**amatodes** Meyrick, 1911b, pp. 61–62 (*Carposina*).  
New synonymy.

Otira River WD, G.V. Hudson; LT ♀ here designated, labelled "Otira New Zealand GVH 12.08", "Carposina amatodes Meyr. 1/3 E. Meyrick det. in Meyrick Coll.", abdomen missing, BMNH.

Hudson 1928, p. 216, not figured.

Note. The forewing patterns of the two type specimens are almost identical, *amatodes* having more pronounced yellow patches. The PLT ♀ also lacks the abdomen.

**cryodana** Meyrick, §1885b, p. 349; 1885g, pp. 148–149 (*Heterocrossa*)

Dunedin DN, E. Meyrick; LT ♂ here designated, labelled "Dunedin New Zealand 27/9/82", "Carposina cryodana Meyr. 2/3 E. Meyrick det. in Meyrick Coll.", BMNH.

Hudson 1928, p. 217, pl. xxii fig. 19 (LT browner, and lacking the median stripe).

Note. Meyrick's Diary of Captures does not mention 27 September 1882, but notes for 28 September "H. cryodana 2".

**epomiana** Meyrick, §1885b, p. 349; 1885g, p. 149  
(*Heterocrossa*)

Otira [Gorge, 1,600 ft] WD, E. Meyrick; HT ♀ unique,  
BM genitalia slide no. 3693 ♀, BMNH.

Hudson 1928, p. 217, as synonym of *Carposina gonosemana*.

Note. See also *philpotti hudsoni*, below.

**eriphylla** Meyrick, 1888d, p. 76 (*Heterocrossa*)

Wellington WN, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 217, pl. xxiv fig. 52.

**exochana** Meyrick, 1888d, p. 76 (*Heterocrossa*)

Nelson NN, E. Meyrick; HT ♀ unique, BMNH.  
Hudson 1928, p. 217, pl. xxiv fig. 7.

**gonosemana** Meyrick, 1882b, pp. 179–180  
(*Heterocrossa*)

Dunedin DN, R.W. Fereday; LT ♂ here designated, labelled “Dunedin New Zealand RWF 21.2.79”, “*Carposina gonosemana* Meyr. 4/7 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, pp. 217–218 (in part), pl. xxii fig. 23 doubtful, as *Carposina gonosemana*.

Note. The ♂ collected by Fereday is the only remaining ST.

**ignobilis** Philpott, 1930c, p. 438 (*Carposina*) new combination

[Governor's Bush], Mount Cook MK, A. Philpott; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 455, not figured, as *Carposina ignobilis*.

**iophaea** Meyrick, 1907c, p. 117 (*Heterocrossa*)

Invercargill SL, A. Philpott; LT ♀ here designated, labelled “Invercargill New Zealand AP. 06”, “*Carposina iophaea* Meyr. 1/4 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 216, pl. xxii fig. 24, as *Carposina iophaea*.

**thalamota** Meyrick, 1909a, p. 12 (*Heterocrossa*). New synonymy.

Invercargill SL, A. Philpott; LT ♀ here designated, labelled “Invercargill New Zealand AP 1.07”, “*Carposina thalamota* Meyr. 1/5 E. Meyrick det. in Meyrick Coll.”, BMNH. Hudson 1928, p. 216, pl. xxii fig. 25 and 26, as *Carposina thalamota*.

Note. No consistent external differences were observed between the 2 type specimens, nor between the specimens in Meyrick's or Philpott's collections, nor the NZAC series from the type locality.

**literata** Philpott, 1930b, p. 11 (*Carposina*) new combination

Defiance Hut, [Mount Moltke], Franz Josef Glacier WD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ. Hudson 1939, p. 454, pl. lix fig. 22.

**maculosa** Philpott, 1927a, pp. 705–706 (*Carposina*) new combination

Cooper's Knob, Banks Peninsula MC, S. Lindsay; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 218, pl. xvii fig. 28.

**morbida** Meyrick, 1912c, p. 120 (*Carposina*) new combination

Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 218, pl. xxiv fig. 20.

**philpotti philpotti** Dugdale, 1971b, p. 75, fig. 9–11 (*Carposina*, as subspecies of *epomiana*) new combination

Camp Cove, Carnley Harbour, Auckland Islands, G.V. Hudson; HT ♂ designated by Dugdale, NMNZ.

Hudson 1909, p. 74, pl. ii fig. 17, as *Carposina gonosemana*; 1928, p. 217 (in part).

“n.sp.” Philpott, 1928f, pp. 477 (key) and 479, fig. 7

**philpotti hudsoni** new name proposed here for *epomiana* of authors but not Meyrick (1885, p. 149) (*Carposina*)

West Plains, [Invercargill] SL, A. Philpott; HT ♂ designated by Dugdale, NZAC.

**epomiana** not of Meyrick, but in the sense of Philpott (1928f, pp. 477 (key) and 479, fig. 6) (*Carposina*)

**rubophaga** new species proposed here for *adreptella* of authors, but not Walker (1864a, p. 564)

[Riccarton Bush], Christchurch MC, E. Meyrick; HT ♂ selected by K.R. Tuck, labelled “Christchurch New Zealand 9/3/82”, “*Carposina adreptella* Wlkr 13/18 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1928, p. 216, pl. xxii fig. 22; Philpott 1928f, p. 480, fig. 9, as *Carposina adreptella*.

**adreptella** Meyrick, 1882b, p. 179 (*Heterocrossa*; as *abreptella*), and Meyrick 1883b, p. 66.

Note. This species is the ‘raspberry budmoth’ of horticulture. The HT is drawn from the series collected by Meyrick at Riccarton Bush MC, because it was on this series that Meyrick based his concept of *adreptella*, and therefore of *Heterocrossa* (Meyrick 1882b, p. 179).

**sanctimonea** Clark, 1926, p. 418 (*Carposina*) new combination

Arthur's Pass NC/WD, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.

Hudson 1928, p. 218, pl. lii fig. 29, as *Carposina sanctimonea*.

**sarcanthes** Meyrick, 1918a, p. 133 (*Carposina*) new combination

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 216, not figured, as *Carposina sarcanthes*.

Also 3 undescribed species (NZAC).

● **Paramorpha** Meyrick, 1881b, pp. 696–697. Type species *Paramorpha aquilana* Meyrick, 1881b, pp. 697–698, by subsequent designation (Meyrick 1910c, p. 154); Australia.

**marginata** Philpott, 1931, p. 33 (*Carposina*) Okoroire WO, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 455, pl. lix fig. 23.

**heptacentra** Meyrick, 1931a, p. 95 (*Paramorpha*). Synonymised by Hudson (1939, p. 455). Whangarei ND, S.C. Patterson; HT not found in BMNH.


● Also 1 undescribed genus and species.

**Superfamily EPERMENIOIDEA**

(Minet, 1983)

**Family EPERMENIIDAE**

(in the sense of Gaedike 1978)



**Epermeniidae**

(168) *Thambotricha vates* Meyrick

● ***Thambotricha*** Meyrick, 1922b, p. 270. Type species *Thambotricha vates* Meyrick, by original monotypy.


***vates*** Meyrick, 1922b, pp. 270–271 (*Thambotricha*) Wellington WN, C.E. Clarke; HT ♂ unique, BMNH. Hudson 1928, p. 325, pl. I fig. 11.  
Note. Meyrick (1924a, pp. 204–205) redescribed the genus and species.

—◎—

**Superfamily PTEROPHOROIDEA**

**Family PTEROPHORIDAE**

(in the sense of Common 1970, p. 838)



**Pterophoridae**

(169) *Stenoptilia lithoxesta* Meyrick

● ***Lantanophaga*** Zimmerman, 1958, p. 400. Type species *Oxyptilus pusillidactylus* Walker, by original designation.

***pusillidactyla*** Walker, 1864b, p. 933 (*Oxyptilus*) West Indies, Jamaica, Mr Goose; HT ♀ in BMNH. Neotropical, “now widely spread in the tropics and subtropics” (Zimmerman 1958, p. 402) for control of *Lantana camara*, a weed shrub.  
New Zealand: NZAC has specimens collected in 1982 (adult) and 1986 (all stages) from Henderson and Mt Albert AK.

● ***Pterophorus*** Schaeffer, 1766, pl. 104 fig. 2 and 3. Type species *Phalaena Alucita pentadactyla* Linnaeus, 1758, p. 542, by subsequent designation (Whalley 1961, p. 159).

Note. See also ICBN 1964, p. 113.

***Aciptilia*** of authors, in the sense of Meyrick (1913b, p. 47, as *Alucita* Linn.). Type species *Phalaena Alucita pentadactyla* Linnaeus, 1758, p. 542, as given by Meyrick (1913b, p. 47).

***furcatalis*** Walker, 1864b, p. 950 (*Aciptilus*) [Nelson NN], T.R. Oxley; LT ♂ here designated, labelled “*furcatalis* Type Coll. Cab. 14 Dr. 5”, “26. *Aciptilus furcatalis*” (printed strip cut from Walker’s proof), “*Aciptilus furcatalis* Wkr Type”, “Auckland N. Zeal. 60-73” (circular), abdomen, forelegs, middle legs, and right hindleg missing, BMNH.  
Hudson 1928, p. 210, pl. xxiii fig. 17, as *Alucita furcatalis*, following Meyrick (1913b, p. 48).

Note. The 4 specimens from Auckland AK collected by Bolton (nos. a-d) could not be found.

**innotatalis** Walker, 1864b, p. 945 (*Pterophorus*)  
[Nelson NN]. T.R. Oxley; LT ♂ here designated, labelled  
“57. *Pterophorus innotatalis*” (printed strip cut from  
Walker’s proof), “Type Coll. Cab. 14 Dr. 6”, “*Pter-*  
*ophorus innotatalis* Wkr Type ♂”, “Auckland N. Zeal.  
60-73”, abdomen missing, BMNH.

Hudson 1928, p. 210, pl. xxiii fig. 1, as *Alucita innotatalis*,  
following Meyrick (1913b, p. 48).

**monospilalis** Walker, 1864b, pp. 950-951 (*Aciptilus*)  
[Auckland AK], D. Bolton; LT ♂ here designated, labelled  
“27. *Aciptilus monospilalis*” (printed strip cut from  
Walker’s proof), “*Aciptilus monospilalis* Wkr Type ♂”,  
“New Zealand 54.4” (circular), BMNH.

Hudson 1928, p. 209, pl. xxiii fig. 5 and 6, as *Alucita monospilalis*, following Meyrick (1913b, p. 47).

**furcatalis** var. β Walker, 1864b, p. 950 (*Aciptilus*). **New synonymy.**

[Nelson NN], T.R. Oxley; HT ♂ unique, labelled “Walk-  
er’s *furcatalis* var. β” [T.B. Fletcher’s writing], BMNH.

**furcatalis** var. γ Walker, 1864b, p. 950 (*Aciptilus*). **New synonymy.**

[Auckland AK], D. Bolton; LT ♂ (designated by Fletcher  
as “Type”) with label “This spec. was marked Type but  
is evidently Walker’s type γ. I have therefore shifted the  
type label off it. T.B.F.”, BMNH.

**patruelis** Felder & Rogenhofer, 1875, pl. cxl fig.  
56 (*Aciptilia*). **Synonymised by Meyrick (1885a, p.  
347; 1885f, p. 124).**

[Nelson NN, T.R. Oxley]; HT ♂ designated by Felder,  
BMNH.

Hudson 1928, p. 209, as synonym.

**lycosema** Meyrick, §1885a, p. 347; 1885f, pp. 123  
(key) and 124 (*Aciptilia*). **Synonymised by Chappell  
(1934, p. 163).**

[Botanic Gardens], Wellington WN, E. Meyrick; LT ♂  
here designated, labelled “Wellington New Zealand  
9/1/80”, “Lectotype JSD 1980”, BMNH.

Hudson 1928, p. 209, pl. xxiii fig. 18, as species.

Note. Philpott (1928h, fig. 11 and 12, and key, p. 649)  
illustrates genital structure differences between *monos-*  
*pilalis* and *lycosema*.

● **Platyptilia** of authors, in the sense of Meyrick  
(1910d, p. 9; 1913b, p. 47; 1913d, p. 10)

**aelodes** Meyrick, 1902c, p. 278 (*Platyptilia*)

Chatham Island, J. Fourgère; LT ♂ here designated,  
labelled “Chatham Is F. /00”, wingspan 15.5 mm,  
BMNH.

Hudson 1928, p. 207, as species.

**campsiptera** Meyrick, 1907c, pp. 112-113  
(*Platyptilia*)

Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 208, pl. xxiii fig. 15.

**carduidactyla** Riley, 1869, p. 180, pl. II fig. 13 and  
14 (*Pterophorus*)

North America.

New Zealand: adventive; in imported artichoke crowns,  
1973, Auckland (AK). Not known to be established.

Note. Lange (1950, pp. 585-592) gives a useful account  
of biology and illustrates structure.

**deprivatalis** Walker, 1864b, p. 946 (*Pterophorus*)

[Auckland AK], D. Bolton; LT ♂ here designated, labelled  
“58. *Pterophorus deprivatalis*” (printed slip cut from  
Walker’s proof), “*Pterophorus deprivatalis* Wkr Type  
♂”, “New Zeal. 54.4” (circular), BMNH.

Hudson 1928, p. 208, pl. xxiii fig. 2.

Note. Examination of HT *haasti* Felder shows it to be  
not synonymous with *deprivatalis*.

**falcatalis** Walker, 1864b, p. 931 (*Platyptilus*)

[Nelson NN], T.R. Oxley; LT ♂ here designated, labelled  
“15. *Platyptilus falcatalis*” (printed slip cut from  
Walker’s proof), “Type” (oblong card), “Auckland N.  
Zeal. 60-73” (circular), BMNH.

Hudson 1928, p. 207, pl. xxiii fig. 7 and 8.

**haasti** Felder & Rogenhofer, 1875, pl. cxl fig. 58  
(*Platyptilia*). **New synonymy.**

[Nelson NN, T.R. Oxley]; HT ♂ [unique], BMNH.  
Hudson 1928, p. 208, as synonym of *Platyptilia depri-*  
*valis*, following Meyrick (1913b, p. 47).

**indubitata** Philpott, 1928g, p. 485 (*Platyptilia*),  
new name for *Platyptilia ferruginea* Philpott, 1923,  
p. 150, preoccupied by *Crocydoscelus ferrugineum*  
(Walsingham, 1897, p. 35). **Synonymised by Hud-**  
**son (1928, p. 207).**

Mount Arthur Tableland NN, A. Philpott; HT ♀ desig-  
nated by Philpott, abdomen missing, NZAC.

Hudson 1928, p. 207, as *ferruginea* Philpott, as synonym;  
1939, p. 429 (refers to Philpott 1928g).

**heliastis** Meyrick, §1885a, p. 347; 1885f, pp. 127  
(key) and 129 (*Platyptilia*)

Porter’s Pass MC, R.W. Fereday; HT ♂ unique, BMNH.  
Hudson 1928, p. 208, pl. xxiii fig. 13.

Note. The published locality is “Castle Hill”, the sheep  
station then owned by the published collector, J.D.  
Enys. Porter’s Pass was on Enys’s run, Fereday and  
Enys collected together, and Meyrick’s labelling was  
occasionally inaccurate.

**hokowhitalis** Hudson, 1939, p. 430 (*Platyptilia*)

Hokowhitu Bush WN/WI, G.V. Hudson; HT ♂ unique,  
labelled “504a”, NMNZ.

Hudson 1928, pl. xxiii fig. 3, as *Platyptilia celidota* (in  
error); 1939, p. 430.

**isoterma** Meyrick, 1909a, p. 10 (*Platyptilia*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 207, as synonym of *Platyptilia falcatalis*.

Note. Palpal length and colour pattern distinguish HT  
*isoterma* from HT *falcatalis* and HT *haasti*, so *iso-*  
*terma* is here removed from synonymy with *falcatalis*.

» Pterophoridae, *Platyptilia*

**pulverulenta** Philpott, 1923, p. 149 (*Platyptilia*)  
Nelson NN, A. Philpott; HT ♂ designated by Philpott,  
NZAC.

Hudson 1928, p. 207, as synonym of *Platyptilia falcatalis*.  
Note. Palpal length and colour pattern clearly distinguish  
HT *pulverulenta* from HT *falcatalis*, *haasti*, and  
*indubitata*, so *pulverulenta* is here removed from syn-  
onymy with *falcatalis*.

**repletalis** Walker, 1864b, p. 931 (*Platyptilia*)  
[Auckland AK], D. Bolton; LT ♂ (as HT) here desig-  
nated, labelled "Type Coll. Cab. 14 drawer 5", "TYPE",  
"16. *Platyptilia repletalis*" (printed slip cut from  
Walker's proof), "New Zealand 54.4" (circular), abdo-  
men missing, BMNH.

Hudson 1928, p. 207, as synonym of *Platyptilia falcatalis*;  
1928, p. 207, pl. xxiii fig. 14, as *Platyptilia aelodes*  
(part).

Note. Dugdale (1971b, pp. 147–148) discusses differences  
between *repletalis*, *aelodes*, and *falcatalis*.

Also 1 undescribed species (NZAC).

● ***Stenoptilia*** in the sense of Meyrick (1910d, p. 18;  
1913b, p. 48; 1913d, p. 28)

**charadrias** Meyrick, §1885a, p. 347; 1885f, p. 126  
(*Mimaeseoptilus*)

[Otira Gorge WD], E. Meyrick; LT ♂ here designated,  
labelled "Otira Gorge New Zealand 1500 ft 29/1/83",  
"Lectotype", BMNH.

Hudson 1928, p. 211, pl. xxiii fig. 4.

**vigens** not of Felder & Rogenhofer, 1875, pl. cxl  
fig. 49, but in the sense of Meyrick (1912c, p. 119)  
(*Stenoptilia*).

Hudson 1928, p. 211, pl. xxiii fig. 9, as species.  
Note. Known from specimens from Humboldt Range OL  
collected by Hudson and sent to Meyrick. Felder &  
Rogenhofer's *Oxyptilus vigens*, although recorded as from  
"Nova Seelandia", is actually from Natal, Republic of  
South Africa.

**celidota** Meyrick, §1885a, p. 347; 1885f, p. 125  
(*Lioptilus*)

Christchurch MC, R.W. Fereday; LT ♂ here designated,  
labelled "Lectotype JSD 1980" (circular), "Christ-  
church New Zealand RWF /82", BMNH.

Hudson 1928, p. 209, pl. xxiii fig. 3, as *Platyptilia celidota*.

Note. None of the type series has black scales on the  
forewing dorsum (cf. Hudson 1928, p. 209), and the  
species is retained in *Stenoptilia* (see Meyrick 1913b,  
p. 48).

**epotis** Meyrick, 1905, pp. 231–232 (*Platyptilia*)  
Humboldt Range OL, G.V. Hudson; HT ♀ unique,  
BMNH.

Hudson 1928, p. 210, pl. xxiii fig. 16.

***lithoxesta*** Meyrick, §1885a, p. 347; 1885f, pp. 126  
(key) and 127 (*Mimaeseoptilus*)

Arthur's Pass NC/WD, E. Meyrick; LT ♂ here desig-  
nated, labelled "Lectotype JSD 1980" (circular),  
"Arthur's Pass New Zealand 3000 ft 25/1/83", BMNH.  
Hudson 1928, p. 210, pl. xxiii fig. 11.

**orites** Meyrick, §1885a, p. 347; 1885f, p. 127  
(*Mimaeseoptilus*)

Clinton SL, G.F. Mathew; HT ♂ unique, BMNH.  
Hudson 1928, p. 211, pl. xxiii fig. 10.

Note. HT bears the label "Dunedin New Zealand Mathew  
1887", but it is likely that Meyrick saw Mathew's col-  
lection in New Zealand before it was received at  
BMNH in 1887 and given the accession number "87-  
50".

***zophodactyla*** Duponchel, 1838, pl. 314 fig. 4  
(*Pterophorus*)

Described from France; HT not examined, ?MNHN.


***canalis*** Walker, 1864b, p. 944 (*Pterophorus*).  
Synonymised by Meyrick (1910d, p. 19).

Sydney N.S.W., Lambert; ST "a" in BMNH.  
Hudson 1928, p. 211, pl. xxiii fig. 12.

Note. There is no evidence that *S. zophodactyla* was  
introduced by man.

—♂—

Superfamily PAPILIONOIDEA  
Family LYCAENIDAE




---

**Lycaenidae**  
(170) *Lycaena salustius* (Fabricius)

---

**Subfamily LYCAENINAE**  
(in the sense of Common & Waterhouse  
1972, p. 308)

• **Lampides** Hübner, [1819], p. 70. Type species *Papilio (Plebejus Rurales) boeticus* Linnaeus, by subsequent designation (Grote 1873a, p. 179).

**boeticus** Linnaeus, 1767, p. 789 (*Papilio P. R.*)  
Type locality given as "Barbaria".  
New Zealand: self-adventive since 1965 (Gibbs 1980, p. 176); ND-NN, locally common.

• **Lycaena** Fabricius, 1807, p. 285. Type species *Papilio phlaeas* Linnaeus, 1776, p. 285, by subsequent designation (Curtis 1828, p. 5, pl. 12).

**Helleia** in the sense of Sibatini (1974, p. 109) but not of Verity (1943). Synonymised by Gibbs (1980, p. 106).

Note. Sibatini (1974, p. 109) gives "A tentative scheme of higher classification of Lycaeninae (s.str.) of the world", placing the New Zealand species *salustius* and *feredayi* in a subgenus of *Helleia* Verity, and *boldenarum* in another (unnamed) genus. Miller & Brown (1979, pp. 5 and 6) refer to "Lycaena" *salustius*.

**boldenarum boldenarum** White, 1862, p. xxvi  
(*Lycaena*)

[HB or TO], W. Colenso; HT (?gender), BMNH.  
Hudson 1898, pp. 118-119, pl. xii fig. 13-17; 1928, pp. 38-39, pl. v fig. 1-6 and 14-17, as *Chrysophanus boldenarum*, after Butler (1874, p. 29). Gibbs 1980b, p. 155, as *Lycaena boldenarum*.

**boldenarum caerulea** Salmon, 1946, pp. 10-11  
(*Chrysophanus*), as subspecies of *boldenarum* (*sic*).

Hollyford Valley FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.  
Salmon 1946, pp. 10-11, pl. 1 fig. 12 and 13. Gibbs (1980b, p. 156) notes Salmon's action, but ignores it nomenclaturally.

**boldenarum ianthina** Salmon, 1946, p. 11 (*Chrysophanus*), as subspecies of *boldenarum* (*sic*).

Milford Sound FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.  
Salmon 1946, p. 11, pl. 1 fig. 7 and 8. Gibbs (1980b, p. 156) notes Salmon's action, but ignores it nomenclaturally.

**feredayi** Bates, 1867, p. 53 (*Chrysophanus*)

Kaiapoi Bush MC, R.W. Fereday; ST ♂ so labelled by P. Ackery (1985), BMNH.

Hudson 1898, p. 116; 1928, p. 37, as possible synonym of *Chrysophanus salustius*. Gibbs 1980a, p. 105, fig. 7-9, 19-21, and 31, as *Lycaena feredayi*.

**enysii** Butler, 1876, p. 153 (*Chrysophanus*). Synonymised by Hutton (1901, p. 97, as *enysii*).

[?Hawkes Bay HB], J.D. Enys; HT ♂ labelled by Butler as "Type", BMNH.  
Hudson 1898, pp. 117-118, pl. xii fig. 22-24; 1928, p. 38, pl. v fig. 10-12; as *Chrysophanus enysii*. Gibbs 1980a, p. 105; 1980b, p. 150; as synonym of *Lycaena feredayi*.

**rauparaha** Fereday, 1877b, p. 462 (*Chrysophanus*)

Kaiapoi Bush MC, R.W. Fereday; HT ♂ so labelled, CMNZ.

Hudson 1898, p. 116, pl. xii fig. 21, pl. 13 fig. 2; 1928, p. 36, pl. v fig. 24 and 25, as synonym of *Chrysophanus salustius*. Gibbs 1980a, pp. 110-113, fig. 4-6, 10, 13-15, 22, and 26-29, as *Lycaena rauparaha*.

Note. Hutton (1901, p. 97) was first to uphold the distinctiveness of *rauparaha* (see Gibbs 1980a, pp. 105-106). Fereday (1878a, p. 255) gave additional details. See note under *salustius*, below.

**salustius** Fabricius, 1793, p. 310 (*Hesperia Rurales*)  
"Habitat in India", Drury; HT not found, depicted in *Jones Icones* 6, tab. 59 fig. 1.

Hudson 1898, pp. 116-117, pl. xiii fig. 3-5, pl. xviii fig. 18-20; 1928, pp. 36-38, pl. v fig. 7-9, 26, and 27, as *Chrysophanus salustius*. Gibbs 1980a, pp. 106-109, fig. 1-3, 16-18, 24, and 30, as *Lycaena salustius*.

Note. As outlined by Andrews (1986), there is strong evidence – based on the *Jones Icones* – to suggest that *salustius* Fabricius = *rauparaha* Fereday. The matter will not be resolved until Drury's specimen, thought to be in the Macleay Museum, Sydney, is located and examined.

**edna** Doubleday, 1843, p. 283 (*Lycaena*). Synonymised by Butler (1869, p. 174).

"New Zealand", E. Dieffenbach; HT not located in BMNH (P. Ackery, pers. comm.).

Hudson 1898, p. 116; 1928, p. 36; as synonym.

## Subfamily DANAINAE

**maui** Fereday, 1877b, pp. 461–462 (*Chrysophanus*). Synonymised by Hudson (1898, p. 116). Wellington WN, R.W. Fereday; HT ♂ designated as “Type” by Fereday, no locality label, CMNZ.

Hudson 1898, p. 116; 1928, p. 36, pl. v fig. 27, as synonym. Gibbs 1980b, p. 142, as Hudson’s “Wellington inland form”. Fereday (1878a, pp. 254–255) gave additional details.

• **Zizina** Chapman, 1910, p. 482. Type species *Polyommatus labradus* Godart, by original designation.

**labradus labradus** Godart, 1824, p. 680 (*Polyommatus*; as species)

“?Australia”; type material in MNHN.

Hudson 1898, p. 119, pl. xii fig. 10 and 11, as *Lycaena phoebe* Murray. Stempffer (1967, p. 258), as *Zizina labradus*; Common & Waterhouse 1972, p. 409, and Gibbs 1980b, pp. 166–172, fig. 60a, as *Zizina otis labradus*.

Note. The name *Lycaena phoebe* Murray, 1873, pp. 107–108, refers to specimens collected in “South Australia” and Brisbane standing under that name in BMNH.

**labradus oxleyi** Felder & Felder, 1865, pp. 280–281, pl. xxxv fig. 6 (*Lycaena*)

[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.

Hudson 1883, pp. 119–120, pl. xii fig. 12, as species; 1928, p. 40, not figured, as synonym of *Lycaena labradus*. Gibbs 1980b, pp. 173–175, fig. 60b, as subspecies of *Zizina otis*.

—◎—

## Family NYMPHALIDAE

## Subfamily BRASSOLINAE

• **Opsiphanes** Doubleday, [1851], p. 344, pl. 57 fig. 2 [illustration]; Westwood in Doubleday & Westwood, 1851, p. 346 [text]. Type species *Opsiphanes sallaei* Doubleday, 1849, pl. 57 fig. 2, by subsequent designation (Scudder 1875, p. 233); Neotropical.

**cassina** Felder & Felder, 1862, p. 122 (*Opsiphanes*) New Zealand: adventive; DN (Harris 1982, p. 330, as *Opsiphanes cassina*).

**tamarindi** Felder & Felder, 1861, p. 111 (*Opsiphanes*)

New Zealand: adventive; WN, DN, SL (Harris 1982, pp. 329–330, fig. 1 and 2.)

Note. Both species have been intercepted alive in shipments of Ecuadorian bananas; neither has established.

• **Danaus** Kluk, 1802, p. 84. Type species *Papilio plexippus* Linnaeus, 1758, by subsequent designation (Hemming 1933, p. 222).

**plexippus** Linnaeus, 1758, p. 471 (*Papilio Danaus*) North America.

New Zealand: self-adventive; throughout, locally established ND-NN.

Hudson 1898, pp. 102–104, pl. xi fig. 1 and 2, as *Anosia erippus* Cramer; 1928, pp. 26–28, pl. iv fig. 10, as *Danaida plexippus*. Gibbs 1980b, pp. 62–67, as *Danaus plexippus plexippus*.

Note. Fereday (1874, p. 183) gave early New Zealand records, using the specific name *berenice* Cramer. The specimen he records as found by Mrs Meinertzhagen is in CMNZ.

**chrysippus petilia** Stoll, 1790, pl. 28 fig. 3 (*Papilio*; as species)

New Zealand: occasionally self-adventive (Gibbs 1980b, pp. 68–69), ND-WD.

Hudson 1928, p. 28, pl. iv fig. 11 and 12, as *Danaida chrysippus*. Gibbs 1980b, pp. 68–70, pl. 41.

• **Tirumala** Moore, 1880, p. 4. Type species *Papilio limniace* Cramer, 1775, by original designation.

**hamatus hamatus** Macleay, 1826, p. 451 (*Euploea*; as species)

Australia.

New Zealand: occasionally self-adventive, non-establishing, ND-WN (Gibbs 1980b, p. 70).

Hudson 1950, p. 66, pl. B fig. 6. Gibbs 1980b, pp. 70–71, pl. 44, as *Danaus hamata*.

—◎—

## Subfamily NYMPHALINAE

(in the sense of Common 1970, p. 843)


• **Bassaris** Hübner, [1821a], pl. 24. Type species *Papilio Nymphalis Gemmata itea* Fabricius, by monotypy (Hemming 1967, p. 74).

Note. Field (1971, p. 28) resurrected Hübner’s genus for Australian and New Zealand species previously placed in *Pyrameis* or *Vanessa* of authors.

**gonerilla gonella** Fabricius, 1775, p. 498 (*Papilio N. G.*, as species)

“Habitat in nova Zelandia”, J. Banks; HT ♂ unique, type no. Rh. 8918, BMNH.

Hudson 1898, pp. 105–107, pl. xii fig. 5 and 6; 1928, pp. 34–35, pl. iv fig. 2 and 9, as *Vanessa gonella*. Gibbs 1980b, pp. 111–115, pl. 108, as *Bassaris gonella gonella*, after Field (1971, p. 33).



**Nymphalidae: Nymphalinae**  
(171) *Bassaris gonerilla* (Fabricius)

**gonerilla ida** Alfken, 1899, pp. 5–8 (*Pyrameis*; as species)

Te One and Maunganui, Chatham Islands, “Schauinsland”; not found, ?Bremen Museum (Field 1971, p. 32).

Hudson 1928, p. 35, pl. v fig. 31, as “Chatham Island form” and “*Vanessa ida* Alfken” as footnote. Gibbs 1980b, pp. 116–118, as *Bassaris g. ida*, after Field (1971, pp. 32–33, fig. 87 and 88, taken from Alfken 1903, p. 602, pl. 32 fig. 12).

**argentata** Alfken, 1903, p. 602, pl. 32 fig. 13 (*Pyrameis ida* var.). Synonymised by Field (1971, p. 32). Excluded name, Type 3, in the sense of Field (1971, p. 4).

Chatham Islands, Schauinsland; type material not found, ?Bremen Museum (Field 1971, p. 32).

**itea** Fabricius, 1775, p. 498 (*Papilio N. G.*)

“Habitat in nova Zealandia”, J. Banks; HT ♂ unique, type no. Rh.8719, BMNH.

Hudson 1898, pp. 107–108, pl. xii fig. 3 and 4; 1928, pp. 35–36, pl. iv fig. 8, as *Vanessa itea*. Gibbs 1980b, pp. 118–121, pl. 109, as *Bassaris itea*, after Field (1971, p. 30).

● **Cynthia** Fabricius, 1807, p. 281. Type species *Papilio Nymphalis cardui* Linnaeus, 1758, p. 475, by subsequent designation (Westwood 1840, p. 87). Note. Field (1971, p. 33) resurrected Fabricius’s genus.

**kershawi** McCoy, 1868, p. 76 (*Cynthia*)

?Melbourne Vict., Kershaw; HT not found (Field 1971, p. 46).

New Zealand: regularly self-adventive throughout, temporarily establishing from ND to NN on adventive Asteraceae.

Hudson 1898, pp. 108–109, pl. xii fig. 1 and 2; 1928, p. 36, pl. iv fig. 1; as *Vanessa cardui*, with *kershawi* as

synonym. Gibbs 1980b, pp. 121–125, pl. 128, as *Cynthia kershawi*, after Field (1971, p. 45), as resurrected combination.

● **Hypolimnas** Hübner, [1819], p. 45. Type species *Papilio Nymphalis pipleis* Linnaeus, 1758, p. 476, by subsequent designation (Scudder 1875, p. 194).

**bolina nerina** Fabricius, 1775, p. 509 (*Papilio Nymphalis Phaleratus*; as species)

“Habitat in nova Hollandia”, J. Banks; HT ♀ unique, BMNH.

New Zealand: irregularly self-adventive but non-establishing, ND–WD/ND.

Hudson 1898, pp. 104–105, pl. xii fig. 7–9, as *Anosia bolina*; 1928, pp. 32–33, pl. v fig. 18 and 19, as *Hypolimnas b. nerina*, after Common & Waterhouse (1972, p. 287).

● **Junonia** Hübner, 1819, p. 34. Type species *Papilio lavinia* Cramer, 1775, p. 32, pl. 21 fig. C and D, by subsequent designation (Scudder 1872, p. 43) (Hemming 1967, p. 239).

Note. *Junonia*, an Indo-Malayan genus, is used here following de Lesse (1952, pp. 74–77, fig. 1). I have examined African *Junonia lavinia coenia* (Hübner) and the Nearctic–Neotropical *Precis octavia* (Cramer), and agree with de Lesse’s findings. Tilden (1971, pp. 101–108) regarded the two genera as synonymous, but all species examined by him are in *Junonia* in the sense of de Lesse. *Junonia* is adopted by Common & Waterhouse (1981, 2nd edn), following Edwards (1977).

**villida calybe** Godart, 1819, p. 317 (*Vanessa*)


Australia.

New Zealand: intermittently adventive, ND–WD–DN. Hudson 1898, pp. 109–110, pl. xi fig. 16 and 17, as *Junonia velleda*; 1928, pp. 33–34, pl. iv fig. 15, as *Precis velleda*; Gibbs 1980b, pp. 128–129, pl. 138, as *Precis villica calybe*, after Common & Waterhouse (1972, p. 294).

Note. The type locality of the nominate subspecies (*P. v. villica* Fabricius) is given as “Insula Amsterdam” (Fabricius 1787, p. 35), in the Macassar Straits. See note under Godart, J.B., 1819 in ‘References’. The species *calybe* may be a Latreille name.

—◎—

**Subfamily SATYRINAE**  
(in the sense of Common & Waterhouse  
1972, p. 232)




---

**Satyrinae**  
(172) *Argyrophenga antipodum* (Doubleday)

---

• ***Argyrophenga*** Doubleday, 1845, p. 307. Type species *Argyrophenga antipodum* Doubleday, by original monotypy.

***antipodum*** Doubleday, 1845, p. 307 (*Argyrophenga*)  
Waikouaiti DN, P. Earl; HT ♂ here designated, labelled  
“New Zealand, purch. fr. Earl 45.30”, “B.M. Type No.  
Rh. 3552”, “Type H.T.”, “Argyrophenga antipodum  
Doubleday det. P. Ackery 1976”, BMNH.  
Hudson 1898, pp. 110–112, pl. xi fig. 4–6; 1928, pp. 29–  
30, pl. iv fig. 18–20. Craw 1978, pp. 754–761, fig. 2,  
3, 6 (part), 7k–o, and 8k–r.

***harrisi*** Craw, 1978, pp. 761–762 (*Argyrophenga*)  
Mount Owen NN, J.S. Dugdale; HT ♂ designated by  
Craw, NZAC.  
Not mentioned by Hudson. Craw 1978, pp. 761–762, fig.  
4, 7f–j, and 8f–j.

***janitae*** Craw, 1978, pp. 763–765 (*Argyrophenga*)  
Lake Sylvester, Cobb Valley NN, J.S. Dugdale; HT ♂  
designated by Craw, NZAC.  
Hudson 1898, pp. 110–111, pl. xi fig. 3 and 7; 1928, p.  
29, pl. iv fig. 6 and 7; as *Argyrophenga antipodum*.  
Craw 1978, pp. 763–765, fig. 5, 6 (part), 7a–e, and 8a–  
e.

• ***Dodonidia*** Butler, 1884a, p. 172. Type species  
*Dodonidia helmsii* Butler, by original designation.

***helmsii*** Butler, 1884a, p. 172 (*Dodonidia*)  
Paparoa Range BR, R. Helms; HT ♂ unique, in pieces,  
CMNZ.  
Hudson 1898, pp. 112–113, pl. xi fig. 14 and 15; 1928, p.  
30, pl. iv fig. 16 and 17; as *Dodonidia helmsii* Fereday.  
Gibbs 1980b, pp. 77–82, pl. 47, as *Dodonidia helmsii*  
Butler.

***helmsii*** Fereday, 1883b, p. 193 (no genus); invalid  
name  
Type locality (as “Paparoa Range”), collector, and HT as  
above.  
Note. Hemming (1967, p. 148), following Butler (1884a,  
p. 172), wrongly ascribed the spelling “helmsii” to Fereday  
(1883b, p. 193). Butler’s authorship takes precedence  
over Fereday’s under the Rules, as does Butler’s spelling  
(*helmsii*), as this is the spelling used the first time the  
binomen (genus + species) was published.

• ***Erebiola*** Fereday, 1879, p. 128. Type species *Erebiola butleri* Fereday, by original designation.

Note. Warren (in Wise 1967, p. 40) gave *Dubierebia* Muschamp, 1915, pp. 12–26 (type species *Erebiola myops* Staudinger, 1881; Ala Tau, Turkestan)  
as a junior subjective synonym of *Erebiola*. Miller  
(1968) assigned each genus to a different tribe, and  
this action is currently recognised (R.C. Craw, pers.  
comm.).

***butleri*** Fereday, 1879, pp. 129–130, pl. 1 fig. 3 and  
4 (*Erebiola*)

Whitcombe Pass MC/WD, J.D. Enys; HT ♂ designated  
by Wise (1967, p. 4), CMNZ.  
Hudson 1898, p. 115, pl. xi fig. 11 and 12; 1928, pp. 31–  
32, pl. iv fig. 3–5. Gibbs 1980b, pp. 95–98, pl. 85–87.

• ***Melanitis*** Fabricius, 1807, p. 282. Type species  
*Papilio Nymphaalis leda* Linnaeus, 1758, p. 474, by  
subsequent designation (Butler 1868, p. 194);  
“Asia”.

***leda bankia*** Fabricius, 1775, p. 499 (*Papilio Nymphaalis Gemmata*; as species)  
“Australia”, J. Banks; HT (?gender) not examined,  
BMNH.  
New Zealand: self-adventive, sporadic, not establishing.  
Holloway 1962, pp. 79–82, fig. 1 and 2.

• ***Oreixenica*** Waterhouse & Lyell, 1914, p. 41. Type  
species *Lasiommata*(?) *lathoniella* Westwood, by  
original designation.

***lathoniella herceus*** Waterhouse & Lyell, 1914, p. 42  
(*Oreixenica*)  
South-eastern Australia. New Zealand: ?adventive; 1 ♂  
found in a gutter, Wellington City WN, 1972 by C.F.  
Mercer (NZAC).

• ***Perknodaimon*** Butler, 1876, p. 152. Type species  
*Erebia merula* Hewitson (as *Erebia pluto* Fereday  
MS.), by original monotypy.

Note. There is strong disagreement (cf. Hemming  
1967, p. 352, and Wise 1967, p. 41) as to whether  
Fereday’s brief mention (“I may also mention a  
black butterfly found on the bare summits of the

snowy mountains, and of which I have several specimens ... I believe it to be a species of *Erebia* and I have named it *E. pluto*") constitutes an indication according to the Rules of Zoological Nomenclature. The arrangement below is that stated (or implied) by Hemming, and cannot be regarded as final since neither Hemming's nor Wise's points have been put before the International Commission for Zoological Nomenclature.

**merula** Hewitson, 1875, p. 10 (*Erebia*); first valid name (Hemming 1967, p. 352)

"New Zealand" [description equates with Canterbury specimens], H. Strecker; HT (?gender), BMNH.

Hudson 1898, pp. 114–115, pl. xi fig. 8–10; 1928, p. 31, pl. v fig. 13, 29, and 30; as *Erebia pluto* Fereday.

Note. Philpott (1928g, p. 481) refers to it as *Erebia merula*, and gives history. Gibbs (1980b, pp. 99–105) refers to it as *Percnodaimon pluto* (Fereday MS.) Butler.

**pluto** Fereday, 1872, p. 217 (*Erebia*); invalid name (Hemming 1967, p. 352)

Craigieburn Range MC, J.D. Enys; LT ♂ designated by Wise (1967, p. 42), CMNZ.

Note. Fereday (1872, p. 217) records that the specimens were collected at altitudes over 6000 feet. This combination was upheld as valid by Wise (1967, p. 41).

**othello** Fereday, 1876, pp. 302–304, pl. ix (*Oreina?*); replacement name for *pluto*, preoccupied by *Erebia pluto* (de Prunner, 1798). Synonymised by Butler (1878, p. 268).

Type data as for *Erebia pluto* Fereday.

**pluto** (Fereday MS.) Butler, 1876, p. 153 (*Percnodaimon*). Synonymised by Hemming (1967, p. 352).

Type data as for *Erebia pluto* Fereday.

Note. Butler here 'formalised' Fereday's (1872) indication.

—◎—

**rapae rapae** Linnaeus, 1758, p. 468 (*Papilio Danaus Candidus*; as species)

Europe.

New Zealand: man-adventive; recorded since 1930, ND-SL, throughout.

Hudson 1939, pp. 389–391, pl. lxii fig. 7 and 27, as *Pieris rapae*. Gibbs 1980b, pp. 44–56, pl. 8, as *Pieris rapae*.

Note. Kudrna (1974) and Higgins (1975) revived the genus *Artogeia* Verity, 1947, pp. 192–193 (type species by original designation *Papilio napi* Linnaeus, 1758), to reflect differences in male genitalia, androconial scales, and chromosome numbers between the crucifer-eating, Palearctic species *rapae* and *brassicae*. The citation given is that adopted by Gibbs (1980b).

—◎—

## Family PIERIDAE

• **Catopsilia** Hübner, [1819], p. 98. Type species *Papilio crocale* Cramer, [1775], by subsequent designation (Scudder, 1871, p. 58).

**pomona pomona** Fabricius, 1775, p. 479 (*Papilio Danaus Candidus*; as species)

"Asia".

New Zealand: recorded once, in Auckland AK.  
Hudson 1898, p. 121, as *Catopsilia catilla* Cramer.

• **Pieris** Schrank, 1801, pp. 152 and 161. Type species *Papilio brassicae* Linnaeus, 1758, by subsequent designation (Latreille 1810, pp. 440 and 351).

**Superfamily BOMBYCOIDEA**  
**Family BOMBYCIDAE**

• **Bombyx** Linnaeus, 1758, p. 496. Type species *Bombyx mori* Linnaeus, as in Kirby (1892, p. 718).

*mori* Linnaeus, 1758, p. 499 (*Bombyx*)

China.

Domesticated worldwide, including New Zealand.

—◎—

**Family SATURNIIDAE**

• **Antheraea** Hübner, [1819], p. 152. Type species *Attacus mylitta* Drury, 1773, by subsequent designation (Kirby 1892, p. 759).

Note. Australian species are now assigned to *Opodiphthera* Wallengren (E. D. Edwards, pers. comm.).

*eucalypti* Scott, 1864, pl. 1 (*Antheraea*)

Eastern Australia.

New Zealand: introduced by man, now established on *Eucalyptus* and *Schinus*, generally north of 39°30'S and coastally south to 41°30'S.

• **Samia** Hübner, [1819], p. 156. Type species *Bombyx cecropia* Linnaeus, 1758, p. 447, by subsequent designation (Kirby 1892, p. 750); North America.

*cynthia* Drury, 1773, pl. 6 fig. 2 (*Attacus*)

Java.

New Zealand: man-adventive; established in Auckland City AK on *Ailanthis* trees around the university and Albert Park.

Note. The following species have been reported – but never for more than two consecutive seasons – from the Auckland metropolitan area. All species listed are available from overseas commercial Lepidoptera dealers.

- *Actias selene* (Hübner, 1819)
- *Antheraea pernyi* (Guérin-Méneville, 1855)
- *Dictyoplaca simla* (Westwood, 1848)
- *Hyalophora cecropia* (Linnaeus, 1758)

—◎—

**Family SPHINGIDAE**

Note. Only *Agrius convolvuli* is resident in New Zealand.

• **Agrius** Hübner, [1819], p. 140. Type species *Sphinx cingulata* Fabricius, 1775, by subsequent designation (Tutt 1902, p. 353); America.

*convolvuli* Linnaeus, 1758, p. 490 (*Sphinx*)

Old World, Pacific.

New Zealand: intermittently resident, on *Convolvulus*, *Calystegia*, and *Ipomoea*.

Hudson 1898, pp. 99–100, pl. xiii fig. 1; 1928, p. 41, pl. vi fig. 16; as *Sphinx convolvuli*, after Meyrick (1917b, p. 271).

*distans* Butler, 1874a, p. 30; 1874b, p. 4 (*Sphinx*, as variety of *convolvuli*, providing a name for “*S. convolvuli* var.  $\gamma$ ” of Walker 1856b, pp. 213–214). Synonymised by Meyrick (1890, p. 213).

[Auckland AK], D. Bolton; ST series BMNH. Hudson 1898, p. 99; 1928, p. 41; as *Protoparce distans*, as synonym.

• **Daphnis** Hübner, [1819], p. 134. Type species *Sphinx nerii* Linnaeus, by subsequent designation (Kirby 1892, p. 671).

*placida placida* Walker, 1856, p. 186 (*Darapsa*) Sumatra.

New Zealand: one record from Auckland AK; the specimen is in AMNZ.

• **Hippotion** Hübner, [1819], p. 134. Type species *Sphinx celerio* Linnaeus, 1758; “Old World”.

*celerio* Linnaeus, 1758, p. 491 (*Sphinx*)

Old World, Pacific.


New Zealand: vagrant, sporadic, non-establishing.

Hudson 1928, p. 42, pl. vi fig. 15, as *Deilephila celerio*, after Meyrick (1917b, p. 271).

Note. Buller (1905, p. 332) and Hudson (1905b, p. 359) give earliest records.

—◎—

**Superfamily THYRIDOIDEA**  
 (in the sense of Minet 1983, p. 203)  
**Family THYRIDIDAE**



**Thyrididae**  
 (173) *Morova subfasciata* Walker

**Subfamily SICULINAE**  
 (in the sense of Whalley 1971, p. 16)

- ***Morova*** Walker, 1865a, p. 523. Type species *Morova subfasciata* Walker, by original monotypy.

***subfasciata*** Walker, 1865a, pp. 523–524 (*Morova*)  
 [Nelson NN], T.R. Oxley; HT ♀ unique, head missing,  
 rest of body glued together, BMNH.

Hudson 1928, p. 206, pl. xxiv fig. 25 and 26.

***gallicolens*** Butler, 1874, p. 46 (*Cacoecia*). Synonymised by Meyrick (§1883d, p. 531; 1884b, p. 108; 1913b, p. 46).

Christchurch MC, C.M. Wakefield; HT ♂ designated by Butler, BMNH.

Hudson 1928, p. 206, as synonym.

Note. *C. gallicolens* HT bears the accession label 74.13, registered as “Christchurch, C.M. Wakefield, sent to Albert Müller, type material of *C. gallicolens* Butler”. Accession label 76.57 refers to additional material sent by Wakefield. The record by Meyrick (1886g, p. 216) of this species from Fiji is based on a misidentification.

—◎—

**Superfamily PYRALOIDEA**  
**Family PYRALIDAE**  
 (in the sense of Minet 1981, p. 267)

**Subfamily GALLERIINAE**

- ***Achroia*** Hübner, [1819], p. 163. Type species *Bombyx cinereola* Hübner, [1803], pl. 23 fig. 91, by subsequent designation (Ragonot 1893, p. x, as *cinerola*).

***grisella*** Fabricius, 1794, p. 289 (*Tinea*)  
 Cosmopolitan.  
 New Zealand: adventive; in beehives.  
 Hudson 1928, p. 157, pl. xix fig. 21, as *Meliphora grisella*.

- ***Galleria*** Fabricius, 1798, p. 419. Type species *Tinea cereana* Blom, 1764, by subsequent designation (Latreille 1810, p. 441, as *cereana* Fab.).

***mellonella*** Linnaeus, 1758, p. 888 (*Tinea*)  
 Cosmopolitan.  
 New Zealand: adventive; in beehives.  
 Hudson 1928, p. 158, pl. li fig. 9.

—◎—

**Subfamily PHYCITINAE**  
 (in the sense of Munroe 1972, pp. 12–13)

- ***Crocydopora*** Meyrick, 1882b, pp. 158–159. Type species *Nephopteryx stenopterella* Meyrick, 1879, by original monotypy.

***cinigerella*** Walker, 1866b, p. 1719 (*Nephopteryx*)  
 Swan River W.A., Richardson; HT ♀ unique, abdomen missing, BMNH.  
 Hudson 1928, p. 156, pl. xix fig. 2, as *Crocydopora cinigerella*.

***stenopterella*** Meyrick, 1879a, pp. 200–201 (*Nephopteryx*). Synonymised by Meyrick (1886f, p. 804).

“Sydney, Bowenfels and Newcastle” N.S.W., E. Meyrick; HT not located.

Hudson 1928, p. 156, as synonym.

Note. *C. cinigerella* probably occurs naturally in New Zealand.

Also 2 undescribed species (NZAC).

- ***Delogenes*** Meyrick, 1918a, p. 132. Type species *Delogenes limodoxa* Meyrick, by original monotypy.

» Pyralidae, *Delogenes*

**limodoxa** Meyrick, 1918a, p. 132 (*Delogenes*)  
Waitati DN, C.E. Clarke (Hudson 1928, p. 157); LT ♂  
here designated, labelled "Wellington New Zealand  
GVH. 2.17", "Delogenes limodoxa Meyrick 2/4 E.  
Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1928, p. 157, pl. xliv fig. 12, but illustration not  
patterned enough, and too blue.

• ***Ephestia*** in the sense of Kloet & Hincks (1972,  
p. 46), not Roesler (1966, pp. 104–160)

**cautella** Walker, 1863b, p. 73 (*Pempelia*)  
Type locality "Ceylon" (Sri Lanka).  
New Zealand: adventive; around human habitation,  
throughout.  
Not mentioned by Hudson. Roesler 1966, pp. 134–135,  
fig. 21 and 43, in *Cadra* Walker.

**elutella** Hübner, 1796, pl. 24 fig. 103 (*Tinea*)  
Type locality Europe.  
New Zealand: adventive; around human habitation,  
throughout.  
Hudson 1939, p. 418, not figured. Roesler 1966, pp. 130–  
131, fig. 18 and 40, in *Ephestia* Guenée.  
Note. Philpott (1928a, p. 361) gives an early record.

**kuehniella** Zeller, 1879, p. 466 (*Ephestia*)  
Europe.  
New Zealand: adventive; around human habitation,  
throughout.  
Hudson 1928, pp. 156–157, pl. xliv fig. 13. Corbet & Tams  
1943, p. 68, fig. 58, 64, 112, and 149, as *Ephestia ser-*  
*icarium* not of Scott (1859, p. 207, pl. 61). Roesler  
1966, pp. 126–127, fig. 15 and 37 (in *Anagasta* Heinrich,  
subgenus of *Ephestia*).

• ***Etiella*** Zeller, 1839, p. 179. Type species *Phycis*  
*zinckenella* Treitschke 1832, by monotypy.

**behrii** Zeller, 1848, p. 883 (*Phycis*) new record  
Type locality Adelaide S.A. (Whalley 1973, p. 18).  
New Zealand: adventive; 2 records, Auckland AK, Jan-  
uary 1981, C.J. Green, and Opunake TK, April 1982,  
F. Chambers; not found again 1982–85 (NZAC).

• ***Homoeosoma*** of authors, not of Curtis (1833, p.  
190). Type species *Phycis gemina* Haworth, 1811,  
by original monotypy; England, "Italia".

**anaspila** Meyrick, 1901, pp. 566–567  
(*Homoeosoma*)  
Christchurch MC, R.W. Fereday; LT ♀ here designated,  
labelled "Christchurch New Zealand RWF /86",  
"Homoeosoma anaspila Meyrick 1/4 E. Meyrick det.  
in Meyrick Coll.", BMNH.

Hudson 1928, p. 157, as synonym of *vagella* of authors;  
1939, p. 418, pl. lvi fig. 34, as species.

**ischnomorpha** Meyrick, 1931a, p. 94 (*Homoeosoma*)  
Whangarei ND, S.C. Patterson; HT ♂ unique, BMNH.  
Hudson 1939, p. 418, pl. lvi fig. 14.  
Note. The status of *Homoeosoma* species in New Zealand  
requires evaluation.

• ***Oligochroa*** Ragonot, 1888, p. 20. Type species  
*Pempelia dionysia* Zeller, 1846, by original  
designation.

**oculiferella** Meyrick, 1879b, p. 222 (*Pempelia*)  
[Morpeth N.S.W.], E. Meyrick; ST ♀ labelled "23/1/78",  
"SYNTYPE", "Pempelia oculiferella Meyr. 1/5 E.  
Meyrick det. in Meyrick Coll.", BMNH.  
Hudson 1939, p. 417, pl. lvi fig. 19, as *Salebria sublignalis*.  
Notes. The other ST ♀ is labelled "8/3/78". In his Diary  
of Captures, Meyrick records for 23 January 1878  
"Morpeth ... Pempelia oculiferella, 1", and for 8 March  
1878 "Randwick ... Pemp. mesonyctella, 1, Pemp. ruf-  
itinctella, 1". *Trachonitis sublignalis* Walker, 1863b,  
p. 41, from Sydney, is now a junior synonym of *Fav-*  
*eria laiasalis* Walker, 1859c, p. 889, and is distinct  
from *O. oculiferella*. Meyrick (1931a, p. 95) gives first  
records – Lake Rotomahana BP in 1915, and Wha-  
ngarei ND [late 1920s].

• ***Patagoniodes*** Roesler, 1969, p. 254. Type species  
*Patagoniodes popescugorji* Roesler, by original  
designation.

**farinaria** Turner, 1904, p. 128 (*Homoeosoma*)  
Tasmania, A.J. Turner; HT ♀ in ANIC.  
Hudson 1928, p. 157, pl. xx fig. 11, as *Homoeosoma*  
*vagella*; 1939, p. 418, as *Homoeosoma farinaria*, after  
Philpott (1928g, p. 485). McQuillan & Ireson 1987, p.  
240, as *Patagoniodes farinaria*.  
Note. Probably occurs naturally in New Zealand; is  
vagrant on The Snares (D.S. Horning Jr, pers. comm.).

• ***Plodia*** Guenée, 1845, p. 318. Type species *Tinea*  
*interpunctella* Hübner, by original monotypy.

**interpunctella** Hübner, [1810–1813], pl. 45 fig. 310  
(*Tinea*)  
Europe.  
New Zealand: adventive; around human habitation,  
throughout.  
Hudson 1928, p. 156, pl. xlvi fig. 7.  
Note. Meyrick (1915a, p. 202) reports this species from  
Wellington WN, taken by Hudson.

• ***Sporophyla*** Meyrick, 1905, p. 224. Type species  
*Crocypodora oenospora* Meyrick, by original  
monotypy.

***oenospora*** Meyrick, 1897b, p. 388 (*Crocydopora*)  
Castle Hill MC, G.V. Hudson; HT ♀ unique, abdomen  
missing, BMNH.  
Hudson 1928, p. 156, pl. xix fig. 1, as *Sporophyla*  
*oenospora*.

Note. Hudson does not mention Castle Hill as a locality  
for this species. It is possible that the HT was col-  
lected by George Howes in Central Otago CO.

- Also 1 undescribed genus and species (NZAC).

—◎—

### Subfamily PYRALINAE

(in the sense of Munroe 1972, pp. 12–14)

• ***Aglossa*** Latreille, 1796, p. 145. Type species *Pha-  
laena Pyralis pingualis* Linnaeus, by subsequent  
monotypy (Latreille, [1802], p. 414).

***caprealis*** Hübner, [1800–1809], pl. 23 fig. 153  
(*Pyralis*)

Europe.

New Zealand: adventive; widespread around human hab-  
itation. Philpott (1931, pp. 29–30) gives early records,  
as *A. cuprealis*.

• ***Diplopseustis*** Meyrick, 1884e, p. 284. Type spe-  
cies *Cymoriza minima* Butler, 1881, by subsequent  
designation (Ragonot 1891, p. 520).

***perieresalis*** Walker, 1859c, pp. 958–959 (?*Ambia*)  
Sarawak, Borneo, Saunders; HT ♂ not seen, HCOE.  
Hudson 1928, p. 205, pl. xxii fig. 44, as *Diplopseustis per-  
ieralis*, after Meyrick (1913c, p. 46).

***minima*** Butler, 1881a, p. 684 (*Cymoriza*). Syn-  
onymised by Hampson (1896, p. 489).  
Formosa (Taiwan); HT ♂, abdomen missing, BMNH.  
Hudson 1928, p. 205, as synonym.

***nana*** Warren, 1896a, p. 225 (*Sufetula*). Synony-  
mised by Hampson (1896, p. 489).  
Bombay, India; HT (?gender) in BMNH.

• ***Endotricha*** Zeller, 1847, p. 593. Type species *Py-  
ralis flammealis* Denis & Schiffermüller, 1775, by  
original designation.

***pyrosalis*** Guenée, 1854, pp. 219–220 (*Endotricha*)  
“Australie, Nouvelle Hollande”; HT in MNHP.  
Note. Philpott (1920, p. 44) records this species once, col-  
lected probably by A. Hamilton on Mt Dennan, Tar-  
arua Range WN. No other specimens have been  
reported.

• ***Gauna*** Walker, 1866a, p. 1252. Type species  
*Gauna subferralis* Walker, 1866a, p. 1253, by origi-  
nal monotypy.

***aegusalis*** Walker, 1859c, p. 912 (*Pyralis*)  
Moreton Bay Qld, Diggles; HT ♂ so labelled, BMNH.  
Hudson 1939, p. 428, pl. lvi fig. 20, as *Gauna aegalis*,  
after Turner (1904, p. 185).

Note. Philpott (1931, p. 30) gives early records.

***subferralis*** Walker, 1865, p. 1253 (*Gauna*). Syn-  
onymised by Hampson (1896, p. 486).

• ***Pyralis*** Linnaeus, 1758, p. 533; see Munroe (1972,  
p. 8). Type species *Pyralis farinalis* Linnaeus, by  
subsequent designation (ICZN 1957, p. 254).

***farinalis*** Linnaeus, 1758, p. 533 (*Pyralis*)

Europe.

New Zealand: adventive; around human habitation,  
throughout. Meyrick (1885a, p. 346) reported it pres-  
ent in Christchurch MC.

—◎—

### Family CRAMBIDAE

(in the sense of Minet 1981, pp. 267–268)

#### Subfamily CRAMBINAE


(in the sense of Munroe 1972, pp. 12–14)

• ***Angustalius*** Marion, 1954, pp. 42 and 50. Type  
species *Angustalius ditaeniellus* Marion, by original  
monotypy; Madagascar.

***malacelloides*** Bleszinski, 1955, p. 229 (*Crambopsis*)  
Tasmania, ?collector; HT ♂ designated by Bleszinski  
(1955, p. 230), IZWP.

Hudson 1939, p. 420, pl. lvi fig. 29, as *Crambus mala-  
cellus* [of authors]. Gaskin 1975, p. 346, fig. 22g and  
83.

Note. First noted at Whangarei ND (Philpott, 1931, p.  
26) in 1927; now generally distributed in lowland North  
Island localities.



Crambidae: Crambinae  
(174) *Orocrambus flexuosellus* (Doubleday)

➤ Crambidae, Crambinæ

● **Culladia** Moore, 1886, p. 383. Type species *Araxes admigratella* Walker, 1863b, by original designation; Ceylon.

**strophaea** Meyrick, 1905, p. 226 (*Argyria*)  
Wellington WN, G.V. Hudson; LT ♂ designated by Gaskin (1973, p. 451), BMNH.  
Hudson 1928, p. 170, pl. xix fig. 35, as *Argyria strophaea*.  
Gaskin 1973, p. 451 (*Culladia*), fig. 19, 20, and 27.

● **Gadira** Walker, 1866b, p. 1742. Type species *Gadira acerella* Walker, by original monotypy.

**Scenoploca** Meyrick, §1882a, p. 186; 1883b, pp. 6 (key) and 9. Type species *Scenoploca petraula* Meyrick, by original monotypy. Synonymised by Gaskin (1973, p. 454).

**Cryptomima** Meyrick, §1882a, p. 186; 1883b, pp. 6 (key) and 8. Type species *Gadira acerella* Walker, by original monotypy. Synonymised by Gaskin (1973, p. 454).

**acerella** Walker, 1866b, p. 1742 (*Gadira*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1928, p. 176, pl. xxii fig. 29. Gaskin 1973, p. 457, fig. 3, 16, 21, and 28.

**mahanga** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 27 (*Botys*). Synonymised by Meyrick (1883, p. 8).

Hudson 1928, p. 176, as synonym. Gaskin 1973, p. 457, as synonym, as "mehanga".

Note. Gaskin (1973, p. 454) wrongly ascribed *Botys* to Felder (1875); the reference is to *Botys* in the sense of Felder, not of Latreille (1802).

**leucophthalma** Meyrick, §1882a, p. 186; 1883b, p. 7 (*Thinasotia*; misspelling of *Thisanotia*)  
[Lyttelton, on hills], Christchurch MC, E. Meyrick; LT ♂ designated by Gaskin (1973, p. 459), BMNH.  
Hudson 1928, p. 172, pl. xxi fig. 35, as *Talis leucophthalma*. Gaskin 1973, pp. 459–460, fig. 17, 22, and 29.

**petraula** Meyrick, §1882a, p. 186; 1883b, pp. 9–10 (*Scenoploca*)  
[Lyttelton, on hills], Christchurch MC, E. Meyrick; LT ♂ designated by Gaskin (1973, p. 460), BMNH.  
Hudson 1928, p. 172, pl. xix fig. 34, as *Scenoploca petraula*. Gaskin 1973, pp. 460–462, fig. 18, 23, and 31.

● **Glaucocharis** Meyrick, 1938, p. 426. Type species *Glaucocharis stella* Meyrick, by original monotypy.  
Note. Gaskin (1985, p. 11) resurrected *Glaucocharis*, as referring to all New Zealand species previously placed (Gaskin 1971) in *Pareromene*, and gives a full synonymy of this widely distributed genus.

**Pareromene** Osthelder, 1941, p. 366. Type species by original designation *Pareromene rebeli* Osthelder, 1941 (= *Pareromene euchromiella* (Ragonot, 1895)); Mediterranean. Synonymised by Gaskin (1985, p. 11).

**auriscriptella** Walker, 1864b, p. 976 (*Eromene*)  
[Auckland AK], D. Bolton; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 768), BMNH.  
Hudson 1928, p. 175, pl. xix fig. 37, as *Diptychophora auriscriptella*. Gaskin 1971, fig. 1, 28, and 46, as *Pareromene auriscriptella*.

**bipunctella** Walker, 1866b, p. 1761 (*Eromene*)  
[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1928, p. 175, pl. lii fig. 30; Gaskin 1971, p. 770, fig. 2, 29, and 47.

**chrysoclyta** Meyrick, §1882a, p. 186; 1883b, pp. 11 (key) and 12 (*Diptychophora*)  
[The Domain], Auckland AK, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 771), BMNH.  
Hudson 1928, p. 173, pl. xix fig. 10, as *Diptychophora chrysoclyta*. Gaskin 1971, pp. 771–772, fig. 3, 29 and 47, as *Pareromene chrysoclyta*.

**elaina** Meyrick, §1882a, p. 187; 1883b, pp. 11 (key) and 17 (*Diptychophora*)  
[Botanic Gardens], Wellington, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 773), BMNH.  
Hudson 1928, p. 176, pl. xix fig. 31, as *Diptychophora elaina*. Gaskin 1971, pp. 773–774, fig. 4, 31, and 49, as *Pareromene elaina*.  
Note. Gaskin gives the LT label date as 31/12/59; this should read 31/12/79.

**epiphaea** Meyrick, §1885a, p. 347; 1885f, p. 132 (*Diptychophora*)  
Arthur's Pass NC/WD, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 776), BMNH.  
Hudson 1928, p. 176, pl. xix fig. 40, as *Diptychophora epiphaea*. Gaskin 1971, pp. 776–777, fig. 5, 36, and 66, as *Pareromene epiphaea*.

**harmonica** Meyrick, 1888c, p. 71 (*Diptychophora*)  
[Waitakere Range], Auckland AK, E. Meyrick, LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 779), BMNH.  
Hudson 1928, p. 175, pl. xix fig. 38, as *Diptychophora harmonica*. Gaskin 1971, pp. 779–780, fig. 7, 30 and 48, as *Pareromene harmonica*.

**helioclyta** Meyrick, §1882a, p. 187; 1883b, pp. 11 (key) and 17 (*Diptychophora*)  
Lake Wakatipu OL, R.W. Fereday; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 780), BMNH.  
Hudson 1928, pp. 175–176, pl. xix fig. 39, as *Diptychophora helioclyta*. Gaskin 1971, pp. 780–782, fig. 8, 32, and 50, as *Pareromene helioclyta*.

**holanthes** Meyrick, §1885a, p. 347; 1885f, p. 131  
(*Diptychophora*)

Otira Gorge WD, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 782), BMNH.

Hudson 1928, p. 175, pl. xix fig. 36, as *Diptychophora holanthes*. Gaskin 1971, pp. 782–783, fig. 9, 37, and 56, as *Pareromene holanthes*.

Note. LT has bright yellow forewings and dark hindwings; D.E. Gaskin (pers. comm.) notes that populations can vary in hindwing colour from dark to pale.

**interrupta** Felder & Rogenhofer, 1875, pl. cxxxv fig. 15 (*Crambus*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.

Hudson 1928, p. 173, pl. xix fig. 13, as *Diptychophora interrupta*. Gaskin 1971, pp. 792–793, fig. 10, 33, and 51, as *Pareromene interrupta*.

**astrosema** Meyrick, §1882a, p. 186; 1883b, pp. 11 (key) and 13 (*Diptychophora*). Synonymised by Meyrick (§1885a, p. 347; 1885f, p. 130).

Christchurch MC, R.W. Fereday; HT ♂ unique, BMNH. Hudson 1928, p. 173, as synonym.

**lepidella** Walker, 1866b, p. 1761 (*Eromene*)

[Nelson NN], T.R. Oxley; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 794), BMNH.

Hudson 1928, p. 174, pl. xix fig. 14 and 15, as *Diptychophora lepidella*. Gaskin 1971, pp. 793–795, fig. 11, 34, and 52, as *Pareromene lepidella*.

**gracilis** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 26 (*Crambus*). Synonymised by Meyrick (1883b, p. 14).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.

Hudson 1928, p. 174, as synonym.

Note. Butler (1877, p. 401) erroneously synonymised *gracilis* with *aurascriptella*.

**leucoxantha** Meyrick, §1882a, p. 186; 1883b, p. 15  
(*Diptychophora*)

Lake Wakatipu OL, R.W. Fereday; HT ♂ unique, BMNH.

Hudson 1928, p. 174, pl. xix fig. 16 and 17, as *Diptychophora leucoxantha*. Gaskin 1971, pp. 795–797, fig. 12, 27, and 45, as *Pareromene leucoxantha*.

**metallifera** Butler, 1877, p. 401 (*Eromene*)

[Dunedin DN], J. Hector; HT ♂ designated by Butler, BMNH.

Hudson 1928, p. 174, pl. xix fig. 32 and 33, as *Diptychophora metallifera*. Gaskin 1971, pp. 797–799, fig. 13, 35, and 53, as *Pareromene metallifera*.

**microdora** Meyrick, 1905, p. 227 (*Diptychophora*)

[Flora Saddle], Mount Arthur NN, E. Meyrick; LT ♀ selected by S. Bleszynski (Gaskin 1971, p. 799), BMNH.

Hudson 1928, p. 173, pl. xix fig. 12, as *Diptychophora microdora*. Gaskin 1971, pp. 799–801, fig. 14, 38, and 54, as *Pareromene microdora*.

Note. The gender of the LT is ♀, not ♂ as reported by Gaskin (1971, p. 799).

**parormene** Meyrick, 1924a, p. 202 (*Diptychophora*)  
Mount Ruapehu TO, G.V. Hudson; LT ♂ selected by S. Bleszynski (Gaskin 1971, p. 801), BMNH.

Hudson 1928, p. 176, not figured, as *Diptychophora parormene*. Gaskin 1971, pp. 801–802, fig. 14, 40, and 57, as *Pareromene parormene*.

**planetopa** Meyrick, 1923, p. 162 (*Diptychophora*)  
[Routeburn Valley], Lake Wakatipu OL, G.V. Hudson;  
HT ♀ unique, slide BM Pyral. 14269 (Gaskin 1974,  
p. 182), BMNH.

Hudson 1928, p. 174, pl. xlvi fig. 19, as *Diptychophora planetopa*. Gaskin 1971, pp. 802–803, fig. 16 and 39 (♀ genitalia not figured); 1974, p. 182.

**pyrsophanes** Meyrick, §1882a, p. 186; 1883b, p. 11  
(*Diptychophora*)

[Botanic Gardens, Wellington WN, E. Meyrick; LT ♀ selected by S. Bleszynski (Gaskin 1971, p. 803, as ♂; corrected, Gaskin 1974, p. 182), BMNH.

Hudson 1928, p. 173, pl. xix fig. 11, as *Diptychophora pyrsophanes*. Gaskin 1971, pp. 803–805, fig. 17, 41, and 58, as *Pareromene pyrsophanes*.

**selenaea** Meyrick, §1885a, p. 347; 1885f, p. 131  
(*Diptychophora*)

Dunedin DN, A. Purdie; LT ♀ selected by P.E.S. Whalley (Gaskin 1971, p. 805, as ♂; corrected, Gaskin 1974, p. 182), BMNH.

Hudson 1928, pp. 174–175, pl. xix fig. 30, as *Diptychophora selenaea*. Gaskin 1971, pp. 805–806, fig. 18, 25, and 43, as *Pareromene selenaea*.

**stella** Meyrick, 1938, pp. 426–427 (*Glaucchocharis*)  
Orongorongo Valley WN, Stella Gibbs [Hudson]; HT ♂ unique, BMNH.

Hudson 1939, p. 428, pl. lxii fig. 10, in Scopariinae.

**gurri** Gaskin, 1971, pp. 777–778 (*Pareromene*).  
Synonymised by Gaskin (1985).

Tapu–Coroglen Road CL, J.S. Dugdale; HT ♂ designated by Gaskin, NZAC.  
Gaskin 1971, fig. 6, 26, and 44.

• **Kupea** Philpott, 1930a, p. 247. Type species *Kupea electilis* Philpott, by original designation and monotypy.

**electilis** Philpott, 1930a, p. 247 (*Kupea*)  
Birdling's Flat MC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 420, pl. lvi fig. 37 (recorded as ♀). Gaskin 1975, p. 345, fig. 22f and 25.

Note. Females of *K. electilis* were unknown to Philpott, Lindsay, and Hudson.

• **Maoricrambus** Gaskin, 1975, p. 344. Type species *Crambus oncobolus* Meyrick, by original designation.

**oncobolus** Meyrick, §1885a, p. 348; 1885f, p. 138 (*Crambus*)

Castle Hill [Basin] MC, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 344), BMNH. Hudson 1928, p. 169, pl. xx fig. 35, as *Crambus oncobolus*. Gaskin 1975, pp. 344–345, fig. 24 ♂, 24 ♀, and 81, as *Maoricrambus oncobolus*.

**oncolobus** Hampson, 1895, p. 940, misspelling of *oncobolus* Meyrick.

• **Orocrambus** Purdie, 1884, p. 168 (no generic description). Type species by original monotypy *Orocrambus melampetrus* (Meyrick MS.) Purdie, 1884, p. 168; available name according to Bleszynski & Collins (1962, p. 329) and Gaskin (1975, p. 277).

Note. This Meyrick manuscript name and the specific epithet were made nomenclaturally available by Purdie before their publication by Meyrick.

**Orocrambus** Meyrick, §1885a, p. 347; 1885f, p. 133. Type species by original monotypy *Orocrambus melampetrus* Meyrick. Available name according to Meyrick (1885f, p. 133; 1913b, p. 32) but regarded as a synonym by Gaskin (1975, p. 277), and as a junior homonym by Fletcher & Nye (1984, p. 107).

Note. Hampson did not ascribe *Orocrambus* to Purdie, and Purdie did not “use details from an abstract”, he used the binomen Meyrick had attached to a specimen (since lost) then in OMNZ, and lacking a locality label (cf. Gaskin 1975, p. 277).

**abditus** Philpott, 1924a, p. 212 (*Crambus*)

Otarama MC, S. Lindsay; HT ♀ designated by Philpott, CMNZ.

Hudson 1928, p. 167, pl. 1 fig. 18, as *Crambus abditus*. Gaskin 1975, p. 301, fig. 3a, 11a, and 29, as *Orocrambus abditus*.

**aethonellus** Meyrick, §1882a, p. 187; 1883b, p. 19 (*Crambus*)

Mount Hutt MC, R.W. Fereday; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 301), BMNH.

Hudson 1928, p. 162, not figured, as *Crambus aethonellus*. Gaskin 1975, pp. 301–302, fig. 3b, 11b, 30, and 31, as *Orocrambus aethonellus*.

**antimorus** Meyrick, 1901, p. 567 (*Crambus*). Synonymised by Gaskin (1975, p. 301).

Mount Cook MK, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 161, pl. xix fig. 23, as *Crambus antimorus*.

**heteranthes** Meyrick, 1901, p. 568 (*Crambus*). Synonymised by Gaskin (1975, p. 301).

Mount Cook MK, G.V. Hudson; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 301), BMNH. Hudson 1928, p. 161, pl. xix fig. 7, as *Crambus heteranthes*.

**saristes** Meyrick, 1909a, p. 8 (*Crambus*). Synonymised by Gaskin (1975, p. 301).

Invercargill SL, A. Philpott; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 302), BMNH.

Hudson 1928, p. 162, pl. xlv fig. 18, as *Crambus saristes*.

**aulistes** Meyrick, 1909a, p. 9 (*Crambus*). Synonymised by Gaskin (1975, p. 301).

Invercargill SL, “Hudson” [?Howes]; HT ♂ unique, BMNH.

Hudson 1928, p. 162, as *Crambus aulistes*.

Note. Hudson (1928) states “I am unacquainted with this species”.

**meristes** Meyrick, 1919, p. 351 (*Crambus*). Synonymised by Gaskin (1975, p. 301).

Longwood Range SL, A. Philpott; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 302), BMNH.

Hudson 1928, p. 162, not figured, as *Crambus meristes*.

**meritus** Philpott, 1929b, p. 496 (*Crambus*); misspelling of *meristes* Meyrick (Gaskin 1975, p. 301).

**angustipennis** Zeller, 1877, p. 15 (*Chilo*)

[locality uncertain, probably Christchurch MC], H.G. Knaggs; HT ♀ unique, BMNH.

Hudson 1928, p. 163, pl. xx fig. 38, as *Crambus angustipennis*, after Meyrick (§1882a, p. 187; 1883b, p. 22).

Gaskin 1975, p. 303, fig. 3d (as 3c), 11c, and 32, as *Orocrambus angustipennis*.

Note. Gaskin (1987) corrects this typographic error; his fig. 3c shows ♂ genitalia of *apicellus*.

**apicellus** Zeller, 1863, p. 31 (*Crambus*)

“Neu Seeland”; original material lost (Gaskin 1975, p. 304), NHMW.

Hudson 1928, p. 167, pl. xx fig. 20, as *Crambus apicellus*. Gaskin 1975, p. 304, fig. 3c (as 3d), 11d, and 33, as *Orocrambus apicellus*.

Note. Gaskin (1987) corrects this typographic error; his fig. 3d shows ♂ genitalia of *angustipennis*.

**callirrhous** Meyrick, §1882a, p. 187; 1883b, pp. 19 (key) and 24 (*Crambus*)

Lake Guyon MB/BR, R.W. Fereday; LT ♂ designated by Gaskin (1975, p. 305), CMNZ.

Hudson 1928, p. 165, pl. xx fig. 34, as *Crambus callirrhous*. Gaskin 1975, p. 305, fig. 3e, 11e, and 34, as *Orocrambus callirrhous*.

**schedias** Meyrick, 1911b, p. 60 (*Crambus*). Synonymised by Gaskin (1975, p. 305).

Wellington WN, G.V. Hudson; HT ♂ unique (Gaskin 1975, p. 305), BMNH.

Hudson 1928, p. 166, pl. xx fig. 33, as *Crambus schedias*.

**callirhous** Bleszynski & Collins, 1962, p. 258, misspelling of *callirrhous*.

**catacaustus** Meyrick, §1885a, p. 347; 1885f, p. 134 (*Crambus*)

Arthur's Pass NC/WD, R.W. Fereday; LT ♂ designated by Gaskin (1975, p. 305), BMNH.

Hudson 1928, p. 159, pl. xx fig. 25. Gaskin 1975, pp. 305–306, fig. 4a, 11f, and 35.

**pervius** Meyrick, 1912c, p. 118 (*Orocrambus*).  
Synonymised by Gaskin (1975, p. 305).  
Lake Wakatipu OL, G.V. Hudson; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 305), BMNH.  
Hudson 1928, p. 159, pl. xx fig. 24, as species.

**clarkei clarkei** Philpott, 1930b, p. 3, line 20 (*Orocrambus*; as species)  
Mount Moltke, Franz Josef WD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, pp. 418–419, pl. lvi fig. 6, as species. Gaskin 1975, pp. 306–307, fig. 4b and 36.

**nebulosa** Philpott, 1930b, p. 3, line 1 (*Tauroscopa*). Synonymised by Gaskin (1975, p. 306).  
Minaret Peaks OL, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 421, pl. lvi fig. 5, as *Tauroscopa nebulosa*.

**clarkei eximia** Salmon, 1946, p. 6 (*Tauroscopa*; as species)  
Homer Cirque FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.  
Hudson 1950, p. 100, pl. vii fig. 4, as *Tauroscopa eximia*.  
Gaskin 1975, p. 307, fig. 4b, 12a, and 37, as *Orocrambus clarkei eximia*.

**corruptus** Butler, 1877, p. 399 (as *Hypochalcia corrupta*)  
[Dunedin DN or mid Canterbury MC], J. Hector or J.D. Enys; HT ♂ designated by Butler, BM Pyral. slide 5567, BMNH.  
Hudson 1928, p. 161, pl. xx fig. 1, as *Crambus corruptus*.  
Gaskin 1975, p. 308, fig. 4c, 12b, and 38, as *Orocrambus corruptus*.

**luridus** Hudson, 1923a, p. 64 (*Crambus*).  
Synonymised by Hudson (1928, p. 161).  
Taylor's Mistake, Banks Peninsula MC, C.E. Clarke; ST ♂ and ♀ not located.  
Hudson 1928, p. 161, as synonym.  
Note. The month of capture (August) is unusual.

**crenæus** Meyrick, §1885a, p. 348; 1885f, pp. 135–136 (*Crambus*)  
Arthur's Pass NC/WD, E. Meyrick; LT ♂ selected by Gaskin (1975, p. 308), CMNZ.  
Hudson 1928, p. 165, pl. xx fig. 9, as *Crambus crenæus*.  
Gaskin 1975, pp. 308–309, fig. 4d and 39, as *Orocrambus crenæus*, ♂; 1987, p. 114, fig. 1, 2, and 5, ♀.

**diploorrhous** Meyrick, §1885a, p. 348; 1885f, pp. 136–137 (*Crambus*).  
Synonymised by Gaskin (1975, p. 308).  
[Ben Lomond] OL, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 308), BMNH.  
Hudson 1928, p. 164, pl. xx fig. 19, as *Crambus diploorrhous*.

**cultus** Philpott, 1917b, p. 242 (*Orocrambus*)  
Cecil Peak OL, M.O. Pasco; PT ♀ labelled "2143", "Cecil Pk 8.1.14", "Lectotype Orocrambus cultus Philpott

1916 label 7.1.81. B. Patrick rediscovered", "Paratype ♀ Orocrambus cultus Philpott, rediscovered by B. Patrick 7.1.81 x Southland Museum", NZAC.  
Hudson 1928, p. 160, pl. xliv fig. 10. Gaskin 1975, p. 309, not seen.

Note. PT ♀ is the only remaining member of the ST series, and was rescued from an old public display cabinet in the Southland Museum by Mr B. Patrick. Neither he nor I could locate the other ♀ and 2 ♂♂ of the type series, and Mr Patrick considers that they had disintegrated during the years the material was on display.

**cyclopicus** Meyrick, §1882a, p. 187; 1883b, pp. 19 (key) and 29 (*Crambus*)

[Lyttelton, on hills] MC, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 309), BMNH.  
Hudson 1928, p. 169, pl. xx fig. 44, as *Crambus cyclopicus*. Gaskin 1975, pp. 309–311, fig. 4e, 12c, 18c, 19d, and 40, as *Orocrambus cyclopicus*.

**sophistes** Meyrick, 1905, p. 226 (*Crambus*).  
Synonymised by Gaskin (1975, p. 309).

[Ida Valley CO], J.H. Lewis; HT ♂ unique, BMNH.  
Hudson 1928, p. 169, pl. xix fig. 24, as species.

**dicrenellus** Meyrick, §1882a, p. 187; 1883b, pp. 19 (key) and 22 (*Crambus*)

Mount Hutt MC, R.W. Fereday; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 312), BMNH.  
Hudson 1928, p. 164, pl. xx fig. 18 ♂ (not ♀ as captioned), as *Crambus dicrenellus*. Gaskin 1975, p. 312, fig. 4f and 41, as *Orocrambus dicrenellus*.

**enchophorus** Meyrick, §1885a, p. 348; 1885f, p. 136 (*Crambus*)

Castle Hill [Basin] MC, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 312), BMNH.  
Hudson 1928, p. 165, pl. xx fig. 16, as *Crambus enchophorus*. Gaskin 1975, pp. 312–314, fig. 5a, 12d, 18d, 19e, and 42, as *Orocrambus enchophorus*.

**pedias** Meyrick, §1885a, p. 348; 1885f, p. 37 (*Crambus*).  
Synonymised by Gaskin (1975, p. 312).  
Wanganui WI, E. Meyrick; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 312), BMNH.

Hudson 1928, p. 166, pl. xx fig. 32, as species.  
**scitulus** Philpott, 1926a, p. 390 (*Crambus*).  
Synonymised by Gaskin (1975, p. 312).  
Mount Arthur NN, 4000 ft, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 165, pl. li fig. 23, as species.

**ephorus** Meyrick, §1885a, p. 348; 1885f, p. 135 (*Crambus*)

Arthur's Pass NC/WD, 4800 ft, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 163, pl. xlvi fig. 17, as *Crambus ephorus*. Gaskin 1975, p. 314, fig. 5b, 12e, and 43, as *Orocrambus ephorus*.

**corylana** Clarke, 1926, p. 417 (*Crambus*).  
Synonymised by Gaskin (1975, p. 314).

➤ Crambidae, *Orocrambus ephorus*

Mount Ida CO, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.  
Hudson 1928, p. 163, not figured, as *Crambus corylanus*.  
Gaskin 1975, p. 314, as synonym.

**flexuosellus** Doubleday, 1843, p. 289 (*Crambus*)  
"New Zealand", E. Dieffenbach; ST ♂ labelled "New  
Zealand 42-55", BMNH.  
Hudson 1928, p. 168, pl. xx fig. 31, as *Crambus flexu-*  
*osellus*. Gaskin 1975, pp. 314-317, fig. 5c, 12f, 18c,  
19f, and 44, as *Orocrambus flexuosellus*.

**fugitivellus** Hudson, 1950, p. 99 (*Crambus*)  
"Mackenzie Country" MK, R. Dick; HT ♂ unique,  
NMNZ.  
Hudson 1950, p. 99, pl. iv fig. 2, as *Crambus fugitivellus*.  
Gaskin 1975, p. 317, fig. 5d and 45.

**haplotomus** Meyrick, §1882a, p. 187; 1883b, pp. 19  
(key) and 23 (*Crambus*)  
Lake Wakatipu OL, R.W. Fereday; LT ♂ selected by S.  
Bleszynski (Gaskin 1975, p. 317), abdomen missing,  
BMNH.  
Hudson 1928, p. 165, pl. xx fig. 7, as *Crambus haplotomus*.  
Gaskin 1975, pp. 317-318, fig. 5e, 13a, and 46,  
as *Orocrambus haplotomus*.

**harpophorus** §Meyrick, 1882a, p. 187; 1883b, pp.  
19 (key) and 30 (*Crambus*)  
Lake Wakatipu OL, R.W. Fereday; LT ♂ selected by S.  
Bleszynski (Gaskin 1975, p. 318), BMNH.  
Hudson 1928, p. 169, pl. xx fig. 36, as *Crambus harpo-*  
*phorus*. Gaskin 1975, p. 318, fig. 5f, 13b, and 47.

**helioites** Meyrick, 1888c, p. 68 (*Crambus*)  
Mount Arthur NN, 3800 ft, E. Meyrick; LT ♂ selected  
by S. Bleszynski (Gaskin 1975, p. 319), BMNH.  
Hudson 1928, p. 161, pl. xix fig. 5 and 6, as *Crambus helioites*.  
Gaskin 1975, p. 319, fig. 6a, 13c, and 48, as  
*Orocrambus helioites*.

**heteraulus** Meyrick, 1905, p. 225 (*Crambus*)  
Humboldt Range OL, G.V. Hudson; HT ♂ unique,  
BMNH.  
Hudson 1928, p. 164, pl. xx fig. 37, as *Crambus heter-*  
*aulus*. Gaskin 1975, p. 320, fig. 6b, 13d, and 49, as  
*Orocrambus heteraulus*.

**horistes** Meyrick, 1902c, p. 276 (*Crambus*)  
Chatham Island, J. Fougere; LT ♂ selected by S. Ble-  
szynski (Gaskin 1975, p. 320), BMNH.  
Hudson 1928, p. 168, not figured, as *Crambus horistes*.  
Gaskin 1975, pp. 320-321, fig. 6c, 13e, and 50, as *Oro-*  
*crambus horistes*.

**isochytus** Meyrick, 1888c, p. 68 (*Crambus*)  
Mount Arthur NN, 4300 ft, E. Meyrick; LT ♂ selected  
by S. Bleszynski (Gaskin 1975, p. 321), BMNH.  
Hudson 1928, p. 164, pl. xx fig. 10, as *Crambus isochytus*.

Gaskin 1975, p. 321, fig. 6d, 13f, and 51, as *Orocram-*  
*bus isochytus*.

**jansoni** Gaskin, 1975, p. 322 (*Orocrambus*)  
Waiaouru TO, D.E. Gaskin; HT ♂ designated by Gaskin,  
NZAC.  
Gaskin 1975, p. 322, fig. 6e, 14a, and 52.

**lectus** Philpott, 1929a, p. 301 (*Crambus*)  
Lake Tekapo MK, A. Philpott; HT ♂ designated by Phil-  
pott, tip of abdomen missing, NZAC.  
Hudson 1939, p. 419, pl. lvi fig. 17, as *Crambus lectus*.  
Gaskin 1975, pp. 322-323, fig. 6f, 14b, and 53, as *Oro-*  
*crambus lectus*.

**lewisi** Gaskin, 1975, p. 323 (*Orocrambus*)  
Titahi Bay WN, G.V. Hudson; HT ♂ designated by Gas-  
kin, NMNZ.  
Gaskin 1975, pp. 323-324, fig. 7a, 14c, and 54.

**lindsayi** Gaskin, 1975, p. 324 (*Orocrambus*)  
Mount Ida CO, C.E. Clarke; HT ♀ designated by Gaskin,  
AMNZ.  
Gaskin 1975, p. 324, fig. 14d, 20d, and 55.

**machaeristes** Meyrick, 1905, p. 224 (*Orocrambus*)  
Mount Earnslaw OL, G.V. Hudson; LT ♂ selected by S.  
Bleszynski (Gaskin 1975, p. 324), BMNH.  
Hudson 1928, p. 160, pl. xx fig. 22. Gaskin 1975, pp. 324-  
325, fig. 7b, 14e, and 56.

**melampetrus** Purdie, 1884, p. 168 (*Orocrambus*)  
Mount Bonpland OL, A. Purdie; 4 STs (?gender) lost,  
OMNZ.  
Gaskin 1975, p. 325, fig. 7c, 14f, and 57.

**melampetrus** Meyrick, §1885a, p. 347; 1885f, p.  
133 (*Orocrambus*). Synonymised by Gaskin (1975,  
p. 325).  
Mount Hutt MC, R.W. Fereday; LT ♂ here designated,  
labelled "Mt Hutt New Zealand RWF /80", BMNH.  
Hudson 1928, p. 158, pl. xx fig. 26.

**melitastes** Meyrick, 1909a, p. 9 (*Crambus*)  
Invercargill SL, E. Meyrick; LT ♂ selected by S. Ble-  
szynski (Gaskin 1975, p. 326), BMNH.  
Hudson 1928, p. 162, pl. xix fig. 8 and 9, as *Crambus melitastes*.  
Gaskin 1976, p. 326, fig. 7d, 15a, 58, and 59.

**mylites** Meyrick, 1888c, p. 67 (*Orocrambus*)  
Mount Arthur NN, 4000 ft, E. Meyrick; LT ♂ selected  
by S. Bleszynski (Gaskin 1975, p. 326), BMNH.  
Hudson 1928, p. 158, pl. xx fig. 23. Gaskin 1975, pp. 326-  
327, fig. 7e, 15b, and 60.

**oppositus** Philpott, 1915, p. 197 (*Crambus*)  
Mount Cleughearn, Hunter Mountains FD, A. Philpott;  
HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 164, pl. xliv fig. 1 and 2, as *Crambus oppositus*.  
Gaskin 1975, p. 327, fig. 7f, 15c, and 61.

**ordishi** Gaskin, 1975, p. 327 (*Orocrambus*)  
Hinds MC, C.E. Clarke; HT ♂ designated by Gaskin,  
AMNZ.  
Gaskin 1975, pp. 327–328, fig. 8a, 15d, and 62.

**ornatus** Philpott, 1927d, p. 82 (*Crambus*)  
Golden Downs NN, A. Philpott; HT ♂ designated by  
Philpott, NZAC.  
Hudson 1939, pp. 419–420, pl. lvi fig. 28, as *Crambus*  
*ornatus*. Gaskin 1975, p. 328, fig. 8b and 63, as *Orocrambus*  
*ornatus*.

**paraxenus** Meyrick, §1885a, p. 348; 1885f, p. 137  
(*Crambus*)  
[Ben Lomond] OL, E. Meyrick; LT ♂ selected by S. Ble-  
szynski (Gaskin 1975, p. 329), BMNH.  
Hudson 1928, p. 167, pl. xx fig. 17, as *Crambus*  
*paraxenus*. Gaskin 1975, p. 329, fig. 8c and 64.

**philpotti** Gaskin, 1975, pp. 329–330 (*Orocrambus*)  
Iron Hill, [Cobb Valley] NN, 4700–5000 ft, J.S. Dugdale;  
HT ♂ designated by Gaskin, NZAC.  
Gaskin 1975, pp. 329–330, fig. 8d, 15e, and 65 [specimen  
from Mount Cedric, BR].

**punctellus** Hudson, 1950, p. 99 (*Crambus*)  
Portobello DN, W.G. Howes; HT ♀ unique, MNZNZ.  
Hudson 1950, p. 99, pl. vii fig. 10 (as ♂), as *Crambus*  
*punctellus*. Gaskin 1975, p. 330, fig. 15f and 66.

**ramosellus** Doubleday, 1843, p. 288 (*Crambus*)  
“New Zealand”, E. Dieffenbach; LT ♂ here designated,  
labelled “New Zealand 42-55”, abdomen missing,  
BMNH.

Hudson 1928, p. 163, pl. xx fig. 29 and 30, as *Crambus*  
*ramosellus*. Gaskin 1975, pp. 330–331, fig. 8e, 16a,  
and 67, as *Orocrambus ramosellus*.

**rangona** Felder & Rogenhofer, 1875, pl. cxxxvii  
fig. 25 (*Crambus*). Synonymised by Meyrick (1883b,  
p. 21).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1928, p. 163, as synonym.

**leucanialis** Butler, 1877, p. 401 (*Chilio*). Synon-  
ymised by Meyrick (1885, p. 139).  
[Dunedin DN or Christchurch MC], J. Hector or J.D.  
Enys; HT ♂ designated by Butler (as “Type”), BMNH.  
Hudson 1928, p. 163, as synonym.

**apsealias** Meyrick, 1907c, p. 109 (*Crambus*). Synon-  
ymised by Gaskin (1975, p. 330).  
Invercargill SL, G.V. Hudson [?W.G. Howes]; LT ♂  
selected by S. Bleszynski (Gaskin 1975, p. 331), BMNH.  
Hudson 1928, p. 162, pl. xx fig. 13, as *Crambus*  
*apsealias*.

**scoparioides** Philpott, 1914, p. 119 (*Orocrambus*)  
Ben Lomond OL, A. Philpott; HT ♂ designated by Phil-  
pott, NZAC.  
Hudson 1928, p. 160, pl. xlvi fig. 1. Gaskin 1975, pp. 332–  
333, fig. 8f, 16b, and 68.

**scutatus** Philpott, 1917b, pp. 242–243 (*Crambus*)  
Longwood Range SL, A. Philpott; HT ♂ designated by  
Philpott, NZAC.  
Hudson 1928, p. 164, pl. xliv fig. 3, as *Crambus*  
*scutatus*. Gaskin 1975, p. 333, fig. 9a and 69, ♂; 1987, p. 117,  
fig. 3, 4, and 6, ♀.

**simplex** Butler, 1877, p. 400 (*Chilo*)  
[Nelson NN], T.R. Oxley; HT ♂ here recognised, labelled  
“60-73 Auckland N. Zeal.”, “*Chilo simplex* Butler  
Type”, “*Pyralidae* B.M. 5562”, BMNH.  
Hudson 1928, p. 166, pl. xx fig. 14 and 15, as *Crambus*  
*simplex*. Gaskin 1975, pp. 333–335, fig. 9b, 16c, and  
70, as *Orocrambus simplex*.

Note. Data on *simplex* type material provided by BMNH  
for Gaskin were mixed, and this is an example of the  
confusion generated when the reviser is unable to see  
the types. Butler (1877, pp. 400–401) clearly states: “I  
have taken the description from an example previ-  
ously in the Museum Collection from Auckland [Nel-  
son, in this instance] and which Walker had  
confounded with *Crambus vitellus* of Doubleday”. The  
specimen bearing Butler’s type designation, and the  
accession number 60-73 (“Auckland, New Zealand,  
Oxley”) is therefore the HT, as it is the specimen on  
which Butler said he based his description.

**siriellus** Meyrick, §1882a, p. 187; 1883b, pp. 19 (key)  
and 25–26 (*Crambus*)  
[heath-like scrub and swamp], Hamilton WO, E. Meyr-  
ick; LT ♂ selected by S. Bleszynski (Gaskin 1975, p.  
335), BMNH.

Hudson 1928, pp. 166–167, pl. xx fig. 28, as *Crambus*  
*siriellus*. Gaskin 1975, p. 335, fig. 9c, 16d, and 71, as  
*Orocrambus siriellus*.

**sophronellus** Meyrick, §1885a, p. 348; 1885f, p. 138  
(*Crambus*)  
?Christchurch MC, R.W. Fereday; HT ♀ unique, BMNH.  
Hudson 1928, p. 169, pl. xx fig. 43, as *Crambus*  
*sophronellus*. Gaskin 1975, pp. 335–336, fig. 16e, as *Oro-  
crambus sophronellus*.

**thymiastes** Meyrick, 1901, p. 567 (*Orocrambus*)  
Invercargill SL, A. Philpott; LT ♀ selected by S. Ble-  
szynski (Gaskin 1975, p. 336), BMNH.  
Hudson 1928, p. 159, pl. xx fig. 21. Gaskin 1975, p. 336,  
fig. 9d, 16f, and 72.

**tritonellus** Meyrick, §1885a, p. 347; 1885f, pp. 134–  
135 (*Crambus*)  
Porter’s Pass MC, J.D. Enys; HT ♀ unique, BMNH.  
Hudson 1928, p. 160, pl. xx fig. 27, as *Orocrambus*  
*tritonellus*, after Meyrick (1913, p. 33). Gaskin 1975, pp.  
336–337, fig. 9e, 18a, and 73.

**tuhualis** Felder & Rogenhofer, 1875, pl. cxxxvii fig.  
18 (*Crambus*)  
[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.  
Hudson 1928, p. 168, pl. xx fig. 8, as *Crambus*  
*tuhualis*.

» Crambidae, *Orocrambus tuhualis*

Gaskin 1975, pp. 337–338, fig. 9f, 17b, and 74, as *Orocrambus tuhualis*.

**thrinodes** Meyrick, 1910a, pp. 64–65 (*Crambus*). Synonymised by Philpott (1926a, p. 391).

Kaitoke WN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 168, as synonym.

Note. Meyrick (1911b, pp. 64–65) redescribed *thrinodes* as a new species apparently on the same specimen, as the Hudson–Meyrick correspondence mentions only one specimen.

**ventosus** Meyrick, 1920a, p. 30 (*Orocrambus*)

Mount Arthur NN, 4200 ft, [Stella] Hudson; LT ♂ selected by S. Bleszynski and here designated (cf. Gaskin 1975, p. 338), labelled “Mt Arthur New Zealand GVH 4200’ 1.18”, BMNH.

Hudson 1928, p. 159, pl. xlvi fig. 15 and 16. Gaskin 1975, p. 338, fig. 10a, 17c, and 75.

Note. Gaskin’s action in selecting a NMNZ specimen as LT was incorrect (see ICZN Article 75 (a) (i), p. 81), as there is no proof that Meyrick had those specimens before him when he wrote the description. Bleszynski’s lectotype designation is here upheld, on the reasonable assumption that the 2 specimens in the Meyrick Collection (now in BMNH) are those 2 specimens (Meyrick 1920, p. 30) on which Meyrick’s description is based.

**vittellus** Doubleday, 1843, p. 289 (*Crambus*)

[Auckland AK], A. Sinclair; 2 ST ♂♂ labelled “.55”, one without abdomen, one lacking labial palpi, BMNH. Hudson 1928, pp. 167–168, pl. xx fig. 2–4, as *Crambus vittellus*. Gaskin 1975, pp. 339–341, fig. 10b, 17d, 22e, 76, and 77, as *Orocrambus vittellus*.

**nexalis** Walker, 1863b, p. 178 (*Crambus*). Synonymised by Meyrick (1883b, p. 27).

[Nelson NN], T.R. Oxley; LT ♀ selected by S. Bleszynski (Gaskin 1975, p. 339), BMNH.

Hudson 1928, p. 167, as synonym.

**transcissalis** Walker, 1863b, p. 178 (*Crambus*). Synonymised by Meyrick (1883b, p. 27).

[Auckland AK], A. Sinclair; HT ♂ unique, BMNH.

Hudson 1928, p. 167, as synonym.

**sublicellus** Zeller, 1863, p. 31 (*Crambus*). Synonymised by Meyrick (1883b, p. 27).

[New Zealand], lost; NT selected by S. Bleszynski (the same specimen as his NT for *vittellus*, but this disregarded by Gaskin 1975, p. 340), BMNH.

Hudson 1928, p. 167, as synonym.

**bisectellus** Zeller, 1863, p. 32 (*Crambus*). Synonymised by Meyrick (1883b, p. 27).

[New Zealand], lost; Castle Hill MC, E. Meyrick; NT ♂ selected by S. Bleszynski, labelled “Castle Hill, 18/1/83”, BMNH.

**incrassatellus** Zeller, 1863, p. 32 (*Crambus*). Synonymised by Meyrick (1883b, p. 27).

[New Zealand], Hugel; HT ♀ unique, abdomen missing, NHMW.

Hudson 1928, p. 167, as synonym.

**rapidus** Butler, 1877, p. 399 (*Crambus*). Synonymised by Meyrick (1883b, p. 27).

[Dunedin DN], J. Hector; ST ♀ labelled “77.34” (Gaskin 1975, p. 339), abdomen missing, BMNH.

Hudson 1928, p. 167, as synonym.

**conopias** Meyrick, 1907c, p. 109 (*Crambus*). Synonymised by Gaskin (1975, p. 339).

Ida Valley CO, J.H. Lewis; HT ♀ unique, BM genitalia slide no. Pyral 5634, BMNH.

Hudson 1928, p. 163, not figured, as *Crambus conopias*. Gaskin 1975, p. 339, as synonym.

Note. HT *conopias* bears the Meyrick label “Dunedin ... GVH 05”, but Hudson (1928, p. 163) points out the actual collector and locality.

**vulgaris** Butler, 1877, p. 400 (*Crambus*)

[?Dunedin DN or Castle Hill MC], J. Hector or J.D. Enys; HT ♂ designated by Butler, but see Gaskin (1975, p. 342, information at end of ‘Type Data’ section), BMNH.

Hudson 1928, pp. 168–169, pl. xx fig. 39, as *Crambus vulgaris*. Gaskin 1975, pp. 342–343, fig. 10c, 17e, 78, and 79, as *Orocrambus vulgaris*.

Note. Meyrick (1883b, p. 28; 1913b, p. 35) synonymised *vulgaris* with his concept of *tuhualis* Felder & Rogenhofer.

**obstructus** Meyrick, 1911a, p. 82 (*Crambus*). Synonymised by Gaskin (1975, p. 342).

[blown into train], Lumsden OL/SL, G.B. Longstaff; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 342), BMNH. Hudson 1928, p. 167, pl. xx fig. 5, as *Crambus obstructus*.

**xanthogrammus** Meyrick, §1882a, p. 187; 1883b, pp. 19 (key) and 32–33 (*Crambus*)

Lake Coleridge MC, R.W. Fereday; LT ♂ selected by Gaskin (1975, p. 343), BMNH.

Hudson 1928, p. 170, pl. xx fig. 6, as *Crambus xanthogrammus*. Gaskin 1975, pp. 343–344, fig. 10d, 17f, and 80, as *Orocrambus xanthogrammus*.

● **Tauroscopa** Meyrick, 1888c, p. 69. Type species *Tauroscopa gorgopis* Meyrick, by original monotypy.

**gorgopis** Meyrick, 1888c, pp. 69–70 (*Tauroscopa*)

Mount Arthur NN, G.V. Hudson; HT ♂ unique (Gaskin 1973, p. 445), BMNH.

Hudson 1928, p. 171, pl. xix fig. 42. Gaskin 1973, pp. 445–446, fig. 1, 8, and 13.

**howesi** Philpott, 1928a, p. 361 (*Tauroscopa*). Synonymised by Gaskin (1973, p. 445).

Obelisk, Old Man Range CO, W.G. Howes; HT ♂ designated by Philpott (Gaskin 1973, p. 445), NZAC.

Hudson 1939, p. 421, pl. lxii fig. 21, as species. Gaskin 1975, fig. 24.

**notabilis** Philpott, 1923, p. 149 (*Tauroscopa*)

Mount Peel NN, H. Hamilton; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 171, pl. I fig. 19. Gaskin 1973, fig. 9, 14, and 25.

**trapezitis** Meyrick, 1905, p. 227 (*Tauroscopa*)  
Mount Earnslaw OL, G.V. Hudson; HT ♂ unique (Gaskin 1973, p. 448), BMNH.  
Hudson 1928, p. 171, pl. xix fig. 41. Gaskin 1973, pp. 448–449, fig. 10, 15, and 26.

● **Tawhitia** Philpott, 1931, pp. 26–28. Type species *Tauroscopa glaucophanes* Meyrick, by original designation.

**Velasquez** Bleszynski, 1962, p. 14. Type species *Crambus pentadactylus* Zeller, by original designation.

**glaucophanes** Meyrick, 1907c, pp. 109–110  
(*Tauroscopa*)

Lake Wakatipu OL, G.V. Hudson; LT ♂ selected by S. Bleszynski (Gaskin 1975, p. 348), BMNH.

Hudson 1928, pp. 171–172, pl. xx fig. 45 and 46, pl. xlvi fig. 5. Gaskin 1975, pp. 348–349, fig. 22i, 26, 27♂, ♀, and 84–86.

**leonina** Philpott, 1931, p. 28 (*Tawhitia*). Synonymised by Gaskin (1975, p. 348).

Takitimu Mountains SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 421, pl. lxi fig. 15, as species.

**pentadactyla** Zeller, 1863, p. 32 (*Crambus*)

[Christchurch MC], H.G. Knaggs; HT ♂ unique, BMNH. Hudson 1928, pp. 170–171, pl. xx fig. 48, as *Argyria pentadactyla*. Gaskin 1975, pp. 349–350, fig. 28♂, ♀ and 87, as *Tawhitia pentadactyla*.

**claviferella** Walker, 1886b, pp. 1546–1547  
(*Aquita*). Synonymised by Meyrick (1913, p. 35). Tasmania, Allport; HT ♂ unique, BMNH.

Hudson 1928, p. 170, as synonym.

**strigosus** Butler, 1877, pp. 398–399 (*Aphomia*). Synonymised (with *claviferella*) by Meyrick (1888c, p. 69).

[Dunedin DN], J. Hector; HT ♂ ?unique, BMNH.

Hudson 1928, p. 170, as synonym.

—♂—

## Subfamily MUSOTIMINAE (in the sense of Minet 1981, p. 269)

● **Musotima** Meyrick, 1884c, pp. 288–289. Type species *Diathrausta aduncalis* Felder & Rogenhofer, by subsequent designation (Hampson 1896, p. 199).

**aduncalis** Felder & Rogenhofer, 1875, pl. cxxxv fig. 11 (*Diathrausta?*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, head and abdomen missing, BMNH.

Hudson 1928, p. 177, pl. xix fig. 19 and 20, as *Musotima aduncalis*, after Meyrick (1884e, p. 289).

**nitidalis** Walker, 1866a, pp. 1317–1318 (*Isopteryx*) “Australia”, Damel; HT ♂ unique, forewings and head missing, abdomen in gelatin capsule, BMNH.

Hudson 1928, pp. 177–178, pl. xix fig. 18, as *Musotima nitidalis*, after Meyrick (1884e, p. 290).

**timarialis** Felder & Rogenhofer, 1875, pl. cxxxv fig. 23 (*Diathrausta*). Synonymised by Meyrick (1884e, p. 290).

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen in gelatin capsule, BMNH.

Hudson 1928, p. 177, as synonym.

Also 1 Australian species establishing sporadically in AK.

—♂—

## Subfamily NYMPHULINAE (in the sense of Minet 1985a, p. 84)

● **Hygraula** Meyrick, §1885a, p. 347; 1885f, p. 129. Type species *Paraponyx nitens* Butler, by original monotypy.

Note. Lange (1956, pp. 65, 66, 68, and 94) regards *Hygraula* as separate from *Paraponyx* [sic] Hübner (type species *Tinea stratiotata* Linnaeus, by subsequent designation of Guenée 1854, p. 269), but showing a probable relationship.


**nitens** Butler, 1880, p. 556 (*Paraponyx*) Blenheim MB, W. Skellon; HT ♂ designated by Butler, abdomen missing, BMNH.

Hudson 1928, p. 177, pl. xix fig. 22, as *Nymphula nitens*, following Meyrick (1913b, p. 38).

Note. Lange (1956, fig. 111 and 139) illustrates ♂ and ♀ genitalia.

—♂—

**Subfamily PYRAUSTINAE**  
(in the sense of Munroe 1972, pp. 12–13)



**Pyralidae: Pyraustinae**  
(175) *Mnesictena flavidalis* (Doubleday)

• **Clepsicosma** Meyrick, 1888c, pp. 63–64. Type species *Clepsicosma iridia* Meyrick, by original monotypy.

Note. Fletcher & Nye (1984, p. 35) place *Clepsicosma* in Nymphulinae.

**iridia** Meyrick, 1888c, p. 64 (*Clepsicosma*)  
Waitakere Range AK, E. Meyrick; HT ♀ unique, BMNH.  
Hudson 1928, pp. 204–205, pl. xxiv fig. 22.

• **Deana** Butler, 1879b, p. 451, new name for *Adena* Walker, preoccupied by *Adena* Agassiz, 1846 (emendation of *Hadena* Schrank, Noctuidae).

**Adena** Walker, 1883b, pp. 1976–198. Type species *Adena xanthialis* Walker, by original monotypy.  
Note. Despite Meyrick (1913b, p. 40), *Deana* is strictly available, and is not a homonym of *Deanea* Reichenbach (Aves).

**Nesarcha** Meyrick, 1884e, p. 330. Type species *Scopula hybreasalis* Walker (as *hybrealis*), by original monotypy.

**hybreasalis** Walker, 1859b, p. 797 (*Scopula*?)  
[?Wellington WN], W. Parry; HT ♀ unique, with accession label “50-71”, BMNH.

Hudson 1928, p. 179, pl. xxi fig. 30 and 31, as *Nesarcha hybrealis*.

**paronalis** Walker, 1859b, pp. 797–798 (*Scopula*?). Synonymised by Meyrick (1884e, p. 330).  
[?Hawkes Bay HB], W. Colenso; LT ♂ here designated, labelled “New Zeal. 53-19”, “52. *Scopula paronalis*”, “Type”, posterior half of abdomen in gelatin capsule, BMNH.

Hudson 1928, p. 179, as synonym.

**xanthialis** Walker, 1863b, p. 198 (*Adena*). Synonymised by Meyrick (1884e, p. 330).  
[Nelson NN], T.R. Oxley; HT ♀ unique, BMNH.  
Hudson 1928, p. 179, as synonym.

• **Diasemia** Hübner, [1825] 1816, p. 348. Type species *Phalaena litterata* Scopoli, 1763, p. 229.

**grammalis** Doubleday, 1843, p. 287 (*Diasemia*)  
[Auckland AK, A. Sinclair]; HT ♂ (so labelled) with accession label “42.55 New Zealand”, “k”, BMNH.  
Hudson 1928, p. 178, pl. xix fig. 26.

• **Heliothela** Guenée, 1854, p. 152, in the sense of Meyrick. Type species *Phalaena atralis* Hübner, [1788] 1786–89, by original monotypy.

Note. *Phalaena atralis* Hübner, [1788] is a junior primary homonym of *P. atralis* Fabricius, 1755. The objective replacement name is *Heliothis huebneri* Kocak, 1980 (Fletcher & Nye 1984, p. 67).

**atra** Butler, 1877, p. 404 (*Orosana*) new combination  
Canterbury Plains MC, J.D. Enys; HT ♂ so labelled, Brit.

Mus. Pyralidae slide no. 3697, BMNH.  
Hudson 1928, p. 182, pl. xix fig. 29, as *Heliothela erebopis*, following Meyrick (1913b, p. 41).

**erebopis** Meyrick, 1913b, p. 41 (*Heliothela*). New synonymy.

Name proposed by Meyrick to avoid confusion with *atralis* Hübner; HT therefore Butler's type specimen.

Hudson 1928, p. 182, as valid name.

Note. The epithet *atra* is sufficiently distinct from *atralis* for it not to be regarded as a homonym, and there is no need for a new name.

• **Hellula** Guenée, 1854, pp. 415–416. Type species *Phalaena undalis* Fabricius, 1781, by original monotypy; “Italia”.

Note. *Hellula* is placed in Glaphyriinae by Munroe (1964, p. 1261).

**hydralis** Guenée, 1854, p. 416 (*Hellula*)

Type locality unknown to Guenée; HT not examined.  
Hudson 1939, p. 422, pl. lxi fig. 9, as *Hellula undalis*.

Note. New Zealand records are of vagrant specimens from eastern Australian populations. Meyrick (1934, p. 152) gives a record from near Greymouth BR in 1932, as *H. undalis*.

• **Hymenia** Hübner, [1825] 1816, p. 361. Type species *Pyralis perspectalis* Hübner, 1796, by subsequent designation of Moore ([1885] 1884–1887, p. 293) (Fletcher & Nye 1984, p. 73; cf. Meyrick 1913b, p. 39); “Ceylon” (Sri Lanka).

**recurvalis** Fabricius, 1775, p. 644 (*Phalaena*)  
“East Indies”.

New Zealand: adventive; recurrent immigrant, occasionally establishing.

Hudson 1928, p. 179, pl. xx fig. 41, as *Hymenia fascialis* Cramer.

Note. Clarke (1971, pp. 69–70) gives full references (under *Spoladea* Guenée). Walker (1859a, p. 396) lists 3 New Zealand records from Bolton and Sinclair.

- ***Loxostege*** Hübner, [1825] 1816, p. 352. Type species *Pyralis aeruginalis* Hübner, 1796, by subsequent designation (Hampson 1918b, p. 189).

***Proternia*** Meyrick, 1884e, pp. 292 (key) and 317. Type species *Proternia philocapna* Meyrick, by original monotypy; placement as in BMNH collection.

***Proteroeca*** Meyrick, 1884e, pp. 292 (key) and 335. Type species *Proteroeca comastis* Meyrick, by original monotypy; placement as in BMNH collection.

***affinitalis*** Lederer, 1863, p. 475, pl. 12 fig. 4 (*Botys*)  
Type locality Australia; HT not examined.  
New Zealand: adventive; established since 1970 (AK, HB).

***comastis*** Meyrick, 1884e, p. 335 (*Proteroeca*)  
Castle Hill MC, E. Meyrick; LT ♂ here designated,  
labelled “Castle Hill New Zealand 2400 ft 18/1/83”,  
“Proteroeca comastis” Meyr. 1/9 E. Meyrick det. in  
Meyrick Coll.”, BMNH.

Hudson 1928, p. 182, pl. xix fig. 25.  
Note. Meyrick (1885j, p. 456) gives locality information  
omitted in the original description.

***intrudens*** Warren, 1892, p. 175 (*Ennychia*).  
Synonymised by M. Shaffer, BMNH.  
Castle Hill MC, E. Meyrick; HT ♀ (ex Raynor Collection)  
selected by M. Shaffer, BMNH.  
Not mentioned by Hudson.

***philocapna*** Meyrick, 1884e, pp. 317–318 (*Proternia*)  
Hamilton WO, E. Meyrick; LT ♀ here designated, labelled  
“Hamilton New Zealand at light 17/1/80”, “Proternia  
philocapna” Meyr. 1/6 E. Meyrick det. in Meyrick  
Coll.”, BMNH.  
Hudson 1928, p. 179, pl. xx fig. 42.

- ***Mnesictena*** Meyrick, 1884e, pp. 293 (key) and 328–329. Type species *Mnesictena marmarina* Meyrick, by subsequent designation (Hampson 1899, p. 231, as *M. quadralis*).  
Note. Fletcher & Nye (1984, p. 95) give details of  
Hampson's actions.

***Mecyna*** in the sense of Meyrick (1913b, p. 40).

***adversa*** Philpott, 1917b, p. 243 (*Mecyna*) new  
combination

Queenstown OL, W.G. Howes; HT ♂ not located, AT ♀  
designated by Philpott, NZAC.  
Hudson 1928, p. 181, pl. xxi fig. 24, as *Mecyna adversa*.

***antipodea*** Salmon in Salmon & Bradley, 1956, p.  
77, fig. 21 (*Mecyna*) new combination  
Antipodes Islands, E.G. Turbott; HT ♂ unique, AMNZ.

***daiclesalis*** Walker, 1859c, p. 1017 (*Scopula*?)  
[?Auckland AK], J.F. Churton; HT ♀ abdomen and right  
forewing missing, BMNH.  
Hudson 1928, p. 180, pl. xxi fig. 23, as *Mecyna daiclesalis*  
(name emended by Meyrick 1889, p. 155), after Meyr-  
ick (1913b, p. 40).

***otagalis*** Felder & Rogenhofer, 1875, pl. cxxxiv  
fig. 35 (*Botys*). New synonymy.  
[?Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen in  
gelatin capsule, BMNH.  
Hudson 1928, p. 181, as synonym of *Mecyna flavidalis*.  
Note. *Botys otagalis* is obviously synonymous with Walk-  
er's *daiclesalis*; the yellowish flush at the base of the  
forewing is not uncommon in *daiclesalis* populations.

***flavidalis*** Doubleday, 1843, p. 287 (*Margaritia*);  
Meyrick, 1884c, p. 330, as *Mnesictena flavidalis*  
[Auckland AK], A. Sinclair; HT ♂ (labelled as HT by  
W.H.T. Tams) labelled “Margaritia flavidalis Dou-  
bleday TYPE”, “42.55 New Zealand”, abdomen miss-  
ing, BMNH.

Hudson 1928, p. 181, pl. xxi fig. 27 and 28, as *Mecyna*  
*flavidalis*, after Meyrick (1913b, p. 40).

***quadralis*** Doubleday, 1843, p. 288 (*Margaritia*).  
Synonymised by Meyrick (1884e, p. 330).  
[Auckland AK], A. Sinclair; HT ♀ (labelled as HT by  
W.H.T. Tams) here designated, labelled “Margaritia  
quadralis Doubleday TYPE”, “New Zealand 42.55”,  
accompanied by 2 PLT ♀♀ labelled by D.J. Carter as  
ST ♀ ♀ of *flavidalis*, BMNH.  
Hudson 1928, p. 181, as synonym of *flavidalis*.

***dipsasalis*** Walker, 1859b, pp. 796–797 (*Scop-  
ula*?). Synonymised by Meyrick (1884e, p. 330).  
[Auckland AK], A. Sinclair; HT ♂ unique, accession no.  
45.61, BMNH.  
Hudson 1928, p. 181, as synonym of *flavidalis*.

***marmorina*** Meyrick, 1884e, p. 329 (*Mnesictena*)  
Palmerston North WI/WN, E. Meyrick; LT ♂ selected  
by M. Shaffer, labelled “Palmerston New Zealand  
4/3/85”, “Mnesictena marmorina Meyrick 1/9 E.  
Meyrick det. in Meyrick Coll.”, BMNH.  
Hudson 1928, pp. 181–182, pl. xxi fig. 26, as *Mecyna*  
*marmorina*, after Meyrick (1913b, p. 40).

***notata*** Butler, 1879a, p. 493 (*Scopula*); Meyrick,  
1884, p. 330, as *Mnesictena notata*  
[Dunedin DNJ], F.W. Hutton; HT ♂ unique, accession  
no. 79.9, BMNH.  
Hudson 1928, p. 181, pl. xliv fig. 21, as *Mecyna notata*,  
after Meyrick (1913b, p. 40).

***pantheropa*** Meyrick, 1902c, p. 277 (*Mecyna*) new  
combination  
Chatham Island, J. Fougeré; LT ♂ designated by M. Shaf-  
fer, BMNH.  
Hudson 1928, p. 181, pl. xxi fig. 29, as *Mecyna pantheropa*.

• ***Sceliodes*** Guenée 1854, p. 400. Type species *Sce-  
liodes mucidalis* Guenée, by original monotypy.

» Pyralidae, *Sceliodes*

**Eretria** Snellen, 1880, p. 206. Type species *Eretria obsistalis* Snellen, by original monotypy.  
Note. Fletcher & Nye (1984, p. 55) record *Eretria* Snellen as a junior homonym of *Eretria* Robineau-Desvoidy, 1863 (Diptera).

**cordalis** Doubleday, 1843, p. 288 (*Margaritia*)  
[Auckland AK], A. Sinclair; ST ♂ so labelled by D.J. Carter, accession no. 42.55, BMNH.  
Hudson 1928, p. 178, pl. xx fig. 47.

**mucidalis** Guenée, 1854, p. 400 (*Sceliodes*). Synonymised by Meyrick (1884e, p. 303).  
“Australia”, ?collector; HT not examined, MNHN.  
Hudson 1928, p. 178, as synonym of *cordalis*.

**extensalis** Walker, 1866a, p. 1311 (*Daraba*). Synonymised by Meyrick (1884e, p. 303).  
[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1928, p. 178, as synonym of *cordalis*.

**obsistalis** Snellen, 1880, p. 206 (*Eretria*). Synonymised by Meyrick (1884e, p. 303).  
Boelekambo, Bothmian (Sulawesi), M.C. Piepers; type material apparently lost.  
Hudson 1928, p. 178, as synonym of *cordalis*.


• **Uresiphita** Hübner, [1825] 1816, p. 353. Type species *Pyralis limbalis* [Denis & Schiffermüller], in the sense of Hübner [1825] (= *Pyralis polygonalis* [Denis & Schiffermüller], 1775, by subsequent designation (Pierce & Metcalfe 1938, p. 64)); Europe.  
Note. Fletcher & Nye (1984, p. 157) note this as a case of a misidentified type species, and have referred it to the ICZN.

**Mecyna** Doubleday, [1849] 1850, p. 14. Type species *Pyralis asinalis* Hübner, [1819] 1796, pl. 29 fig. 185, by monotypy.  
Note. Fletcher & Nye (1984, p. 88) give details on the usage of *Mecyna*.

**polygonalis maorialis** Felder & Rogenhofer, 1875, pl. cxxxiv fig. 34 (*Botys*)  
[Nelson NN, T.R. Oxley]; HT ♀ unique, abdomen in gelatin capsule, BMNH.  
Hudson 1928, p. 180, pl. xxi fig. 25, as *Mecyna maorialis*.  
Clarke 1971, p. 75, as *maorialis*, by implication as subspecies of *polygonalis*.  
Note. *Mecyna deprivalis* Walker, 1859, p. 806, was listed by Meyrick (1884e, p. 326) as the senior synonym of *maorialis*. *M. deprivalis* was listed by Walker as from “Ceylon”. Meyrick (1913b, p. 40) removed *deprivalis* from the New Zealand list by implication. Walker (1859c, p. 807), under *Mecyna ornithopteralis* Guenée, noted “var. minor; alae posticae margine angustiore”, “J. New Zealand. Presented by Dr Sinclair” [Auckland]. The specimen is *U. p. maorialis*.

—⊗—

Subfamily SCOPARIINAE  
(in the sense of Munroe 1972, pp. 14–17)



Pyralidae: Scopariinae  
(176) *Eudonia diphtheralis* (Walker)

• **Antiscopa** Munroe, 1964, pp. 261 (key) and 265–266. Type species *Scoparia epicomia* Meyrick, by original designation.

**acompa** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 72 (key) and 100 (*Scoparia*)  
Lake Wakatipu OL, E. Meyrick; HT ♂, BM Pyral. slide no. 3673 ♂, BMNH.  
Hudson 1928, p. 195, pl. xlvi fig. 6, as *Scoparia acompa*.  
Munroe 1964, p. 266, as *Antiscopa acompa*.

**elaphra** Meyrick, §1884d, p. 236; 1885e, pp. 72 (key) and 105 (*Scoparia*)  
Palmerston [North] WI/WN, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3686 ♂, BMNH.  
Hudson 1928, p. 197, pl. xxii fig. 5, as *Scoparia elaphra*.  
Munroe 1964, p. 265, as *Antiscopa elaphra*.

**epicomia** Meyrick, §1884d, p. 236; 1885e, pp. 72 (key) and 99 (*Scoparia*)  
Dunedin DN, A. Purdie; LT ♂ selected by E. Munroe, BMNH.  
Hudson 1928, p. 195, pl. xxi fig. 33, as *Scoparia epicomia*.

• **Eudonia** Billberg, 1820, p. 93. Type species *Tinea mercurella* Linnaeus, 1758, p. 902, by original monotypy (listed as *mercuriella* Linnaeus); Europe.

**Witlesia** Chapman, 1912, p. 507. Type species *Eudorea pallida* Curtis, 1827 (as *pallida* Stephens), by original monotypy; Europe. Synonymised by Munroe (1972, p. 47).

Note. Species included here have the ♂ valva lacking a process on the ventral margin (see Munroe 1964, p. 266; 1972, p. 48).

**alopecias** Meyrick, 1901, p. 570 (*Scoparia*) new combination

Mount Cook MK, R.W. Fereday; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3672 ♂, BMNH.  
Hudson 1928, p. 196, not figured, as *Scoparia alopecias*.

**aspidota** Meyrick, §1884d, p. 237; 1885e, pp. 110 (key) and 115–116 (*Xeroscopa*) new combination  
[Ben Lomond] OL, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3643 ♂, BMNH.  
Hudson 1928, p. 201, pl. xxiv fig. 10, as *Scoparia aspidota*, after Meyrick (1913b, p. 45).

**asterisca** Meyrick, §1884d, p. 237 (no description); 1885e, pp. 110 (key) and 118–119 (*Xeroscopa*) new combination  
Lake Wakatipu OL, R.W. Fereday; LT ♂ labelled as HT, "L. Wakatipu New Zealand RWF 11/181", BM Pyral. slide no. 3700 ♂, BMNH.  
Hudson 1928, p. 204, pl. xxi fig. 47, as *Scoparia asterisca*, after Meyrick (1913b, p. 45).

**atmogramma** Meyrick, 1915a, pp. 202–203 (*Scoparia*) new combination  
?Invercargill SL, A. Philpott; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3670 ♂, BMNH.  
Hudson 1928, p. 197, pl. xlvi fig. 25, as *Scoparia atmogramma*.

**axena** Meyrick, §1884d, p. 236; 1885e, pp. 73 (key) and 103 (*Scoparia*) new combination  
Arthur's Pass NC/WD, 4500 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3628 ♂, BMNH.  
Hudson 1928, p. 196, pl. xliv fig. 22, as *Scoparia axena*.

**bisinialis** Hudson, 1928, p. 194 (*Scoparia*) new combination  
Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "271a", NMNZ.  
Hudson 1928, p. 194, pl. xxi fig. 45.

**cataxesta** Meyrick, §1884d, p. 236; 1885e, pp. 73 (key) and 96–97 (*Scoparia*) new combination  
Castle Hill MC, 3000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3660 ♂, BMNH.  
Hudson 1928, p. 193, pl. xxi fig. 19, as *Scoparia cataxesta*.

**chalara** Meyrick, 1901, p. 570 (*Scoparia*) new combination  
Mount Cook MK, G.V. Hudson; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3644 ♂, BMNH.  
Hudson 1928, p. 204, pl. xxiv fig. 41, as *Scoparia chalara*.

**characta** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 72 (key) and 90 (*Scoparia*) new combination  
Makatoku WA, E. Meyrick; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3645 ♂, BMNH.  
Hudson 1928, p. 189, pl. xxii fig. 4, as *Scoparia characta*.

**chlamydota** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 72 (key) and 82 (*Scoparia*) new combination  
Arthur's Pass NC/WD, 3000 ft, E. Meyrick; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3727 ♂, BMNH.

Hudson 1928, p. 184, pl. xix fig. 28, as *Scoparia chlamydota*.

**choristis** Meyrick, 1907c, p. 111 (*Scoparia*) new combination  
Kaitoke WN, G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3656 ♂, BMNH.  
Hudson 1928, pp. 191–192, pl. xxi fig. 2, as *Scoparia choristis*.

**colpota** Meyrick, 1888c, pp. 65–66 (*Scoparia*) new combination  
Wellington WN, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3655 ♂, BMNH.  
Hudson 1928, p. 191, pl. xxi fig. 1, as *Scoparia colpota*.

**critica** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 72 (key) and 88 (*Scoparia*) new combination  
Arthur's Pass NC/WD, 3,000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3743, BMNH.  
Hudson 1928, p. 189, pl. xxii fig. 42, as *Scoparia critica*.

**crypsinoa** Meyrick, §1884d, p. 236 (no description, as *cryspsioa*); 1885e, pp. 72 (key) and 102–103 (*Scoparia*) new combination  
Castle Hill MC, 3000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3625, BMNH.

Hudson 1928, p. 196, pl. xxii fig. 9 (dubious), as *Scoparia crypsinoa*.

**agana** Meyrick, 1912c, p. 119 (*Scoparia*). Synonymised by Meyrick (1921, p. 334).  
Lake Wakatipu OL, G.V. Hudson; LT ♂ selected by E. Munroe, BMNH.  
Hudson 1928, p. 119, pl. xxii fig. 9, as *crypsinoa*, as synonym.

**cymatias** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 72 (key) and 86–87 (*Scoparia*) new combination  
Arthur's Pass NC, 2500 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3734, BMNH.

Hudson 1928, p. 187, pl. xxii fig. 40.

**cryptastis** Meyrick, 1909a, p. 7 (*Scoparia*) new combination  
Invercargill SL, A. Philpott; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3674 ♂, BMNH.  
Hudson 1928, pp. 195–196, pl. xxii fig. 30.

**deltophora** Meyrick, §1884d, p. 236; 1885e, pp. 74 (key) and 106 (*Scoparia*) new combination  
Arthur's Pass NC/WD, 4000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3630 ♂, BMNH.  
Hudson 1928, pp. 197–198, pl. xxi fig. 39, as *Scoparia deltophora*.

**dinodes** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 73 (key) and 85 (*Scoparia*) new combination

» Pyralidae, *Eudonia dinodes*

[Riccarton Bush], Christchurch MC, E. Meyrick; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3733 ♂, BMNH.  
Hudson 1928, pp. 185–186, pl. xxii fig. 3, as *Scoparia dinodes*.

**dochmia** Meyrick, 1905, pp. 229–230 (*Scoparia*) new combination

Lake Wakatipu OL, G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3720 ♂, BMNH.  
Hudson 1928, p. 184, pl. xxii fig. 2.

**epicremna** Meyrick, §1884d, p. 237 (no description); 1885e, pp. 110 (key) and 117 (*Xeroscopa*) new combination

Castle Hill MC, 2500 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3702, BMNH.  
Hudson 1928, p. 203, pl. xlvi fig. 4 (dubious), as *Scoparia epicremna*.

**feredayi** Knaggs, 1867, p. 80 (*Scoparia*) new combination

Rakaia MC, H.G. Knaggs; HT ♂ unique, accession no. 71.30, BM Pyral. slide no. 3616, BMNH.

Hudson 1928, p. 195, pl. xxi fig. 34, as *Scoparia feredayi*.

**moanalis** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 34 (*Scoparia*). Synonymised by Meyrick (§1884d, p. 236; 1885e, p. 100).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1928, p. 195, as synonym.

**gressitti** Munroe, 1964, p. 269 (*Witlesia*) new combination

Tucker Cove, Campbell Island, J.L. Gressitt; HT ♂ designated by E. Munroe, [still in CNCI as at March 1987] NZAC.

Munroe 1964, pp. 269–270, fig. 4 (♂ genitalia).

**gyrotoma** Meyrick, 1909a, p. 7 (*Scoparia*) new combination

Lake Tekapo MK, G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3669 ♂, BMNH.

Hudson 1928, p. 193, pl. xxi fig. 18 (a poor depiction); 1939, p. 424, pl. lvi fig. 27; as *Scoparia gyrotoma*.

**repercussa** Philpott, 1929a, pp. 301–302 (*Scoparia*). Synonymised by Philpott (1931, p. 29).

Lake Tekapo MK, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 424, pl. lvi fig. 27, as synonym.

**hemicycla** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 73 (key) and 87 (*Scoparia*) new combination

Arthur's Pass NC/WD, 3000 ft, E. Meyrick; HT ♂ unique, BM Pyral. slide no. 3738 ♂, BMNH.

Hudson 1928, p. 187, pl. xxii fig. 7, as *Scoparia hemicycla*.

**hemiplaca** Meyrick, 1889b, pp. 155–156 (*Scoparia*)

new combination

Wellington WN, G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3729 ♂, BMNH.

Hudson 1928, p. 184, pl. xxii fig. 11, as *Scoparia hemiplaca*.

**legnota** Meyrick, §1884d, p. 237 (no description); 1885e, pp. 110 (key) and 117–118 (*Xeroscopa*) new combination

Lake Wakatipu OL, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3684, BMNH.

Hudson 1928, p. 203, pl. xxi fig. 49, as *Scoparia legnota*.

**leptalaea** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 73 (key) and 98–99 (*Scoparia*) new combination

[heath-like scrub and swamp], Hamilton WO, E. Meyrick; LT ♂ selected by E. Munroe, BMNH.

Hudson 1928, p. 194, not figured, as *Scoparia leptalaea*.

**leptophaea** Meyrick, 1902c, pp. 277–278 (*Scoparia*). Synonymised by Hudson (1928, p. 194).

Chatham Island, J. Fourgère; LT ♀ selected by E. Munroe, BM Pyral. slide no. 3668 ♀, BMNH.

Hudson 1928, p. 194, as synonym.

**leucogramma** Meyrick, §1884d, p. 237; 1885e, pp. 110 (key) and 119 (*Xeroscopa*) new combination

Mount Hutt MC, R.W. Fereday; LT ♂ selected by E. Munroe, BMNH.

Hudson 1928, p. 204, pl. xxii fig. 41, as *Scoparia leucogramma*.

Note. The ♂ genitalia on BM Pyral. slide no. 3706 are of an Otira WD specimen.

**linealis** Walker, 1866a, pp. 1503–1504 (*Scoparia*) new combination

[Nelson NN], T.R. Oxley; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 192, as synonym of *Scoparia submarginalis*.

**locularis** Meyrick, 1912c, pp. 118–119 (*Scoparia*) new combination

Lake Wakatipu OL, G.V. Hudson; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3653 ♂, BMNH

Hudson 1928, pp. 190–191, pl. xxi fig. 15.

**luminatrix** Meyrick, 1909a, p. 8 (*Scoparia*) new combination

Invercargill SL, A. Philpott; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3699 ♂, BMNH.

Hudson 1928, p. 203, pl. xxi fig. 43 (a poor likeness), as *Scoparia luminatrix*.

**manganeutis** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 72 (key) and 102 (*Scoparia*) new combination

Otira Gorge WD, 1600 ft, E. Meyrick; LT ♂ selected by

E. Munroe, head missing, BM Pyral. slide no. 3679 ♂, BMNH.  
Hudson 1928, p. 196, not figured, as *Scoparia manganeutis*.

**melanaegis** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 73 (key) and 92 (*Scoparia*) new combination

Arthur's Pass NC, 2600 ft, E. Meyrick; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3649 ♂, BMNH.

Hudson 1928, p. 190, pl. xxi fig. 51, as *Scoparia melanaegis*.

**meliturga** Meyrick, 1905, p. 228 (*Scoparia*) new combination

Wellington WN, E. Meyrick; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3725 ♂, BMNH.  
Hudson 1928, pp. 183–184, pl. xxii fig. 12, as *Scoparia meliturga*.

**microphthalmia** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 72 (key) and 87 (*Scoparia*) new combination

Lake Wakatipu OL, 1000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3732 ♂, BMNH.  
Hudson 1928, p. 187 (but not pl. 1 fig. 8), as *Scoparia microphthalmia*.

**minualis** Walker, 1866a, p. 1504 (*Scoparia*) new combination

[Auckland AK], D. Bolton; HT ♀ so labelled, BM Pyral. slide no. 3750, BMNH.

Hudson 1928, p. 185, pl. xxii fig. 37 (a poor likeness), as "Scoparia minualis Meye." (cf. Meyrick 1913b, p. 42, correct citation).

**chimeria** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 72 (key) and 84 (*Scoparia*). New synonymy.

Masterton WA, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3731, BMNH.

Hudson 1928, p. 185, pl. xxii fig. 43 (a poor likeness), as *Scoparia chimeria*.

Note. HT *minualis* and LT *chimeria* are identical in colour pattern. Neither resembles Hudson's figures.

**octophora** Meyrick, §1884d, p. 237 (no description); 1885e, pp. 110 (key) and 118 (*Xeroscopa*) new combination

Castle Hill MC, 3000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3702, BMNH.

Hudson 1928, pp. 203–204, pl. xxiv fig. 21 (very dubious), as *Scoparia octophora*.

**oculata** Philpott, 1927d, pp. 32–83 (*Scoparia*) new combination

Nelson NN, A. Philpott; HT ♀ designated by Philpott, NZAC.

Hudson 1939, p. 427, pl. lvi fig. 16, as *Scoparia oculata*.

Note. A topotypic PT ♂ in BMNH (BM Pyral. slide no. 3701) has the characteristic *Eudonia* valvae.

**oreas** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 72 (key) and 81 (*Scoparia*) new combination

[Ben Lomond], Lake Wakatipu OL, E. Meyrick; HT ♂ unique, BM Pyral. slide no. 3724, BMNH.  
Hudson 1928, p. 183, not figured, as *Scoparia oreas*.

**organaea** Meyrick, 1901, p. 569 (*Scoparia*) new combination

Mount Cook MK, G.V. Hudson; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3703 ♂, BMNH.

Hudson 1928, pp. 202–203, pl. xxii fig. 27, as *Scoparia organaea*.

**pachyerga** Meyrick, 1927a, p. 697 (*Scoparia*) new combination

Mount Holdsworth WN, 2500 ft, G.V. Hudson; HT ♂ unique (altitude given on label as "4000 ft"), BM Pyral. slide no. 3627, BMNH.

Hudson 1928, pp. 196–197, pl. lii fig. 9, as *Scoparia pachyerga* (corroborates collecting altitude of c. 2000 ft).  
Munroe 1964, p. 266, as *Witlesia pachyerga*.

**paltomacha** Meyrick, §1884d, p. 236; 1885e, pp. 74 (key) and 105–106 (*Scoparia*) new combination

Castle Hill MC, 2500 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3629 ♂, BMNH.

Hudson 1928, p. 197, pl. xxi fig. 36 and 37, as *Scoparia paltomacha*. Munroe 1964, p. 266, as *Witlesia paltomacha*.

**periphanes** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 74 (key) and 94 (*Scoparia*) new combination

Lake Wakatipu OL, R.W. Fereday; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3657 ♂, BMNH.

Hudson 1928, p. 191, pl. xxi fig. 3.

**philerga** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 73 (key) and 81–82 (*Scoparia*) new combination

Lake Wakatipu OL, 1000 ft, E. Meyrick; HT ♂ (?LT; see Meyrick's description), BM Pyral. slide no. 3726, BMNH.

Hudson 1928, p. 183, pl. xxi fig. 12, as *Scoparia philerga*.

**philetaera** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 72 (key) and 93–94 (*Scoparia*) new combination

Bealey River NC, E. Meyrick; HT ♂ unique, BM Pyral. slide no. 3652, BMNH.

Hudson 1928, p. 190, not figured, as *Scoparia philetaera*.

**pongalis** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 33 (*Scoparia*) new combination

[Nelson NN, T.R. Oxley]; HT ♂ unique, BM Pyral. slide no. 3648, BMNH.

Hudson 1928, pp. 189–190, pl. xxii fig. 1.

> Pyralidae, *Eudonia*

***psammitis psammitis*** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 73 (key) and 99 (*Scoparia*; as species) new combination

Arthur's Pass NC/WD, 4500 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3667 ♂, BMNH. Hudson 1928, p. 194, pl. xxiv fig. 51, as *Scoparia psammittis* (sic). Munroe 1964, p. 266, as *Witlesia psammittis*.

***psammitis campbellensis*** Munroe, 1964, pp. 267–269 (*Witlesia*) new combination

Tucker Cove, Campbell Island, J.L. Gressitt; HT ♂ designated by E. Munroe, [still in CNCI as at March 1987] NZAC.

Munroe 1964, pp. 267–269, fig. 3 (♂ genitalia).

***quaestoria*** Meyrick, 1929, p. 487 (*Scoparia*) new combination

Waitati DN, C.E. Clarke; LT ♂ (labelled as HT, but see Meyrick's description), BM Pyral. slide no. 3651 ♂, BMNH.

Hudson 1939, p. 424, pl. lvi fig. 7, as *Scoparia quaestoria*.

***rakaiaensis*** Knaggs, 1867, p. 80 (*Scoparia*) new combination

Rakaia MC, H.G. Knaggs; HT ♂ unique, BM Pyral. slide no. 3746 ♂, BMNH.

Hudson 1928, pp. 193–194, pl. xxi fig. 10, as synonym of *Scoparia indistinctalis* not of Walker but in the sense of Meyrick (1885e, p. 97).

***sabulosella*** Walker, 1863b, p. 178 (as *Crambus sabulosellus*) new combination

[Auckland AK], A. Sinclair; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3613 ♂, BMNH.

Hudson 1928, p. 198, pl. xxi fig. 38, as *Scoparia sabulosella*. Munroe 1964, p. 266, as *Witlesia sabulosella*.

Note. LT ♂ genitalia on slide no. 3613 differ markedly from those of a ♂ (locality unspecified, coll. G.F. Mathew) on slide no. 3622, suggesting that two species may be confused under the name *sabulosella*.

***steropaea*** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 74 (key) and 103–104 (*Scoparia*) new combination

Castle Hill MC, 3000 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3680 ♂, BMNH.

Hudson 1928, p. 197, pl. xxii fig. 8, as *Scoparia steropaea*.

***subditella*** Walker, 1866b, p. 1720 (*Nephopteryx*) new combination

[Auckland AK], D. Bolton; LT ♂ labelled as HT, BM Pyral. slide no. 3751 ♂, BMNH.

Not mentioned by Hudson.

***submarginalis*** Walker, 1863b, p. 48 (*Hypochalcia*) new combination

[Auckland AK], A. Sinclair; LT ♀ labelled as HT, BM Pyral. slide no. 3748 ♀, BMNH.

Hudson 1928, p. 192, pl. xxi fig. 7–9, as *Scoparia submarginalis*, after Meyrick (§1884d, p. 236; 1885e, p. 95).

***maoriella*** Walker, 1866b, p. 1720 (*Nephopteryx*). Synonymised by Meyrick (§1884d, p. 236; 1885e, p. 95).

[Auckland AK], D. Bolton; HT ♂ unique, BM Pyral. slide no. 3747 ♂, BMNH.

Hudson 1928, p. 192, as synonym.

***thyridias*** Meyrick, 1905, p. 229 (*Scoparia*) new combination

"New Zealand", G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3737 ♂, BMNH.

Hudson 1928, p. 183, pl. xxii fig. 36, as *Scoparia thyridias*.

***torodes*** Meyrick, 1901, p. 568 (*Scoparia*) new combination

Mount Cook MK, R.W. Fereday; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3654 ♂, BMNH.

Hudson 1928, p. 191, not figured, as *Scoparia torodes*.

***galactilis*** Hudson, 1913, p. 250 (*Scoparia*). Synonymised by Hudson (1939, p. 424).

Routeburn Valley OL, G.V. Hudson; HT ♂ unique, NMNZ.

Hudson 1928, p. 190, pl. xxi fig. 46.

***triclera*** Meyrick, 1905, p. 230 (*Scoparia*) new combination

Wellington WN, G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3728 ♂, BMNH.

Hudson 1928, p. 184, pl. xix fig. 27, as *Scoparia triclera*.

***trivirgata*** Felder & Rogenhofer, 1875, pl. cxxxvii fig. 29 (as *Crambus trivirgatus*) new combination

"Ins. Neumunster" [Nelson NN, T.R. Oxley]; HT ♀ unique, BM Pyral. slide no. 3620 ♀, BMNH.

Hudson 1928, p. 198, pl. xxi fig. 42, as *Scoparia trivirgata*, after Meyrick (§1884d, p. 236; 1885e, p. 107).

***ustiramis*** Meyrick, 1931a, p. 95 (*Scoparia*) new combination

Whangarei ND, S.C. Patterson; HT ♂ unique, BM Pyral. slide no. 3632 ♂, BMNH.

Hudson 1939, p. 423, pl. lvi fig. 33, as *Scoparia ustiramis*.

***xysmatias*** Meyrick, 1907c, p. 110 (*Scoparia*) new combination

Old Man Range CO, J.H. Lewis; HT ♂ unique, BM Pyral. slide no. 3739 ♂, BMNH.

Hudson 1928, p. 187, pl. xxi fig. 5, as *Scoparia sysmatias*.

***zophoclaena*** Meyrick, 1923, pp. 162–163 (*Scoparia*) new combination

Takapuna AK, G.V. Hudson; HT ♂ unique, BM Pyral. slide no. 3745 ♂, BMNH.

Hudson 1928, p. 186, pl. xliv fig. 3, as *Scoparia zophoclaena*.

● ***Exsilirarcha*** Salmon & Bradley, 1956, pp. 73–75.  
Type species *Exsilirarcha graminea* Salmon & Bradley, by original designation.

***graminea*** Salmon & Bradley, 1956, pp. 75–77  
(*Exsilirarcha*)

Campbell Island, J.H. Sorensen; HT ♂ designated by Salmon & Bradley, NMNZ.

● ***Protyparcha*** Meyrick, 1909b, p. 71. Type species *Protyparcha scaphodes* Meyrick, by original monotypy.

***scaphodes*** Meyrick, 1909b, p. 71 (*Protyparcha*)  
Carnley Harbour, Auckland Island, G.V. Hudson; 3 ST ♂♂ labelled "Kermadec Island", BMNH.

Hudson 1928, p. 170, pl. xx fig. 40, in Crambinae, after Meyrick (1909).

Note. The ♂ selected by M. Shaffer as the LT cannot be, as it was presented to BMNH in 1916 (accession no. 232) by Hudson. Meyrick mixed his labelling on specimens he received through Hudson from the Auckland Islands (1907) and Kermadec Islands (1908).

● ***Scoparia*** Haworth, 1811, p. 498. Type species *Scoparia pyralea* Haworth, 1811 (as *Tinea pyrarella* Hübner, 1796, a junior primary homonym of *Tinea pyrarella* [Denis & Schiffermüller], 1775), by subsequent designation (Curtis 1827, folio 170).  
Note. Fletcher & Nye (1984, p. 139) give details of the homonymy.

***Xeroscopa*** Meyrick, 1884e, p. 349. Type species *Scoparia ejuncida* Knaggs, by subsequent designation (Hampson 1897, p. 226). Synonymised by Meyrick (1913, p. 41).

Note. Species included here have the ♂ valva with a process on the ventral margin (see Munroe 1964, p. 266; 1972, p. 29).

***acharis*** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 73 (key) and 85–86 (*Scoparia*)  
Akaroa MC, R.W. Fereday; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3735 ♂, BMNH.

Hudson 1928, p. 186, pl. xxi fig. 14.

***aphelis*** Meyrick, §1884d, p. 237; 1885e, pp. 110 (key) and 115 (*Xeroscopa*)

Arthur's Pass NC/WD, 4500 ft, E. Meyrick; HT ♂ unique, BM Pyral. slide no. 3661, BMNH.

Hudson 1928, p. 201, not figured. Meyrick 1913b, p. 45, as *Scoparia aphelis*.

***augastis*** Meyrick, 1907c, p. 112 (*Scoparia*)  
Invercargill SL, A. Philpott; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3641 ♂, BMNH.

Hudson 1928, p. 199, pl. xxiv fig. 8.

***autochroa*** Meyrick, 1907c, pp. 110–111 (*Scoparia*)  
Invercargill SL, A. Philpott; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3742, BMNH.

Hudson 1928, p. 188, pl. xxi fig. 16.

***chalicodes*** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 72 (key) and 98 (*Scoparia*)  
[sandhills], Christchurch MC, E. Meyrick; LT ♂ labelled as HT, BM Pyral. slide no. 3665 ♂, BMNH.

Hudson 1939, p. 424, pl. lxii fig. 14, corrected description, as *Scoparia chalicodes*; see "*Scoparia*" *limatula*, below.

***ciserodes*** Meyrick, 1920a, p. 30 (*Scoparia*). Synonymised by Hudson (1928, p. 194).

Wellington WN, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.

Hudson 1928, p. 194, as synonym.

***cyameuta*** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 110 (key) and 112–113 (*Xeroscopa*)  
Mount Hutt MC, R.W. Fereday; LT ♂ labelled as HT, BM Pyral. slide no. 3636 ♂, BMNH.

Hudson 1928, p. 199, pl. xxi fig. 54, as *Scoparia cyameuta*, after Meyrick (1913b, p. 44).

***dryphactis*** Meyrick, 1911b, p. 61 (*Scoparia*)  
[Wallacetown SL], A. Philpott; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3637 ♂, BMNH.

Hudson 1928, p. 200, pl. xxi fig. 53.

***ejuncida*** Knaggs, 1867, p. 81 (*Scoparia*)  
?Rakaia MC, H.G. Knaggs; HT ♂ unique, BM Pyral. slide no. 3604, BMNH.

Hudson 1928, p. 201, pl. xxi fig. 52 (a poor representation).

***encapna*** Meyrick, 1888c, p. 65 (*Scoparia*)  
Mount Arthur NN, 3800 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3741 ♂, BMNH.

Hudson 1928, p. 188, pl. xxii fig. 6 (a rather dark depiction).

***halopis*** Meyrick, 1909b, p. 72 (*Scoparia*)  
Carnley Harbour, Auckland Islands, G.V. Hudson; LT ♂ labelled as HT, BM Pyral. slide no. 3635 ♂, BMNH.

Hudson 1928, p. 199, pl. xxiv fig. 50 (not an accurate depiction).

***harpalaea*** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 110 (key) and 114 (*Xeroscopa*)  
Otira Gorge WD, 1600 ft, E. Meyrick; HT ♂ unique, BM Pyral. slide no. 3639 ♂, BMNH.

Hudson 1928, p. 201, not figured, as *Scoparia harpalaea*, after Meyrick (1913b, p. 44).

***nomeutis*** Meyrick, §1884d, p. 237 (no description); 1885e, pp. 110 (key) and 116–117 (*Xeroscopa*)  
[Ben Lomond] OL, 3,500 ft, E. Meyrick; LT ♂ selected by E. Munroe, BM Pyral. slide no. 3705 ♂, BMNH.

Hudson 1928, p. 202, pl. I fig. 28 (a poor depiction), as *Scoparia nomeutis*, after Meyrick (1913b, p. 45).

» Pyralidae, *Scoparia*

**parachalca** Meyrick, 1901, p. 569 (*Scoparia*)  
Lake Tekapo MK, G.V. Hudson; HT ♂ unique, BM Pyral.  
slide no. 3707 ♂, BMNH.  
Hudson 1928, p. 202, pl. xxii fig. 10.

**parmifera** Meyrick, 1909b, p. 72 (*Scoparia*)  
Carnley Harbour, Auckland Islands, G.V. Hudson; HT  
♂ unique, BM Pyral. slide no. 3646 ♂, BMNH.  
Hudson 1928, p. 186, pl. xxi fig. 13.  
Note. HT label reads "Kermadec Is GVH. 08".

**petrina** Meyrick, §1884d, p. 236 (no description);  
1885e, pp. 110 (key) and 111–112 (*Xeroscopa*)  
Mount Hutt MC, R.W. Fereday; LT ♂ selected by E.  
Munroe, BM Pyral. slide no. 3634 ♂, BMNH.  
Hudson 1928, p. 199, pl. xxi fig. 11 and 22, as *Scoparia*  
*petrina*, after Meyrick (1913, p. 44).

**legionaria** Philpott, 1928a, p. 362 (*Scoparia*).  
Synonymised by Hudson (1939, p. 426).  
Gordon's Pyramid, Mount Arthur NN, A. Philpott; HT  
♂ designated by Philpott, NZAC.  
Hudson 1939, p. 426, as synonym.

**rotuella** Felder & Rogenhofer, 1875, pl. cxxxvii fig.  
30 (as *Crambus rotuellus*)  
[Nelson NN, T.R. Oxley], "Ins. Neumunster"; LT ♂  
selected by E. Munroe, BM Pyral. slide no. 3618 ♂,  
BMNH.  
Hudson 1928, p. 200, pl. xxi fig. 41, as *Scoparia rotuella*,  
after Meyrick (1913b, p. 44).

**sideraspis** Meyrick, 1905, p. 231 (*Scoparia*)  
Mount Earnslaw OL, G.V. Hudson; LT ♂ selected by E.  
Munroe, BM Pyral. slide no. 3708 ♂, BMNH.  
Hudson 1928, p. 202, pl. xxiv fig. 24.

**triscelis** Meyrick, 1909b, p. 71 (*Scoparia*)  
Carnley Harbour, Auckland Islands, G.V. Hudson; HT  
♂ unique, BM Pyral. slide no. 3736 ♂, BMNH.  
Hudson 1928, p. 191, pl. xxi fig. 4.

**ustimacula** Felder & Rogenhofer, 1875, pl. cxxxv  
fig. 17 (*Scoparia*)  
[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen miss-  
ing, BMNH.

Hudson 1928, p. 189, pl. xxii fig. 39.

**conifera** Butler, 1879a, p. 493 (*Scoparia*). Syn-  
onymised by Meyrick (§1884d, p. 236; 1885e,  
p. 91).  
[Dunedin DN], F.W. Hutton; HT ♂ so labelled, BM Pyral.  
slide no. 3623 ♂, BMNH.  
Hudson 1928, p. 189, as synonym.

● **Scoparia** of authors

Note. Valval structures of males of the following  
species have not been examined.

**albafascicula** Salmon in Salmon & Bradley, 1956,  
p. 78 (*Scoparia*)  
"Campbell Island", J.H. Sorensen; HT ♂ unique, abdo-  
men missing, not found in NMNZ.  
Note. AMNZ has 3 specimens agreeing with the descrip-  
tion that were collected by E.G. Turrott on Antipodes  
Island. Collectors on Campbell Island have consist-  
ently failed to find *S. albafascicula*.

**animosa** Meyrick, 1914a, p. 103 (*Scoparia*)  
West Plains SL, A. Philpott; HT ♀ unique, BM Pyral.  
slide no. 3730 ♀, BMNH.  
Hudson 1928, pp. 186–187, pl. xlvi fig. 5.

**asaleuta** Meyrick, 1907c, pp. 111–112 (*Scoparia*)  
Lake Wakatipu OL, G.V. Hudson; LT ♀ selected by E.  
Munroe, BM Pyral. slide no. 3662 ♀, BMNH.  
Hudson 1928, p. 193, pl. xxi fig. 17.

**astragalota** Meyrick, §1884d, p. 236 (no descrip-  
tion); 1885e, pp. 110 (key) and 113 (*Xeroscopa*)  
Mount Hutt MC, R.W. Fereday; LT ♀ selected by E.  
Munroe, BM Pyral. slide no. 3638 ♀, BMNH.  
Hudson 1928, p. 200, pl. xxiv fig. 49, as *Scoparia astragalota*,  
after Meyrick (1913b, p. 44).

**autumna** Philpott, 1927d, p. 83 (*Scoparia*)  
Nelson NN, A. Philpott; HT ♂ designated by Philpott,  
NZAC.  
Hudson 1939, p. 425, pl. lvi fig. 35.

**caesia** Philpott, 1926a, p. 390 (as *Orocrambus*  
*caesius*)  
Gordon's Pyramid, Mount Arthur NN, 5000 ft, A. Phil-  
pott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, pp. 159–160, pl. li fig. 16, as *Orocrambus*  
*caesius*. Gaskin 1975, p. 266, excluded from Cram-  
binae, as ?*Scoparia caesia*.

**caliginosa** Philpott, 1918, p. 127 (*Scoparia*)  
?Matakanui, Manuherikia Valley CO, J.H. Lewis; HT ♂  
designated by Philpott, NZAC.  
Hudson 1928, p. 202, pl. xlvi fig. 2.

**cinefacta** Philpott, 1926a, p. 391 (*Scoparia*)  
Gordon's Pyramid, Mount Arthur NN, A. Philpott; HT  
♂ designated by Philpott, NZAC.  
Hudson 1928, p. 188, pl. xlvi fig. 7.

**claranota** Howes, 1946, p. 147 (*Scoparia*)  
Homer Cirque FD, W.G. Howes; HT ♂ designated by  
Howes, NMNZ.  
Howes 1946, p. 147, pl. 8 fig. 2, pl. 9 fig. 4.

**clavata** Philpott, 1912, p. 116 (*Scoparia*)  
Hump Ridge FD, A. Philpott; HT ♂ designated by Phil-  
pott, NZAC.  
Hudson 1928, p. 198, pl. xxi fig. 40.

**contexta** Philpott, 1931, p. 28 (*Scoparia*)  
Mount Moltke, Franz Josef WD, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.  
Hudson 1939, pp. 425–426, pl. lvi fig. 24.

**crepuscula** Salmon, 1946, pp. 7–8 (*Scoparia*)  
Homer Cirque FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.  
Salmon 1946, pp. 7 and 8, pl. 1 fig. 14.

**declivis** Philpott, 1918, p. 126 (*Scoparia*)  
Commissioner's Creek, Wakatipu OL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 200, pl. xliv fig. 23.

**diphtheralis** Walker, 1866a, p. 1501 (*Scoparia*)  
[Nelson NN], T.R. Oxley; LT ♀ labelled as HT, BM Pyral. slide no. 3749 ♀, BMNH. Hudson 1928, p. 192, pl. xxii fig. 20.

**ergatis** Meyrick, §1884d, p. 235 (no description); 1885e, pp. 74 (key) and 88 (*Scoparia*)  
Castle Hill MC, 3000 ft. E. Meyrick; LT ♀ selected by E. Munroe, BM Pyral. slide no. 3740 ♀, BMNH.  
Hudson 1928, p. 188, pl. xxii fig. 28.

**exilis** Knaggs, 1867, p. 81 (*Scoparia*)  
?Rakaia MC, H.G. Knaggs; HT ♀ unique, BMNH.  
Hudson 1928, p. 197, pl. xxi fig. 44 (a poor depiction).

**falsa** Philpott, 1924a, p. 208 (*Scoparia*)  
Gouland Downs NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 200, pl. 1 fig. 23.

**famularis** Philpott, 1930b, pp. 3–4 (*Scoparia*)  
Kepler Mountains FD, C.E. Clarke; HT ♀ designated by Philpott, AMNZ.  
Hudson 1939, p. 427, pl. lvi fig. 25.

**fimbriata** Philpott, 1917b, p. 243 (*Scoparia*)  
Mount Cleughearn FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 194, pl. xiv fig. 3.

**fumata** Philpott, 1915, p. 198 (*Scoparia*)  
Longwood Range SL, A. Philpott; HT ♂ designated by Philpott, abdomen missing, NZAC.  
Hudson 1928, p. 204, pl. xliv fig. 11.

**gracilis** Philpott, 1924a, p. 209 (*Scoparia*)  
Mount Arthur NN, 4500 ft, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 189, pl. 1 fig. 27.

**humilialis** Hudson, 1950, pp. 102–103 (*Scoparia*)  
Sinclair Head WN, G.V. Hudson; LT ♂ here designated, labelled "1288a", NMNZ.  
Hudson 1950, pp. 102–103, pl. v fig. 4.

**illota** Philpott, 1919, pp. 224–225 (*Scoparia*)  
Bluecliffs FD, A. Philpott; HT ♂ missing, AT ♀ designated by Philpott, NZAC.  
Hudson 1928, p. 196, pl. xx fig. 12.

**indistinctalis** Walker, 1863b, p. 48 (*Hypochalcia*)  
[Auckland AK], D. Bolton; HT ♀ (?LT; see Walker's description), BM Pyral. slide no. 3602 ♀, BMNH.  
Hudson 1928, pp. 193–194 (but pl. xxi fig. 10 refers to *Eudonia rakaiaensis*).

**limatula** Philpott, 1930c, p. 436 (*Scoparia*)  
Tisbury SL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 194, pl. xxii fig. 13, as *Scoparia chalicoetes*; 1939, p. 424, as *Scoparia limatula*.

**lychnophanes** Meyrick, 1927a, p. 697 (*Scoparia*)  
Mount Holdsworth WN, G.V. Hudson; HT ♀ unique, BM Pyral. slide no. 3647 ♀, BMNH.  
Hudson 1928, p. 188, pl. lii fig. 10.  
Note. Hudson notes that the specimen was found at 4000 ft (as does Meyrick), but the label on the HT states "2500 ft".

**minuscinalis** Walker, 1866a, p. 1503 (*Scoparia*)  
[Hawkes Bay HB or Taupo area TO], W. Colenso; HT ♂ unique, BMNH.  
Hudson 1928, pp. 184–185, pl. xxii fig. 38.

**molifera** Meyrick, 1926b, p. 415 (*Scoparia*)  
Ashhurst, Manawatu WI, G.V. Hudson; HT ♀ unique, BM Pyral. slide no. 3659 ♀, BMNH.  
Hudson 1928, p. 187, pl. li fig. 18.

**monochroma** Salmon, 1946, p. 7 (*Scoparia*)  
Arthur River, Milford Sound FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.  
Salmon 1946, p. 7, pl. 1 fig. 9.

**niphospora** Meyrick, §1884d, p. 237; 1885e, pp. 110 (key) and 115 (*Xeroscopa*)  
Castle Hill MC, 2500 ft, E. Meyrick; HT ♂ (?LT; see Meyrick's description – "five specimens"), BMNH.  
Hudson 1928, p. 201, pl. xxiv fig. 9, as *Scoparia niphospora*, after Meyrick (1913b, p. 45).

**pallidula** Philpott, 1928a, p. 363 (*Scoparia*)  
Mount Grey NC, W. Heighway; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 427, pl. lvi fig. 12.

**panopla** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 74 (key) and 107 (*Scoparia*)  
Mount Hutt MC, R.W. Fereday; LT ♂ selected by E. Munroe, BMNH.  
Hudson 1928, p. 198, pl. xxi fig. 43 (but the specimen figured lacks the yellowish suffusion around the basal streak present on the LT).

➤ Pyralidae, *Scoparia*

**parca** Philpott, 1928a, pp. 362–363 (*Scoparia*)  
Mount Grey NC, S. Lindsay & W. Heighway; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 427, pl. lvi fig. 15.

**pascoella** Philpott, 1920, pp. 43–44 (*Scoparia*)  
Tooth Peaks OL, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 203, pl. xlviii fig. 8.

**phalerias** Meyrick, 1905, pp. 230–231 (*Scoparia*)  
Wellington WN, G.V. Hudson; HT ♀ unique, BM Pyral. slide no. 3658 ♀, BMNH.  
Hudson 1928, p. 192, pl. xxi fig. 6.

**pura** Philpott, 1924a, p. 208 (*Scoparia*)  
Mount Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 202, pl. xxiv fig. 39 and 40.

**scripta** Philpott, 1918, pp. 126–127 (*Scoparia*)  
[Mount Burns], Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, pp. 200–201, pl. xliv fig. 24.

**sinuata** Philpott, 1930c, p. 436 (*Scoparia*)  
Mount Cook MK, A. Philpott; HT ♂ designated by Philpott, CMNZ.  
Hudson 1939, p. 426, pl. lvi fig. 26.

**subita** Philpott, 1912, p. 116 (as *Orocrambus subitus*)  
Hump Ridge FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 159, pl. xix fig. 3 and 4, as *Orocrambus subitus*. Philpott 1929, p. 497, footnote, “*Orocrambus subitus* Philpott ... proves to be a *Scoparia*”. Hudson 1939, p. 423, as *Scoparia subita*.

**sylvestris** Clarke, 1926, p. 418 (*Scoparia*)  
Otira River WD, C.E. Clarke; HT ♀ designated by Clarke, AMNZ.  
Not mentioned by Hudson.  
Note. The colour pattern resembles that of *Eudonia choristis*.

**tetracycla** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 73 (key) and 97 (*Scoparia*)  
Christchurch MC, R.W. Fereday; LT ♀ selected by E. Munroe, BM Pyral. slide no. 3663 ♀, BMNH.  
Hudson 1928, p. 193, pl. xxi fig. 21 (a rather dark depiction).

**trapezophora** Meyrick, §1884d, p. 236 (no description); 1885e, pp. 73 (key) and 93 (*Scoparia*)  
Castle Hill MC, 3000 ft, E. Meyrick; HT ♀ unique, BM Pyral. slide no. 3650 ♀, BMNH.  
Hudson 1928, p. 190, pl. xxi fig. 50.

**tuicana** Clarke, 1926, p. 418 (*Scoparia*)  
Waitati DN, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.  
Hudson 1939, p. 427, pl. lvi fig. 9.

**turneri** Philpott, 1928a, p. 362 (*Scoparia*)  
Arthur's Pass NC/WD, A.J. Turner; HT ♂ designated by Philpott, NZAC.  
Hudson 1939, p. 425, not figured.

**valenternota** Howes, 1946, pp. 146–147 (*Scoparia*)  
Homer Cirque FD, W.G. Howes; HT ♂ designated by Howes, NMNZ.  
Howes 1946, p. 146, pl. 8 fig. 1, pl. 9 fig. 1.

**declavata** Hudson, 1950, 101–102 (*Scoparia*). New synonymy.

Homer Tunnel, Homer Cirque FD, W.G. Howes; LT ♂ here designated, labelled “1291a”, NMNZ.  
Hudson 1950, pp. 101–102, pl. vi fig. 8 (cf. Howes 1946, pl. 8 fig. 1).


Note. The specimens seen by Hudson (1950, p. 101) are part of the series taken by Howes in December 1942 – January 1943. J.T. Salmon, a close friend of Hudson's, collected with Howes at this time (Howes 1946, p. 139). Specimens in the Howes and Hudson collections at NMNZ are indistinguishable, and the above synonymy must be proposed.

**vulpecula** Meyrick, 1927a, pp. 697–698 (*Scoparia*)  
Bold Peak OL, G.V. Hudson; HT ♀ unique, BM Pyral. slide no. 3666 ♀, BMNH.  
Hudson 1928, p. 195, pl. xxi fig. 32.

**Excluded species:** Klima (1939, p. 353) lists under *Ischnurges* Lederer the species *Nesolocha autolitha* Meyrick as from “?New-Seeland”. The original description by Meyrick (1886g, p. 240) clearly states “Port Moresby, New Guinea”.



**Superfamily GEOMETROIDEA**  
**Family GEOMETRIDAE**  
**Subfamily ENNOMINAE**  
(in the sense of McGuffin 1972, pp. 5–6)



**Geometridae: Ennominae**  
(177) *Declana junctilinea* (Walker)

● ***Chalastra*** Walker, 1862b, pp. 1429–1430. Type species *Chalastra pellurgata* Walker, by original monotypy.

Note. Under this name I place species included by Meyrick in *Selidosema* of authors, with a reduced, rounded uncus and no spinose process on the disc of the valva.

***aristarcha*** Meyrick, 1892, pp. 216–217 (*Selidosema*) new combination

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1898, p. 85, pl. ix fig. 17 and 18; 1928, pp. 137–138, pl. xvi fig. 30 and 31; as *Selidosema aristarcha*.

***ochrea*** Howes, 1911, p. 127 (*Selidosema*) new combination

Dunedin DN, W.G. Howes; HT ♂ designated by Howes, BMNH. Hudson 1928, p. 142, pl. xlvi fig. 27.

***prototoxa*** Meyrick, 1919, pp. 350–351 (*Selidosema*). New synonymy.

Tokau TO, C. O'Connor; HT ♀ unique, BMNH. Hudson 1928, pp. 142–143, pl. xlvi fig. 23 and 24, as species.

Note. Material in NZAC shows clear intergradation of the two colour forms; genitalia are indistinguishable, and therefore synonymy is proposed.

***pellurgata*** Walker, 1862b, p. 1430 (*Chalastra*)

[Nelson NN], T.R. Oxley; HT ♀ unique, BMNH. Hudson 1898, pp. 88–89, pl. ix fig. 33–36, as *Chalastra pelurgata*, after Meyrick (§1884c, p. 235; 1885d, p. 66), an unjustified emendation; 1928, p. 137, pl. xvi fig. 14–17, as *Selidosema pelurgata*.

***cinerascens*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 1 (*Itama?*). Synonymised by Meyrick (§1884c, p. 235; 1885d, p. 66). [Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH. Hudson 1898, p. 88; 1928, p. 137; as synonym.

***streptophora*** Meyrick, §1883d, p. 530; 1884b, p. 106 (*Stratocleis*). Synonymised by Meyrick (§1884c, p. 235; 1885d, p. 66).

Otira Gorge WD, 1500 ft, E. Meyrick; LT ♀ here designated, labelled "Otira Gorge New Zealand 1500 ft 24/1/83", "Lectotype ♀ Stratocleis streptophora Meyrick, J.S. Dugdale 1981", BMNH.

Hudson 1898, p. 88; 1928, p. 137; as synonym.

***pelurgata*** Meyrick, §1884c, p. 235; 1885d, p. 66 (*Chalastra*), unjustified emendation.

● ***Cleora*** Curtis, 1825, p. 88 (Fletcher 1967, pp. 4–5). Type species *Geometra cinctaria* Denis & Schiffermüller, 1775, by original designation; Palearctic.

***Barsine*** Meyrick, §1883d, p. 530; 1884b, p. 100; not Walker, 1854, p. 546. Type species *Scotosia panagrata* Walker, by original monotypy and subsequent designation (Meyrick 1917b, p. 266). Synonymised by Fletcher (1967, p. 5).

***Meyrickia*** Butler, 1884b, p. 133, replacement name for *Barsine* Meyrick, preoccupied by *Barsine* Walker, 1854 (Arctiidae). Synonymised by Fletcher (1967, p. 5).

Note. Fletcher (1967, pp. 4–5) gives synonymies applying beyond New Zealand.

***scriptaria*** Walker, 1860b, p. 422 (*Tephrosia*) [Wellington WN]. Parry; HT ♀ unique, BMNH.

Hudson 1898, p. 86; 1928, p. 145; as synonym of *Selidosema dejectoria*, after Meyrick (§1883d, p. 530; 1917b, p. 268). Hudson 1898, p. 87, pl. ix fig. 25–30; 1928, pp. 144–145, pl. xvii fig. 20–23; as *Selidosema panagrata*, after Meyrick (§1883d, p. 530; 1917b, p. 268). Fletcher 1967, p. 112, as *Cleora scriptaria*.

***stigmatica*** Walker, 1862b, pp. 1359–1360 (*Scotosia*). Synonymised by Fletcher (1967, p. 112). [Auckland AK], D. Bolton; HT ♀ unique, BMNH.

Hudson 1898, p. 86; 1928, p. 145; as synonym of *Selidosema dejectoria*.

***panagrata*** Walker, 1862b, p. 1360 (*Scotosia*). Synonymised by Fletcher (1967, p. 112).

[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH. Hudson 1898, p. 87; 1928, pp. 144–145; as *Selidosema panagrata*, after Meyrick (1917b, p. 268).

***menanaria*** Walker, 1863a, p. 1500 (*Angerona*). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 100).

[?Auckland AK], J.F. Churton; HT ♀ unique, BMNH. Hudson 1898, p. 87; 1928, p. 144; as synonym.

***antipodaria*** Felder, 1875, pl. cxxvi fig. 2 (*Epirranthis*). Synonymised by Meyrick (§1883d, p. 530;

➤ Geometridae, *Cleora scriptaria*

1884b, p. 100).

[Nelson NN, T.R. Oxley]; HT ♀ unique, head missing, BMNH.

Hudson 1898, p. 87; 1928, p. 144; as synonym.

**desiccata** Butler, 1879a, p. 459 (*Hyperythra*).

Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 100).

[Dunedin DN], F.W. Hutton; HT ♂ designated by Butler, BMNH.

Hudson 1898, p. 87; 1928, p. 144; as synonym.

**arenacea** Butler, 1879a, p. 459 (*Hyperythra*).

Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 100).

[Dunedin DN], F.W. Hutton; HT ♀ unique, BMNH.

Hudson 1898, p. 87; 1928, p. 144; as synonym.

• **Declana** Walker, 1858c, p. 1649. Type species *Declana floccosa* Walker, 1858c, by original monotypy.

**Ipana** Walker, 1858c, pp. 1661–1662. Type species *Ipana leptomera* Walker, 1858, by original monotypy. Synonymised by Meyrick (1917b, p. 270).

**Argua** Walker, 1863c, p. 448. Type species *Argua scabra* Walker, 1863, by original monotypy. Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 102).

**Detunda** Walker, 1865a, pp. 618–619. Type species *Detunda atronivea* Walker, 1865, by original monotypy. Synonymised by Hudson (1898, p. 95).

**Politeia** Walker, 1865a, p. 642. Type species *Politeia juncitilinea* Walker, 1865, by original monotypy. Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 65).

**Amphitape** Felder & Rogenhofer, 1875, pl. cix fig. 10 (“Erklärung”). Type species *Amphitape crassitibia* Felder & Rogenhofer, 1875, by original monotypy. Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 103).

**Epicasis** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Atossa* Meyrick, §1883d, p. 530; 1884b, pp. 103–104, preoccupied by *Atossa* Thomson, 1864 (Coleoptera). Type species *Declana niveata* Butler, 1879, p. 500, by original monotypy. Synonymised by Hudson (1898, p. 98).

**Anatossa** Warren, 1894, p. 466, [needless] replacement name for *Atossa* Meyrick.

**atronivea** Walker, 1865a, p. 619 (*Detunda*)

[Wellington WN], Captain Parry; HT ♂ unique, BMNH. Hudson 1898, p. 95, pl. x fig. 33 and 34; 1928, pp. 153–154, pl. xviii fig. 18; as *Declana atronivea*.

**manxifera** Fereday, 1880, pp. 268–270 (?*Chlenias*). Synonymised by Meyrick (§1883d, p. 530;

1884b, p. 101).

Wellington WN, W.T.L. Travers; HT ♀ labelled as “TYPE”, CMNZ.

Hudson 1898, p. 95; 1928, p. 153; as synonym.

**egregia** Felder & Rogenhofer, 1875, pl. cxxxii fig. 24 (*Chlenias*)

[Nelson NN, T.R. Oxley], “Australia meridionalis”; HT ♂ unique, BMNH.

Hudson 1898, p. 96, pl. x fig. 35; 1928, p. 154, pl. xviii fig. 17; as *Declana egregia*.

**feredayi** Butler, 1877, p. 398, pl. 43 fig. 5 (*Declana*)

[Castle Hill] MC, J.D. Enys; HT ♂ unique, BMNH.

Hudson 1898, p. 96, as synonym of *Declana floccosa*, after Meyrick (§1883d, p. 530; 1884b, p. 102); 1928, p. 152, pl. xvii fig. 20–22, as species.

**sinuosa** Philpott, 1915, p. 197 (*Declana*). Synonymised by Prout (1927, p. 79).

Ben Lomond OL, M.O. Pasco; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 152, as synonym of *D. feredayi* (cf. Meyrick 1917b, p. 270, as species).

**floccosa** Walker, 1858c, pp. 1649–1650 (*Declana*)

[Auckland AK], D. Bolton; LT ♀ designated by D.S. Fletcher, BM Geometridae genitalia slide no. 5361, BMNH.

Hudson 1898, pp. 96–97, pl. x fig. 39–47; 1928, pp. 151–152, pl. xviii fig. 23–34.

**scabra** Walker, 1863c, p. 448 (*Argua*). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 102).

[Nelson NN], T.R. Oxley; LT ♀ here designated, labelled “Auckland N. Zeal. 60–73” (circular), “Type” (circular, green margin), “Geometridae genitalia slide no. 5362”, antennae missing, BMNH.

Hudson 1898, p. 96; 1928, p. 151; as synonym.

**nigrosparsa** Butler, 1879a, p. 500 (*Declana*). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 102).

[Dunedin DN], F.W. Hutton; HT ♀ designated by Butler, abdomen missing, BMNH.

Hudson 1898, p. 96; 1928, p. 151; as synonym.

**callista** Salmon, 1946, p. 4 (*Declana*). New synonymy.

Gertrude Cirque, Upper Hollyford Valley FD, J.T. Salmon; HT ♂ unique, NMNZ.

Hudson 1950, pp. 96–97, pl. vii fig. 3.

**glacialis** Hudson, 1903, p. 245 (*Declana*)

Sealey Range MK, G.V. Hudson; LT ♂ here designated, labelled “317e”, “[Mueller terminal moraine and lower spurs of Sealey Range, Dec 15 + 18, 1899]”, NMNZ. Hudson 1928, p. 153, pl. xviii fig. 9 and 10.

**griseata** Hudson, 1898, p. 98 (*Declana*)

Lake Wakatipu OL, G.V. Hudson; LT ♂ [Hudson label 334b: “Head of L. Wakatipu Jan 8 1894”] in Hudson Collection, NMNZ.

Hudson 1898, p. 98, pl. x fig. 32; 1928, p. 151, pl. xviii fig. 15 and 16.

**hermione** Hudson, 1898, p. 98 (*Declana*)  
[Khandallah WN], G.V. Hudson; HT ♂ unique, labelled  
“140a”, NMNZ.  
Hudson 1898, p. 98, pl. x fig. 36; 1928, p. 152, pl. xvii  
fig. 19.

**junctilinea** Walker, 1865a, p. 643 (*Politeia*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1898, p. 98, pl. x fig. 37 and 38, as *Declana junc-*  
*tilinea* Felder; 1928, p. 152, pl. xvii fig. 11–14, ascribed  
to Walker.

**verrucosa** Felder & Rogenhofer, 1875, pl. cxxxii  
fig. 22 (*Chlenias*). Synonymised by Meyrick (§1884c,  
p. 234).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BM Geometri-  
dae genitalia slide no. 4211, BMNH.  
Hudson 1928, p. 152, as synonym.  
Note. Hudson (1928, pl. xviii fig. 11) gives a good match  
for the colour pattern of HT *junctilinea*, and in pl. xviii  
fig. 13 likewise for the colour pattern of HT *verrucosa*.

**leptomera** Walker, 1858c, p. 1662 (*Ipana*)  
[Auckland AK], D. Bolton; LT ♂ so labelled, BMNH.  
Hudson 1898, p. 94, pl. x fig. 29, 31, and 31a, as *Ipana*  
*leptomera*; 1928, pp. 150–151, pl. xviii fig. 35–37, as  
*Declana leptomera*.  
**crassitibia** Felder & Rogenhofer, 1875, pl. cix fig.  
10 (*Amphitape*). Synonymised by Meyrick (§1884c,  
pp. 234–235; 1885b, p. 66).  
Hudson 1928, p. 150, as synonym.

**niveata** Butler, 1879a, p. 500 (*Declana*)  
[Dunedin DN], F.W. Hutton; HT ♂ designated by Butler,  
BMNH.  
Hudson 1898, pp. 98–99, not figured; 1928, p. 151, pl.  
xviii fig. 19.

**toreuta** Meyrick, 1929, pp. 486–487 (*Declana*)  
Arthur's Pass NC/WD, W.H. Burrow; LT ♀ here desig-  
nated, labelled “Selidosema toreuta Meyrick type ♀  
det. D.S. Fletcher 1960”, “Arthurs Pass New Zealand  
W.H. bred '28”, abdomen missing, BMNH.  
Hudson 1939, p. 417, pl. lv fig. 24 (♀); 1950, pl. viii fig.  
9 (♂; as *Declana floccosa*).  
Note. The specimen figured by Hudson (1950, p. 97, pl.  
v fig. 5) as *D. toreuta* ♂ is a characteristic subalpine  
*D. hermione*.

Also 2 undescribed species (NZAC).

● **Gellonia** Meyrick, §1884c, p. 234. Type species  
*Boarmia dejectaria* Walker, by original monotypy.

**Gelonia** Meyrick, 1885d, p. 65, incorrect sub-  
sequent spelling (Fletcher 1979, p. 90).  
Note. Fletcher (1979, p. 90) discusses the validity  
of *Gellonia* vs *Gelonia*. The genus is revived here,  
as on facies and genitalia it differs markedly from  
other species placed by Meyrick in *Selidosema* of  
authors.

**dejectaria** Walker, 1860b, p. 394 (*Boarmia*)  
[Wellington WN], Captain Parry; HT ♂ unique, BMNH.  
Hudson 1898, p. 86, pl. ix fig. 21–24; 1928, pp. 145–146,  
pl. xvii fig. 24–26; as *Selidosema dejectaria*, after  
Meyrick (1917b, p. 268).

**attracta** Walker, 1860b, pp. 394–395 (*Boarmia*).  
Synonymised by Meyrick (§1883d, p. 530; 1884b,  
p. 100).  
“New Zealand, Mr Sowerby, Major Parry, and Dr Sin-  
clair”; LT(?) ♀ so labelled, bearing accession label 45.30  
(i.e., Waikouaiti DN, P. Earl), BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym.  
Note. As the LT was not collected by Sowerby, Parry, or  
Sinclair, its status is doubtful.

**exprompta** Walker, 1860b, p. 395 (*Boarmia*).  
Synonymised by Meyrick (§1883d, p. 530; 1884b,  
p. 100).  
[Auckland AK], A. Sinclair; HT ♂ unique, BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym.

**patularia** Walker, 1860b, p. 422 (*Tephrosia*).  
Synonymised by Meyrick (§1883d, p. 530; 1884b,  
p. 100).  
[Auckland AK], A. Sinclair; HT ♀ unique, BM Geome-  
tridae genitalia slide no. 4206, BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym.

**erebinata** Walker, 1862b, pp. 1358–1359 (*Scot-  
osia*). Synonymised by Meyrick (§1883d, p. 530;  
1884b, p. 100).

[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym.

**lignosata** Walker, 1862b, p. 1361 (*Scotosia*).  
Synonymised by Meyrick (§1883d, p. 530; 1884b,  
p. 100).

[Auckland AK], D. Bolton; HT ♀ unique, abdomen miss-  
ing, BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym.

**caprimulgata** Felder & Rogenhofer, 1875, pl.  
cxxvi fig. 12 (*Hemerophila*). Synonymised by  
Meyrick (§1883d, p. 530; 1884b, p. 100).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym.

**pannularia** Guenée, 1868, p. 42 (*Gnophos*)  
[Christchurch MC], R.W. Fereday; HT ♂ so labelled,  
BMNH.

Hudson 1898, p. 86, pl. ix fig. 19 and 20, as *Selidosema*  
*dejectaria* (part), following Meyrick (1917b, p. 268),  
and as synonym; 1928, p. 145, not figured, as synonym.  
Note. This name is revived for the “stumpy winged” form  
of *dejectaria* of authors, distinct on pattern and wing  
shape – as noted by Meyrick (1884b, p. 100) and illus-  
trated by Hudson (1898, pl. ix fig. 19 and 20) – and  
on genitalia.

**maoriata** Felder & Rogenhofer, 1875, pl. cxxvi  
fig. 4 (*Scotopteryx*). Synonymised by Meyrick  
(§1883d, p. 530, as *masriata*; 1884b, p. 100), as  
synonym of *Boarmia dejectaria*.

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1898, p. 86; 1928, p. 145; as synonym of *Seli-  
dosema dejectaria*, after Meyrick (1917b, p. 268).

» Geometridae, *Gellonia pannularia*

**sulpitiata** Felder & Rogenhofer, 1875, pl. cxxvi fig. 7 (*Hemerophila*). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 100), as synonym of *Boarmia dejectaria*.

[Nelson NN, T.R. Oxley]; HT ♀ unique, tip of abdomen missing, BMNH.

Hudson 1898, p. 86; 1928, p. 145; as synonym of *Selidosema dejectaria*, after Meyrick (1917b, p. 268).

● ***Ischalis*** Walker, 1863a, p. 1749. Type species *Ischalis thermochromata* Walker, by original monotypy.

**Polygona** Guenée, 1868, p. 41, preoccupied by *Polygona* Geyer, 1837 (Epiplemidae). Type species *Polygona fortinata* Guenée, by original monotypy. Synonymised by Meyrick (1884b, p. 106).

**Stratocleis** Meyrick, §1883d, p. 530; 1884b, p. 105. Type species *Selenia gallaria* Walker, by subsequent designation (Poole 1970, p. 135). Synonymised by Poole (1970, p. 135).

**Gonophylla** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Phyllodyce* Meyrick, §1883d, p. 530; 1884b, p. 104; preoccupied by *Phyllodoce* Ranzani, 1817 (Vermes). Type species *Gonodontis nelsonaria* Felder & Rogenhofer. Synonymised by Meyrick (1917b, p. 269).

**Azelina** in the sense of Meyrick (1917b, p. 269) but not Guenée (1857). Synonymised by Poole (1970, p. 135).

**fortinata** Guenée, 1868, p. 41 (*Polygona*) [?Akaroa, Banks Peninsula MC], R.W. Fereday; HT ♂ labelled "ex typicalibus specimenibus" by Guenée, BMNH.

Hudson 1898, p. 93, pl. x fig. 24 and 25; 1928, pp. 148–149, pl. xvii fig. 7 and 8; as *Azelina fortinata*, after Meyrick (§1883d, p. 531; 1884b, p. 106; 1917b, p. 269).

**ziczac** Felder & Rogenhofer, 1875, pl. cxxvi fig. 4 (*Caustaloma*?). Synonymised by Meyrick (§1883d, p. 531; 1884b, p. 106).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Hudson 1898, p. 93; 1928, p. 148; as synonym.

**gallaria** Walker, 1860a, p. 185 (*Selenia*) [Waikouaiti DN], P. Earl; HT ♀ unique, BMNH. Hudson 1898, p. 92, pl. x fig. 13–23; 1928, pp. 149–150, pl. xviii fig. 1–6; as *Azelina gallaria*, after Meyrick (1917b, p. 269).

**thermochromata** Walker, 1863a, p. 1750 (*Ischalis*). Synonymised by Prout (1927, p. 79).

[Wellington WN], Major Parry; HT ♂ unique, abdomen broken, BMNH.

Hudson 1928, p. 149, as synonym, after Meyrick (1917b, p. 269).

**palthidiata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 21 (*Euchlaena*?). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 105).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Hudson 1898, p. 92, as synonym, after Meyrick (1884b, p. 105).

**palthidiata** var. ***cinerea*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 22 (*Euchlaena*?). Synonymised by Prout (1927, p. 79).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Not mentioned by Hudson.

**venustula** Salmon, 1956, pp. 574–575 (as subspecies of *Azelina gallaria*). **New synonymy.** Little Barrier Island CL, J.T. Salmon; HT ♂ unique, designated by Salmon, "Type in the author's collection", not found in NMNZ.

Salmon 1956, pp. 574–575, pl. 22 fig. 1.

Note. The characteristic *venustula* colour pattern is present in most populations of *I. gallaria*.

**nelsonaria** Felder & Rogenhofer, 1875, pl. cxxviii fig. 3 (*Gonodontis*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Hudson 1898, pp. 90–91, pl. x fig. 3–6, as *Gonophylla nelsonaria*, after Meyrick (§1883d, p. 530; 1884b, p. 104; §1885h, p. 589; 1886, p. 184); 1928, p. 150, pl. xvii fig. 7 and 8, as *Azelina nelsonaria*, after Meyrick (1917b, p. 269).

**felix** Butler, 1877, p. 389 (*Gonodontis*). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 104). [Castle Hill Station MC], J.D. Enys; HT ♀ designated as "type" by Butler, ♂ abdomen glued on, BMNH. Hudson 1898, p. 90; 1928, p. 150; as synonym.

**variabilis** Warren, 1895, p. 153 (*Polygona*)

[Nelson NN, T.R. Oxley]; LT ♂ selected by L.B. Prout and so labelled, BMNH.

Hudson 1898, p. 93, pl. x fig. 26–28, as *Azelina ophiopa*; 1928, p. 148, pl. xvii fig. 15–18, as *Azelina variabilis*.

**ophiopa** Meyrick, 1897b, p. 387 (*Gonophylla*). Synonymised by Prout (1927, p. 79).

Wellington WN, G.V. Hudson; HT ♂ unique, abdomen missing, BMNH.

Hudson 1898, p. 93, as species; 1928, p. 148, as synonym.

Also 1 undescribed species (NZAC).

● ***Pseudocoremia*** Butler, 1877, p. 394; reinstated by Prout (1912, p. 53). Type species *Selidosema?* *fragosata* Felder & Rogenhofer, by original designation.

**Zylobara** Butler, 1897, p. 498. Type species *Selidosema fenerata* Felder & Rogenhofer, 1975, by original designation. Synonymised by Meyrick (1888b, p. 61).

***Selidosema*** in the sense of Meyrick (1917b, p. 266), not Hübner ([1823], p. 299) (Prout 1927, p. 79).

***albafasciata*** Philpott, 1915, pp. 196–197 (*Selidosema*) new combination

Taihape RI, H. Hamilton; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 142, pl. xvii fig. 3, as *Selidosema albifasciata*.

Note. Although both the original spelling and the emendation “*albifasciata*” by Meyrick (1917, p. 267) are wrong formations (*albofasciata* is correct), *albafasciata* is retained as the correct original spelling under ICZN Rules, Article 32a (ii) and 32c. See also *Asaphodes albalineata*. HT ♂ externally is very similar to *lactiflava*; both have a characteristically shirred (transversely finely strigose) wing pattern.

***berylia*** Howes, 1943, p. 372 (*Selidosema*)

Homer Cirque FD, J. Sutherland; HT ♂ designated by Howes, NMNZ.

Hudson 1950, pp. 90–91, pl. v fig. 1.

***campbelli*** Philpott, 1927a, p. 705 (*Selidosema*) new combination

Blackball BR, J.W. Campbell; HT ♀ unique, designated by Philpott, NZAC.

Hudson 1928, p. 138, pl. xxxviii fig. 34, as *Selidosema campbelli*.

Note. *P. campbelli* is unusual in having pectinate antennae in the ♀. It has not been reported subsequently.

***cineracia*** Howes, 1942, pp. 277–278 (*Selidosema*) new combination

[Ben Mohr], Moke Lake OL, W.G. Howes; HT ♂ designated by Howes, NMNZ.

Hudson 1950, p. 92, pl. iii fig. 9, as *Selidosema cineracia*.

***colrogramma*** Meyrick, 1936, p. 281 (*Selidosema*) new combination

Arthur's Pass NC/WD, G.V. Hudson; LT ♀ here designated, labelled “Arthurs Pass New Zealand GVH bred '33”, “*Selidosema colrogramma* Meyr. allotype ♀” (L.B. Prout's writing), “LECTOTYPE ♀ *Selidosema colrogramma* Meyrick J.S. Dugdale 1981”, head and abdomen missing, BMNH.

Hudson 1939, p. 413, pl. lv fig. 31 and 32, as *Selidosema colrogramma*.

Note. Prout's original LT designation is a ♂ which shows none of the colour pattern characteristics depicted by Hudson, who based his illustrations of ♂ and ♀ on the original reared series. Designation of the LT ♀ as above fixes the name *colrogramma* on a specimen agreeing with both Meyrick's description and Hudson's illustration, bred from *Cassinia* (Asteraceae).

***fascialata*** Philpott, 1903, pp. 248–249, pl. xxxii fig. 7 (*Selidosema*) new combination

West Plains SL, A. Philpott; LT ♂ here designated, labelled “West Plains” (no date), one of 2 ♂♂ in Philpott Collection, NZAC.

Hudson 1928, p. 139, pl. xvi fig. 1 and 2, as *Selidosema fascialata*.

***fenerata*** Felder & Rogenhofer, 1875, pl. cxxxii fig.

7 (*Selidosema*) new combination

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.

Hudson 1898, p. 82, pl. viii fig. 50 and 51; 1928, p. 144, pl. xvii fig. 13 and 14; as *Selidosema fenerata*.

Note. Butler (1879, p. 498) referred to this species as *Zylobara fenerata*.

***argentaria*** Philpott, 1913, p. 77 (*Selidosema*). New synonymy.

West Plains SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 144, pl. xvii fig. 11 and 12, as species.

***adusta*** Philpott, 1930c, pp. 435–436 (*Selidosema*). New synonymy.

Governor's Bush MK, A. Philpott; HT ♂ designated by Philpott, CMNZ.

Hudson 1939, p. 415, pl. lv fig. 8, as species.

Note. No consistent differences in genitalia or larvae were observed between FRNZ specimens from Southland (= *argentaria*), subalpine localities in the North and South islands (= *adusta*), and lowland localities from Dunedin northwards (= *fenerata*), hence the proposed synonymy.

***flava*** Warren, 1896b, p. 406 (*Pseudocoremia*)

Greymouth WD, R. Helms; HT ♂ unique, BMNH.

Hudson 1939, p. 413, pl. lv fig. 23, as *Selidosema flava*, after Philpott (1928a, pp. 360–361).

***fluminea*** Philpott, 1926a, p. 389 (*Selidosema*) new combination

Flora River, Mount Arthur NN, A. Philpott; HT ♂ designated by Philpott, tip of abdomen missing, NZAC.

Hudson 1928, p. 139, as synonym of *Selidosema productata*; 1939, p. 414, and 1928, pl. xlvi fig. 26, as species, after Philpott (1928g, p. 485, fig. 9 and 10).

***indistincta*** Butler, 1877, p. 394, pl. 43 fig. 8 (*Pseudocoremia*)

[Dunedin DN], J. Hector; HT ♂ designated by Butler, BMNH.

Hudson 1928, p. 140, pl. xvii fig. 1 and 2, as *Selidosema indistincta*.

Note. A ♀ specimen from Blenheim MB, collected by W. Skellon, bears the manuscript name “var. *confusa* Butler”, and refers to “*Pseudocoremia productata* n.sp.” (Butler 1880, p. 551, line 5).

***insignita*** Philpott, 1930b, p. 2 (*Selidosema*) new combination

Kao ND, C.E. Clarke; HT ♂ unique, designated by Philpott, AMNZ.

Hudson 1939, pp. 414–415, pl. lv fig. 15, as *Selidosema insignita*.

Note. Males in NZAC have the apical 9–13 antennal segments non-pectinate, and the wing pattern differences noted by Philpott and figured by Hudson are variable.

***pergrata*** Philpott, 1930b, p. 2 (*Selidosema*). New synonymy.

» Geometridae, *Pseudocoremia insignita*

Sandymount DN, C.E. Clarke; HT ♂ unique, designated by Philpott, AMNZ.

Hudson 1939, p. 415, pl. lv fig. 16, as species.

**lactiflua** Meyrick, 1912c, pp. 117–118 (*Selidosema*) [Routeburn Valley], Lake Wakatipu OL, G.V. Hudson; LT ♂ here designated, labelled “L. Wakatipu New Zealand G.V.H. 2.11”, “Selidosema lactiflua Meyr. ♂ Type” (L.B. Prout’s writing), BMNH.

Hudson 1928, p. 140, pl. xvi fig. 27 and 28, as *Selidosema lactiflua*.

**leucelaea** Meyrick, 1909a, pp. 6–7 (*Selidosema*) Invercargill SL, A. Philpott; LT ♂ selected by L.B. Prout, BMNH.

Hudson 1898, pl. ix fig. 10 and 14, as *Selidosema productata*; 1928, p. 141, pl. xvi fig. 12 and 13, as *Selidosema leucelaea*.

**lupinata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 19 (*Cidaria*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Meyrick §1883d, p. 530; 1884b, p. 99; as *Pseudocoremia lupinata*. Hudson 1928, p. 143, pl. xvii fig. 5 and 6, as *Selidosema lupinata*, after Meyrick (1909, p. 7).

**humillima** Hudson, 1898, pp. 83–84 (*Selidosema*). Synonymised by Meyrick (1909a, p. 7). Wellington WN, G.V. Hudson; LT ♂ here designated, labelled “213a” [“at rest on fence W’ton Terrace Dec. 22 ‘88” – entry in Hudson Register], NMNZ.

Hudson 1898, pp. 83–84, pl. ix fig. 5.

**lutea** Philpott, 1914, p. 119 (*Selidosema*) new combination

Bold Peak, Humboldt Range FD, C.C. Fenwick; HT ♂ not designated by Philpott, one of 2 ♂♂ labelled “Bold Peak 24 Dec. 1913 C.C. Fenwick”, NMNZ.

Hudson 1928, p. 142, pl. xvi fig. 32 and 33, as *Selidosema lutea*.

**melinata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 9 (*Numeria*)

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH. Hudson 1898, p. 85, pl. ix fig. 15 and 16, as *Selidosema melinata*.

Note. Meyrick (§1883d, p. 530; 1884b, p. 99) referred to this species as *Pseudocoremia melinata*.

**pungata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 23 (*Selidosema*). Synonymised by Prout (1927, p. 79).

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen missing, BMNH.

Hudson 1898, p. 84, as synonym of *Selidosema productata*; 1928, p. 138, as synonym.

**cremnopa** Meyrick, 1897b, p. 387 (*Selidosema*). Synonymised by Prout (1927, p. 79).

Auckland AK, E. Meyrick; LT ♂ selected by L.B. Prout, abdomen missing, BMNH.

Hudson 1928, p. 138, as synonym.

**scariphota** Meyrick, 1915a, p. 202 (*Selidosema*).

Synonymised by Philpott (1931, p. 26).

Makara WN, R.M. Sunley; HT ♂ unique, BMNH.

Hudson 1928, p. 138, pl. xvi fig. 29, as species.

Note. Hudson (1939, p. 413) was unsure of distinctions between *melinata*, *cremnopa*, and *scariphota*. Material in NZAC from several localities and reared from *Carmichaelia* shows colour patterns of all three, thus upholding Prout’s synonymies.

**modica** Philpott, 1921, pp. 339–340 (*Selidosema*) new combination

Port Hills, Banks Peninsula MC, E.S. Gourlay; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 140, not figured; 1939, p. 414, pl. lvi fig. 1 and 2.

**monacha** Hudson, 1903, p. 245, pl. 30 fig. 4 (*Selidosema*)

?locality, ?collector; HT ♀ unique, head missing, CMNZ. Hudson 1928, p. 141, pl. xvi fig. 25 and 26, as *Selidosema monacha*.

**maculosa** Howes, 1914, p. 96 (*Pseudocoremia*). Synonymised by Hudson (1928, p. 141).

Queenstown OL, M.O. Pasco; HT ♂ unique, AMNZ. Hudson 1928, p. 141, as synonym.

**ampla var. *bistonaria*** Hudson, 1950, p. 93 (*Selidosema*). New synonymy.

Governor’s Bush, Mount Cook MK, G.V. Hudson; HT ♂ unique, NMNZ.

Hudson 1950, p. 93, pl. vii fig. 5, as variety.

Note. Hudson’s HT is a somewhat clouded but otherwise unexceptional specimen of *monacha*, the only known *Pseudocoremia* restricted to *Phyllocladus*.

**ombrodes** Meyrick, 1902c, pp. 275–276 (*Selidosema*) new combination

Chatham Island, J. Fougere; LT ♂ selected as HT by L.B. Prout, BMNH.

Hudson 1928, p. 144, pl. xliv fig. 6, as *Selidosema ombrodes*.

Note. See Holloway (1977, pp. 136–138, fig. 104, 107, and 108, pl. 29 fig. 11 and 12) for comparison with *P. christiana* Holloway from Norfolk Island.

**productata** Walker, 1862a, p. 1197 (*Larentia*)

[Nelson NN], T.R. Oxley; HT ♀ unique, abdomen glued on, BMNH.

Hudson 1898, p. 84, pl. ix fig. 6–14; 1928, p. 139, pl. xvi fig. 9–11; as *Selidosema productata*.

**fragosata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 28 (*Selidosema*?). Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 98).

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen in capsule, BMNH.

Hudson 1898, p. 84; 1928, p. 139; as synonym.

Note. See Holloway (1977, pp. 136–138, fig. 104, 107, and 108, pl. 29 fig. 11 and 12) for a comparison between *P. productata* and Norfolk Island *P. christiana* Holloway.

***rudisata rudisata*** Walker, 1862b, p. 1420 (*Cidaria*)  
new combination

[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1898, p. 82, pl. ix fig. 1 and 2; 1928, p. 143, pl.  
xvi fig. 23; as *Selidosema rudiata*, after Meyrick (1891,  
p. 101), an unjustified emendation.

***astrapia*** Meyrick, 1890, pp. 218–219 (*Boarmia*).  
Synonymised by Meyrick (1891, p. 101).

Wellington WN, G.V. Hudson; LT ♂ selected by L.B.  
Prout, BMNH.  
Hudson 1898, p. 82; 1928, p. 143; as synonym.

***rudisata ampla*** Hudson, 1923b, p. 129 (*Selido-*  
*sema*; as species)

Stoney Creek, Glenorchy OL, F.S. Oliver; HT (?gender)  
in Oliver Collection, lost.  
Hudson 1928, p. 143, pl. li fig. 8, pl. xvi fig. 23, as species.  
Note. North and South island populations of *rudisata* dif-  
fer in forewing pattern (see Hudson 1928, pl. xvi fig.  
23 and 24), North Island specimens having the lines  
strongly zig-zag.

***suavis*** Butler, 1879a, p. 497 (*Pseudocoremia*)  
[Christchurch MC], F.W. Hutton; HT ♂ designated by  
Butler as “♂ type”, BMNH.

Hudson 1898, p. 83, pl. ix fig. 3 and 4; 1928, p. 142, pl.  
xvi fig. 18–22; as *Selidosema suavis*.

***usitata*** Butler, 1879a, p. 501 (*Pachycnemia*).  
Synonymised by Meyrick (1891, p. 107).

[Dunedin DN], F.W. Hutton; HT ♀ designated by Butler  
as “type”, BMNH.  
Hudson 1898, p. 83; 1928, p. 142; as synonym.  
Note. HT *usitata* appears rather narrow-winged, but this  
may be due to rolling of the dorsum margin.

***terrena*** Philpott, 1915, p. 196 (*Selidosema*) new  
combination

Bold Peak, Humboldt Mountains OL, H. Hamilton; HT  
♂ designated by Philpott, NMNZ.  
Hudson 1928, p. 140, pl. xvi fig. 34, as *Selidosema terrena*.

Also 2 undescribed species (NZAC).

● ***Sarisa*** Fletcher, 1979, p. 185, replacement name  
for *Gargaphia* Walker. Type species *Gargaphia*  
*muriferata* Walker, by original monotypy.

***Gargaphia*** Walker, 1863a, p. 1364, preoccupied  
by *Gargaphia* Stål, 1862 (Hemiptera).

***muriferata*** Walker, 1863a, p. 1365 (*Gargaphia*)  
[Nelson NN], T.R. Oxley; HT ♀ unique, abdomen miss-  
ing, BMNH.

Hudson 1898, pp. 91–92, pl. x fig. 7–11, as *Drepanodes*  
*muriferata*, after Meyrick (§1883d, p. 531; 1884b, p.  
107); 1928, p. 147, pl. xvii fig. 34–36, as *Gargaphia*  
*muriferata*.

***ephyraria*** Walker, 1863a, p. 1761 (*Panagra*).  
Synonymised by Meyrick (§1883d, p. 531; 1884b,  
p. 107).

[Nelson NN], T.R. Oxley; HT ♀ unique, an abdomen  
glued on, BMNH.

Hudson 1898, p. 91; 1928, p. 147; as synonym.

***cookaria*** Felder & Rogenhofer, 1875, pl. cxxiii  
fig. 26 (*Zanclopteryx?*). Synonymised by Meyrick  
(§1883d, p. 531; 1884b, p. 107).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1898, p. 91; 1928, p. 147; as synonym.

***haastaria*** Felder & Rogenhofer, 1875, pl. cxxiii  
fig. 32 (*Zanclopteryx?*). Synonymised by Meyrick  
(§1883d, p. 531; 1884b, p. 107).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1898, p. 91; 1928, p. 147; as synonym.

***neoselena*** Meyrick, 1909b, p. 70 (*Drepanodes*).  
Synonymised by Dugdale (1971b, p. 91).

Carnley Harbour, Auckland Island, G.V. Hudson; HT ♀  
unique, previously labelled as PT, BMNH.  
Hudson 1928, p. 148, pl. xvii fig. 27, as *Gargaphia*  
*neoselena*.

Note. Dugdale (1971b, pp. 91 and 93) wrongly ascribed  
type status to Hudson's material, as Meyrick did not return  
this specimen or others from the subantarctic islands.

● ***Sestra*** Walker, 1863a, p. 1750. Type species *Ses-*  
*tra fusiplagiata* Walker, by original monotypy.

***Pseudosestra*** Butler, 1882b, p. 389. Type species  
*Lozogramma obtusaria* Walker (as *obtusata*), by  
original designation. Synonymised by Hudson  
(1898, p. 89) by implication through synonymy of  
*obtusaria* with *humeralia*.

***Amastris*** Meyrick, §1883d, p. 530; 1884b, p. 104;  
preoccupied by *Amastris* Stål, 1860 (Hemiptera)  
(Fletcher 1979, p. 190). Type species *Amastris*  
*encausta* Meyrick. Synonymised by Meyrick  
(§1885h, p. 589; 1886c, p. 184).

***flexata*** Walker, 1862b, p. 1421 (*Cidaria*)

[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1898, pp. 89–90, pl. x fig. 1 and 2, as *Sestra*  
*humeralia*; 1928, p. 146, pl. xvii fig. 30–33, as *Sestra*  
*flexata* (after Prout 1912, p. 54).

***fusiplagiata*** Walker, 1863a, p. 1751 (*Sestra*).  
Synonymised by Prout (1912, p. 54).

[Auckland AK], D. Bolton; HT unique, abdomen miss-  
ing, BMNH.  
Hudson 1898, p. 89; 1928, p. 146; as synonym.

***encausta*** Meyrick, §1883d, p. 530; 1884, p. 105  
(*Amastris*). Synonymised by Prout (1912, p. 54).

Otira Gorge WD, 1,500 ft, E. Meyrick; LT ♂ here des-  
ignated, labelled “Otira Gorge New Zealand 1500 ft  
24/1/83”, “Lectotype ♂ *Amastris encausta* Meyrick, J.S.  
Dugdale 1981”, BMNH.  
Hudson 1898, p. 89; 1928, p. 146; as synonym.

***humeralia*** Walker, 1861, p. 940 (*Macaria?*)

[Auckland AK], A. Sinclair; HT lost; specimen labelled  
as “Type” from Auckland, collected by D. Bolton (i.e.,  
label “New Zeal. 54-4”), BMNH.

» Geometridae, *Sestra humeraria*

Hudson 1898, p. 90, pl. ix fig. 37, as *Sestra flexata*; 1928, pp. 146–147, pl. xvii fig. 28 and 29, as *Sestra humeraria*, after Prout (1912, p. 54).

**obtusaria** Walker, 1861, p. 985 (*Lozogramma*).  
Synonymised by Meyrick (§1884c, p. 235; 1885d, p. 66); see also Prout (1912, p. 54).  
[Auckland AK], A. Sinclair; HT ♂ unique, BMNH.  
Hudson 1928, p. 146, as synonym.

**obtruncata** Walker, 1862b, p. 1421 (*Cidaria*?).  
Synonymised by Prout (1912, p. 54).

[Auckland AK], D. Bolton; LT ♂ here designated, labelled “New Zeal. 54.4” (circular), “92 Cidaria? obtruncata” (printed strip), “Type” (red-margined, circular), BMNH.  
Hudson 1928, p. 146, as synonym.

**punctilineana** Walker, 1866b, p. 1780 (*Teras*).  
Synonymised by Prout (1912, p. 54).

[Auckland AK], D. Bolton; HT unique, abdomen missing, BMNH.

Not mentioned by Hudson.

● **Zermizinga** Walker, 1863a, p. 1530. Type species *Zermizinga indocilisaria* Walker, by original monotypy.

**indocilisaria** Walker, 1863a, p. 1530 (*Zermizinga*) [Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂ unique, BMNH.

Hudson 1898, p. 88, pl. ix fig. 31 and 32; 1928, p. 147, pl. xvii fig. 9 and 10; as *Hybernia indocilis*, an unjustified emendation by Meyrick (§1883d, p. 530; 1884b, p. 97).

**boreophilaria** Guenée, 1868, p. 61 (*Hybernia*).  
Synonymised by Meyrick (§1883d, p. 530; 1884b, p. 97).


[Christchurch MC], R.W. Fereday; HT ♂ so labelled, BMNH.

Hudson 1898, p. 88; 1928, p. 147; as synonym.

Note. New Zealand and mainland Australian specimens differ in ♀ hindwing shape, but no Tasmanian material was available. In CMNZ there is a ♀ identical in wing shape to New Zealand ♀♀ “bred from *Acacia* from Tasmania”.

—◎—

**Subfamily LARENTIINAE**  
(in the sense of McGuffin 1967, p. 7)



**Geometridae: Larentiinae**  
(178) *Tatosoma tipulata* (Walker)

● **Anachloris** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Arsinoe* Meyrick.

**Arsinoe** Meyrick, §1883d, p. 527; 1884b, pp. 58 (key) and 73; preoccupied by *Arsinoe* Rafinesque, 1815 (Hymenoptera). Type species *Aspilates*(?) *subochraria* Doubleday, by subsequent designation (Meyrick 1917b, p. 255).

**subochraria** Doubleday, 1843, p. 285 (*Aspilates*?)  
[Auckland AK], A. Sinclair; HT ♀ unique, BMNH.  
Hudson 1898, p. 48, pl. vi fig. 45 and 46; 1928, p. 102, pl. xii fig. 37; as *Hydriomena subochraria*.

**strangulata** Guenée, 1857b, p. 423 (?*Camptogramma*; species credited to Zeller). Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 73).  
Adelaide S.A., P.C. Zeller; HT ♂ unique, BMNH.  
Hudson 1898, p. 48; 1928, p. 102; as synonym.

**euboliaria** Walker, 1863a, p. 1684 (*Aspilates*).  
Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 64).

[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1928, p. 102, as synonym.

**fuscinata** Guenée, 1868, p. 92 (*Camptogramma*). Synonymised by Meyrick (§1883d, p. 528, as *pusinata*; 1884b, p. 73).

[Christchurch MC], R.W. Fereday; HT not located.  
Hudson 1898, p. 48; 1928, p. 102; as synonym.

● **Aponotoreas** Craw, 1986, p. 136. Type species *Larentia anthracias* Meyrick, 1883, by original designation.

Note. Craw (1986, p. 138) includes 4 Australian species.

**anthracias** Meyrick, §1883d, p. 529; 1884b, pp. 77 (key) and 84 (*Larentia*)

Lake Wakatipu OL, 5,400 ft, E. Meyrick; LT ♂ labelled as "Type ♂", BMNH.

Hudson 1898, pp. 67–68, as *Xanthorhoe anthracias*; 1928, p. 124, pl. xiv fig. 35, as *Notoreas anthracias*.

**dissimilis** Philpott, 1914, pp. 118–119 (*Venusia*)

Ben Lomond OL, M.O. Pasco; ST series (♂, ♀) lost, formerly in Pasco Collection, SMNZ.

Hudson 1928, p. 119, pl. xiii fig. 8, as *Xanthorhoe dissimilis*, following Prout (1927, p. 76).

Note. Topotypic specimens collected in 1914 are in NZAC.

**incompta** Philpott, 1918, p. 126 (*Notoreas*)

Kepler Mountains FD, R. Gibb; HT ♂ designated by Philpott, "no. 1483", not located in SMNZ.

Hudson 1928, p. 124, pl. xv fig. 48.

Note. A ♂ designated by Philpott as "Paratype" is in NZAC.

**insignis** Butler, 1877, p. 393, pl. 43 fig. 1 (*Aspilates*)

[?Castle Hill MC or "Otago" CO], J.D. Enys or J. Hector; HT ♂ labelled by Butler as "type", BMNH.

Hudson 1898, p. 71, pl. viii fig. 3; 1928, p. 123, pl. xiv fig. 27 and 28; as *Notoreas insignis*, after Meyrick (§1885h, p. 589; 1886c, p. 184).

**orphnaea** Meyrick, §1883d, p. 529; 1884b, pp. 85–86 (*Pasithea*)

Ben Lomond OL, 5,600 ft, E. Meyrick; LT ♀ labelled as "Type" by L.B. Prout, BMNH.

Hudson 1898, pp. 71–72, not figured; 1928, p. 123, pl. xiv fig. 49 and 50; as *Notoreas orphnaea*, after Meyrick (§1885h, p. 589; 1886c, p. 184).

**synclinalis** Hudson, 1903, p. 244 (*Notoreas*)

Seaward Moss SL, A. Philpott; HT ♂ (specimen no. 544b) not in NMNZ.

Hudson 1928, pp. 122–123, pl. xiv fig. 33 and 34, as *Notoreas synclinalis*.

Note. NZAC has 1 ♂ labelled "Seaward Moss" but with no date, and "Notoreas synclinalis Huds.", but there is no indication that the specimen Philpott sent to Hudson was returned.

**villosa** Philpott, 1917b, pp. 241–242 (*Notoreas*)

The Hump FD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 123, pl. xv fig. 23 and 24, as *Notoreas villosa*.

• **Arctesthes** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Stratonice* Meyrick. As *Arcteuthes* in Meyrick 1888b, p. 47 (misprint).

**Stratonice** Meyrick, §1883d, p. 527; 1884b, pp.

58 (key) and 64; preoccupied by *Stratonice* Malmgren, 1867 (Vermes). Type species *Fidonia?* *catapyrrha* Butler, by original monotypy.

Note. Craw (1986, p. 136) reinstated this genus.

**catapyrrha** Butler, 1877, p. 392, pl. 43 fig. 2 (*Fidonia?*)

[?Castle Hill MC], J.D. Enys; HT ♂ designated by Butler as "Type", BMNH.

Hudson 1898, pp. 68–69, pl. viii fig. 35, as *Lythria euclidiata*, after Meyrick (§1884d, p. 234; 1885d, pp. 63–64 (error)); 1928, p. 131, pl. xv fig. 6, as *Lythria catapyrrha*, after Meyrick (1917b, p. 264).

Note. HT has the hindwings strongly red-flushed below, as have all Canterbury specimens. Otago specimens are less red-flushed, and the discal stripe is not red.

**catapyrrha ab. fasciata** Prout, 1939, p. 246 (*Arctesthes*). New synonymy.

Ben Lomond OL, A. Philpott; HT ♀ designated by Prout, BMNH.

**catapyrrha ab. kaikourensis** Prout, 1939, p. 246 (*Arctesthes*).

Mount Tapuaenuku KA, G.V. Hudson; LT ♂ here designated, labelled "Mt Tapuae-nuku (slopes of) 1.3.16", "219z", NMNZ.

Hudson 1928, p. 131, pl. xv fig. 4 and 5, as *Lythria siris*.

**siris** Hudson, 1908, p. 106 (*Lythria*)

Old Man Range CO, J.H. Lewis; LT ♂ here designated, labelled "725a" ["Old Man Range at about 4000 ft. Feb. '06 (J.H. Lewis)" – Hudson's Register], NMNZ.

Hudson 1928, p. 131, pl. xv fig. 4 and 5, as *Lythria siris*.

Prout 1939, p. 246, pl. 24 line h, as *Arctesthes siris*.

• **Asaphodes** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Thyone* Meyrick

**Thyone** Meyrick, §1883d, p. 527; 1884b, pp. 58 (key) and 61; preoccupied by *Thyone* Phillipi, 1840 (Crustacea). Type species *Aspilates abrogata* Walker, by original monotypy.

Note. *Asaphodes* was reinterpreted by Dugdale (1971b, pp. 93–95,) and most New Zealand "Xanthorhoe" species were apportioned either to *Helastria* (p. 180) or to *Asaphodes*. Species that were not apportioned, but which on subsequent examination are unequivocal in character, are listed under the relevant genus; others are listed under "Xanthorhoe of authors", p. 190.

**abrogata** Walker, 1862a, p. 1075 (*Aspilates*)

Waikouaiti DN, P. Earl; HT ♂ unique, abdomen missing, BMNH.

Hudson 1898, p. 55, pl. vii fig. 21; 1928, pp. 107–108, pl. xii fig. 19 and 20.

**servularia** Guenée, 1868, p. 43 (*Fidonia?*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 61).

Christchurch MC, R.W. Fereday; HT ♂ labelled "ex Typicum specimen" by Guenée, BMNH.

Hudson 1898, p. 55; 1928, p. 107; as synonym.

➤ Geometridae, *Asaphodes*

**adonis** Hudson, 1898, p. 63 (*Xanthorhoe*)  
?Castle Hill MC or Routeburn OL, G.V. Hudson; type series not located in NMNZ.  
Hudson 1898, p. 63, pl. vii fig. 49; 1928, p. 117, pl. xiv fig. 5; as *Xanthorhoe adonis*.  
Note. Hudson sent 3 specimens of *adonis* (Hudson register no. 423a-c) to Meyrick. The ST series is presumably now in BMNH.

**aegrota** Butler, 1879a, p. 499 (*Selidosema*)  
Wairarapa WA, F.W. Hutton; HT ♂ designated by Butler, BMNH.  
Hudson 1898, p. 64, pl. vii fig. 37; 1928, p. 120, pl. xiv fig. 18; as *Xanthorhoe aegrota*.

**albolineata** Philpott, 1915, p. 194 (*Xanthorhoe*)  
Table Hill SI, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, pp. 120-121, pl. xiv fig. 7, as *Xanthorhoe albolineata*.  
Note. Although both the original spelling and the emendation "albolineata" by Meyrick (1917b, p. 261) are wrong formations (*albolineata* is correct), *albolineata* is retained as the correct original spelling under ICZN Rules, Articles 32a (ii) and 32c. See also *Pseudocoremia albafasciata*, p. 167.

**aphelias** Prout, 1939, p. 248 (*Xanthorhoe*), replacement name for *obscura* Philpott, 1921 but not Butler, 1882  
Note. Prout provided a new name because *Scotocoremia obscura* Butler, 1882, p. 418 (from Chile) "at present stands in *Xanthorhoe*", and Philpott (1928g, p. 484) raised *obscura* from subspecific to specific rank, formally (cf. Prout 1939, p. 248). Prout's description and figure (1939, in Seitz vol. 12, pl. 24 line i) clearly refer to Philpott's type series.

**obscura** Philpott, 1921, p. 338 (*Xanthorhoe*; as subspecies of *helias*). Synonymised by Prout (1939, p. 248).  
Hump Ridge FD, A. Philpott; HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 120, as variety; 1939, p. 410, pl. lvi fig. 10, as species.

**beata** Butler, 1877, p. 397, pl. 43 fig. 6 (*Cidaria*)  
?Castle Hill MC, J.D. Enys; HT ♂ designated by Butler, BMNH.  
Hudson 1898, p. 63, pl. vii fig. 35 and 36, as *Xanthorhoe beata*; 1928, p. 117, pl. xiv fig. 3 and 4, as *Xanthorhoe benedicta*.  
Note. The original illustration (Butler, pl. 43 fig. 6) omits the black speck on the white discal area on the forewing, present on the HT ♂. This omission has led to confusion in treatment of the 'green' *Asaphodes* species.

**benedicta** Meyrick, 1914a, p. 102 (*Xanthorhoe*).  
Synonymised by Prout (1927, p. 77).  
Riccarton Bush, Christchurch MC, E. Meyrick; LT ♂ designated by Fletcher, labelled "Christchurch New

Zealand 18/2/82", "benedicta Meyr." (Meyrick's handwriting), "Xanthorhoe benedicta Meyr. lectotype ♂" (D.S. Fletcher's handwriting), BMNH.  
Hudson 1928, p. 117, pl. xiv fig. 3 and 4, as species.

**camelias** Meyrick, 1888b, p. 58 (*Larentia*)  
Whangarei Heads ND, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1898, p. 65, not figured; 1928, p. 114, pl. xiii fig. 12 and 13; as *Xanthorhoe camelias*.

**campbellensis** Dugdale, 1964, pp. 616-618, fig. 15, 15a, and 19 (*Xanthorhoe*)  
Shoal Point, Campbell Island, K. Rennell; HT ♂ designated by Dugdale, NZAC.  
Dugdale 1971, p. 95, as *Asaphodes campbellensis*.

**cataphracta** Meyrick, §1883d, p. 528; 1884b, pp. 74 (key) and 79 (*Larentia*)  
Arthur's Pass NC/WD, E. Meyrick; LT ♀ here designated, labelled "Arthurs Pass New Zealand 3000 ft 23/1/83", "Xanthorhoe cataphracta Meyr., Type", BMNH.  
Hudson 1898, p. 61, pl. vii fig. 33 and 34; 1928, p. 119, pl. xiv fig. 28 and 29; as *Xanthorhoe cataphracta*.  
Note. LT ♀ is the only survivor of an original ST series of 6 specimens.

**chionogramma** Meyrick, §1883d, p. 528; 1884b, pp. 77 (key) and 82 (*Larentia*)  
Mount Hutt MC, R.W. Fereday; LT ♂ here designated, labelled "Mt Hutt New Zealand RWF 12/81" "Xanthorhoe chionogramma Meyr. type ♂", BMNH.  
Hudson 1898, p. 65, pl. vii fig. 42 and 43; 1928, p. 114, pl. xiii fig. 44 and 45; as *Xanthorhoe chionogramma*.

**chlamydota** Meyrick, §1883d, p. 527; 1884b, pp. 71 (key) and 73-74 (*Epyaxa*)  
Akaroa, Banks Peninsula MC, R.W. Fereday; LT ♂ here designated, labelled "30/1/73 Akaroa bush on spur south of waterfall", "Fereday Colln", CMNZ.  
Hudson 1898, p. 59, pl. vii fig. 28; 1928, p. 109, pl. xiii fig. 39, as *Xanthorhoe chlamydota*.  
Note. This species may prove not to be an *Asaphodes* (R.C. Craw, pers. comm.).

**chlorocapna** Meyrick, 1925a, p. 271 (*Xanthorhoe*)  
Mangere Island, Chatham Islands, S. Lindsay; HT ♂ designated by Meyrick, CMNZ.  
Hudson 1928, p. 114, pl. xvi fig. 11, as *Xanthorhoe chlorocapna*.

Note. Craw (1987) based this combination on genital structure.

**cinnabari** Howes, 1912, p. 203 (*Larentia*)  
Nevis CO, W.G. Howes; HT ♂ designated by Howes, BMNH.  
Hudson 1928, p. 111, pl. xi fig. 44, as *Xanthorhoe cinnabari*.  
**obsoleta** Prout, 1939, p. 251 (as aberration of *Xanthorhoe cinnabari*). New synonymy.  
Nevis CO, W.G. Howes; HT ♂ designated by Prout, BMNH.

Note. Prout's "ab. *obsoleta*" is a very faintly marked specimen collected at the same place and on the same date as HT *cinnabari*.

***crotnoena*** Clarke, 1934, pp. 11–12 (*Xanthorhoe*) new combination

Waiho Gorge WD, C.E. Clarke; HT ♂ designated by Clarke, AMNZ.

Hudson 1939, p. 410, pl. lvi fig. 3 and 4, as *Xanthorhoe crotnoena*.

Note. This species is an obvious member of the *Asaphodes clarata* group on colour pattern, antennal structure, and valval appendage characters.

***clarata*** Walker, 1862a, p. 1197 (*Larentia*)

Waikouaiti DN, P. Earl; HT ♀ unique, BMNH.

Hudson 1898, pp. 61–62, pl. vii fig. 31 and 32; 1928, pp. 118–119, pl. xiv fig. 26 and 27; as *Xanthorhoe clarata*.

***pyramaria*** Guenée, 1868, p. 93 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 79) [mid Canterbury MC], R.W. Fereday; no Guenée specimens in BMNH.

Hudson 1898, p. 61; 1928, p. 118; as synonym.

***cosmodora*** Meyrick, 1888b, p. 57 (*Larentia*)

[Gordon's Pyramid], Mount Arthur NN, E. Meyrick; HT ♀ unique, BMNH.

Hudson 1898, p. 62, not figured; 1928, p. 115, not figured; as *Xanthorhoe cosmodora*, "possibly the other sex of ... *bryopis*".

***bryopis*** Meyrick, 1888b, pp. 57–58 (*Larentia*)

[Gordon's Pyramid], Mount Arthur NN, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 4500 ft 16/1/86", "Xanthorhoe bryopis Meyr. Type ♂", BMNH.

Hudson 1898, p. 62, not figured; 1928, p. 115, pl. xliv fig. 5; as *Xanthorhoe bryopis*.

Note. Both sexes have been collected together on Mt Owen NN. The differences in pattern noted by Meyrick are not constant, and therefore the above synonymy is proposed.

***declarata*** Prout, 1914, p. 122 (*Xanthorhoe*)

Ben Lomond OL, W.G. Howes; HT ♂ designated by Prout, BMNH.

Hudson 1928, p. 119, pl. xiv fig. 25, as *Xanthorhoe declarata*.

Note. Hudson (1928) was mistaken in reporting that J.H. Lewis discovered this species on the Old Man Range CO.

***dionysias*** Meyrick, 1907c, p. 108 (*Xanthorhoe*)

Old Man Range CO, J.H. Lewis; HT ♂ unique, head and abdomen missing, BMNH.

Hudson 1928, p. 120, pl. xiv fig. 9, as *Xanthorhoe dionysias*.

***exoriens*** Prout, 1912, p. 54 (*Larentia*)

Glenorchy OL, W.G. Howes; HT ♂ labelled as "♂ Type" by Prout, BMNH.

Hudson 1928, p. 121, pl. xiv fig. 8.

***frivola*** Meyrick, 1913a, p. 26 (*Xanthorhoe*) new combination

Invercargill SL, A. Philpott; HT ♂ unique, BMNH.

Hudson 1928, pp. 119–120, not figured.

Note. Antennal and genital characters are unequivocally of the *Asaphodes* condition.

***glaciata*** Hudson, 1925, p. 220 (*Xanthorhoe*)

Mount Moltke WD, 5,600 ft, C.E. Clarke; HT ♂ unique, AMNZ.

Hudson 1928, p. 118, pl. lii fig. 11, as *Xanthorhoe glaciata*.

Note. Craw (1987) based this combination on genital structure.

***helias*** Meyrick, §1883d, p. 528; 1884b, pp. 77 (key) and 81 (*Larentia*)

Dunedin DN, E. Meyrick; LT ♂ labelled "New Zealand /80", "Type" [labelled by Prout? See Prout (1939, p. 248), "Dunedin (loc. typ.)"], BMNH.

Hudson 1898, p. 64, pl. vii fig. 40; 1928, p. 120, pl. xiv fig. 10; as *Xanthorhoe helias*.

***ida*** Clarke, 1926, p. 417 (*Xanthorhoe*)

Eweburn Stream, Mount Ida CO, W.G. Howes; HT ♂ designated by Clarke, AMNZ.

Hudson 1928, p. 116, pl. xi fig. 18, as *Xanthorhoe ida*.

Note. Craw (1987) based this combination on genital structure.

***imperfecta*** Philpott, 1905, p. 330 (*Xanthorhoe*)

West Plains SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 121, pl. xiv fig. 16, as *Xanthorhoe imperfecta*.

***limonodes*** Meyrick, 1888b, pp. 54–55 (*Epyaxa*)

Wellington WN, G.V. Hudson; LT ♂ selected by D.S. Fletcher, labelled "Wellington New Zealand GVH .85", "Epyaxa limonodes Meyr. Lectotype", BMNH.

Hudson 1898, p. 57, pl. vii fig. 46; 1928, p. 116, pl. xiii fig. 32 and 33; as *Xanthorhoe limonodes*.

***mnesichola*** Meyrick, 1888b, p. 56 (*Larentia*)

Mount Arthur NN, [Plateau], 4000–4500 ft, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 4000 ft 18/1/86", "Colostigia mnesichola Meyr. Type ♂", BMNH.

Hudson 1898, p. 60, pl. vii fig. 39; 1928, p. 122, pl. xiv fig. 31 and 32; as *Xanthorhoe mnesichola*.

***nephelias*** Meyrick, §1883d, p. 528; 1884b, pp. 77 (key) and 78 (*Larentia*)

Arthur's Pass NC/WD, 4,600 ft, E. Meyrick; LT ♂ labelled as "♂ type" by Prout, "Xanthorhoe nephelias Meyr. ♂ type", "Arthurs Pass New Zealand 29/1/83", BMNH.

Hudson 1898, p. 61, not figured; 1928, p. 121, pl. xiv fig. 45; as *Xanthorhoe nephelias*.

» Geometridae, *Asaphodes nephelias*

**subflava** Howes, 1917, p. 274 (*Xanthorhoe*). Synonymised by Hudson (1928, p. 121). Arthur's Pass NC/WD, W.G. Howes; location of type material unknown. Hudson 1928, p. 121, as synonym.

**obarata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 33 (*Cidaria*)

Nelson NN, T.R. Oxley; HT ♀ unique, BMNH. Hudson 1898, p. 66 (but not pl. vii fig. 45); 1928, p. 117, pl. xiii fig. 40; as *Xanthorhoe obarata*.

**chorica** Meyrick, 1888b, pp. 58–59 (*Larentia*). Synonymised by Prout (1927, p. 78).

Akaroa, Banks Peninsula MC, R.W. Fereday; HT ♀ unique, CMNZ. Hudson 1898, p. 66, pl. vii fig. 55, as species; 1928, p. 117, as synonym.

Note. Fereday's specimen of *chorica* was collected in "bush above Morgan's".

**omichlias** Meyrick, §1883d, p. 529; 1884b, pp. 85 (key) and 90 (*Pasithea*) new combination

Castle Hill MC, J.D. Enys; LT ♂ here designated, labelled "1878 Castle Hill from J.D. Enys", "Fereday Coll.", CMNZ.

Hudson 1898, p. 76, pl. viii fig. 25; 1928, p. 127, pl. xiv fig. 41; as *Notoreas omichlias*, after Meyrick (§1885h, p. 589; 1886c, p. 184).

Note. Craw (1986, p. 132) places this species in *Asaphodes* on genital characters.

**oraria** Philpott, 1903, p. 248 (*Xanthorrhoe*)

New River, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 121, pl. xiv fig. 30, as *Xanthorhoe oraria*.

**oxyptera** Hudson, 1909, p. 67, pl. ii fig. 23 (*Xanthorrhoe*)

North Arm, Carnley Harbour, Auckland Island, A.A. Dorrien-Smith; HT ♂ unique, NMNZ.

Hudson 1928, p. 122, pl. xii fig. 50, as *Xanthorhoe oxyptera*. Dugdale 1964, p. 617, fig. 16; 1971, pp. 95–97, fig. 47.

**peripheraea** Meyrick, 1905, p. 220 (*Xanthorrhoe*)

Humboldt Mountains OL, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 113, pl. xiii fig. 36, as *Xanthorhoe peripheraea*.

**philpotti** Prout, 1927, pp. 77–78 (*Xanthorrhoe*) new combination

Lake Wakatipu OL, A. Philpott; LT ♂ here designated, labelled "L. Wakatipu New Zealand AP.13", "pt. series mentioned by Meyrick 1914:102", "Lectotype ♂ *Xanthorhoe philpotti* Prout, sel. J.S. Dugdale, 1981", BMNH.

Hudson 1928, p. 116, pl. xiv. fig. 1, as *Xanthorhoe beata*. Prout 1939, p. 265, pl. 26 line h, as *Larentia philpotti*.

Note. See *beata* Butler, above.

**prasiniás** Meyrick, §1883d, p. 528; 1884b, pp. 77 (key) and 81–82 (*Larentia*)

Castle Hill MC, J.D. Enys; LT ♂ here designated, labelled "Castle Hill New Zealand JDE /81", "Larentia prasiniás Meyr. Type ♂" [L.B. Prout's handwriting], BMNH.

Hudson 1898, p. 65, pl. vii fig. 41; 1928, p. 116, pl. xiii fig. 49, as *Xanthorhoe prasiniás*.

**prymnaea** Meyrick, 1911b, p. 73 (*Xanthorrhoe*)

Mount Arthur NN, 3,600–4,200 ft, G.V. Hudson; LT ♂ here designated, labelled "Mt Arthur New Zealand GVH .10", "Xanthorhoe prymnaea Meyr. Type ♂" [L.B. Prout's handwriting], BMNH.

Hudson 1928, p. 118, pl. xiv fig. 6 and 7, as *Xanthorhoe prymnaea*.

**recta** Philpott, 1905, p. 330 (*Xanthorrhoe*)

Ida Valley CO, J.H. Lewis; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 120, pl. xiv fig. 19, as *Xanthorhoe recta*.

**sericodes** Meyrick, 1915a, p. 202 (*Xanthorrhoe*)

Mount Earnslaw OL, G.V. Hudson; LT ♂ here designated, labelled "Mt Earnslaw New Zealand GVH 1.14", "Xanthorhoe sericodes Meyr. ♂ Type" [Prout's handwriting], "Figured in Seitz vol. xii", BMNH.

Hudson 1928, p. 121, pl. xiv fig. 20, as *Xanthorhoe sericodes*. Prout 1939, p. 264, pl. 26 line f, as *Larentia sericodes*.

**apicata** Prout, 1939, p. 264 (*Larentia*; as aberration of *sericodes*). New synonymy.

Mount Earnslaw OL, G.V. Hudson; HT ♂ unique, NMNZ.

Hudson 1928, p. 121, pl. xiv fig. 20, as *Xanthorhoe sericodes*; Prout's type is Hudson's figure.

**stephanitis** Meyrick, 1907c, pp. 107–108 (*Asaphodes*)

Invercargill SL, sandhills, A. Philpott; LT ♂ selected by D.S. Fletcher, labelled "Invercargill New Zealand GVH .06", "Asaphodes stephanitis Meyr. ♂ lectotype DSF 10/47", BMNH.

Hudson 1928, p. 107, pl. xiii fig. 21.

**stinaria** Guenée, 1868, p. 92 (*Camptogramma*)

"Canterbury", R.W. Fereday; ?HT ♂ unique, labelled "Ex Musaeo Ach. Guenée", "Ex Oberthür Coll. Brit. Mus. 1927-1", BMNH.

Hudson 1898, p. 60, pl. vii fig. 29; 1928, p. 122, pl. xiii fig. 14; as *Xanthorhoe stinaria*.

Also 3 undescribed species (NZAC).

• **Austrocidaria** Dugdale, 1971b, p. 97. Type species *Cidaria similata* Walker, by original designation.

Note. Under this name are gathered those species

with a broad, Y-shaped juxta, a diverticulum on the corpus bursae (as depicted in Dugdale 1964, fig. 1–8; 1971, fig. 48–52), and with larvae on *Coprosma* (Rubiaceae). Included are all species previously placed (by Hudson 1928) in *Eucymatoge* of authors.

***anguligera*** Butler, 1879a, p. 506 (*Phibalapteryx*) new combination

Dunedin DN, F.W. Hutton; HT ♂ unique, labelled as ♀, abdomen as Tams prep. no. 203, 1927, BMNH.  
Hudson 1898, p. 47, pl. vi fig. 43, as synonym of *Hydriomena gobiata*; 1928, p. 98, pl. xii fig. 22, as *Eucymatoge anguligera*, after Meyrick (1909a, p. 5).

***arenosa*** Howes, 1911, p. 127 (*Eucymatoge*; as *arenosus*) new combination

Titahi Bay WN, W.G. Howes; HT ♀ designated by Howes, BMNH.

Hudson 1928, p. 98, pl. xii fig. 20, as *Eucymatoge arenosa*.

***bipartita*** Prout, (1941) 1958, p. 389 (*Horisme*; as aberration of *anguligera*) new combination

Wellington WN, W.G. Howes; HT ♂ designated as "Type ♂" by Prout, BMNH.

Hudson 1928, p. 98, pl. xii fig. 23, as *Eucymatoge anguligera* [part]. Note. Males of *anguligera*, *arenosa*, and *bipartita* all have a socketed, sickle-shaped blade in the eversible corematal tufts on abdominal segment 8. As well as differences in colour pattern, there are differences in ♂ juxtal shape and cornutal length between *anguligera* and *bipartita*.

***callichlora*** Butler, 1879a, p. 509 (*Cidaria*)

Dunedin DN, F.W. Hutton; HT ♂ labelled by Butler as "Type", BMNH. Hudson 1898, p. 50, pl. vii fig. 13; 1928, p. 100, pl. xii fig. 47, as *Hydriomena callichlora*.

***harmonica*** Clarke, 1926, p. 417 (*Hydriomena*; as subspecies of *callichlora*). Synonymised by Hudson (1928, p. 100).

Waitati DN, C.E. Clarke; HT ♂ unique, AMNZ.

Hudson 1928, p. 100, pl. xlvi fig. 19, as synonym.

***cedrinodes*** Meyrick, 1911b, p. 72 (*Xanthorhoe*) new combination

Mount Arthur NN, G.V. Hudson; LT ♂ labelled as "type ♂" by Prout, "Mt Arthur New Zealand GVH 2.10", "*Xanthorhoe cedrinodes* Meyr. type ♂", abdomen missing, BMNH.

Hudson 1928, p. 114, pl. xiii fig. 27 and 28, as *Xanthorhoe sedrinodes*.

***undulata*** Philpott, 1913, pp. 76–77 (*Xanthorhoe*). Synonymised by Meyrick (1917b, p. 260).

Tisbury SL, A. Philpott; HT ♂ unique, NZAC.

Hudson 1928, p. 114, as synonym.

***episema*** Prout, 1939, p. 250 (*Xanthorhoe*; as subspecies of *cedrinodes*). New synonymy.

Flagstaff Hill DN, W.G. Howes; HT ♂ designated by Prout, BMNH.

Hudson 1928, p. 114, pl. xiii fig. 27, as a form occurring around Dunedin DN.

Note. *A. cedrinodes* is extremely variable in colour pattern at most localities, and specimens referable to *undulata* and *episema* have been collected from TO to SL.

***gobiata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 2 (*Cidaria*) new combination

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH. Hudson 1928, p. 97, pl. xii fig. 21, as *Eucymatoge gobiata*.

***simulans*** Butler, 1879a, p. 506 (*Phibalapteryx*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).

Dunedin DN, F.W. Hutton; HT ♀ labelled as "♂ type", spurious abdomen glued on, BMNH. Hudson 1928, p. 97, as synonym.

***undulifera*** Butler, 1879a, pp. 506–507 (*Phibalapteryx*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).

Dunedin DN, F.W. Hutton; HT ♂ labelled by Butler as "type", BMNH. Hudson 1928, p. 97, as synonym.

***rivularis*** Butler, 1879a, pp. 507–508 (*Phibalapteryx*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).

Dunedin DN, F.W. Hutton; HT ♂ labelled by Butler as "type", BMNH. Hudson 1928, p. 97, as synonym.

***dryocyma*** Meyrick, 1938, p. 426 (*Eucymatoge*). New synonymy.

Mount Hutt SC, Lawford White; HT ♀ unique, CMNZ. Hudson 1939, p. 406, pl. lxii fig. 13, as *Eucymatoge dryocyma*.

Note. HT *dryocyma* is a good match for *undulifera*.

***gobiata ab. fasciata*** Prout, 1939 (*Horisme*). New synonymy.

Niagara DN, W.G. Howes; HT ♀ designated by Prout, BMNH.

Note. This form, which has the lines bent into rounded teeth between the wing veins, occurs in populations from ND to SL.

***haemophaea*** Meyrick, 1925a, p. 270 (*Hydriomena*)

Chatham Island, S. Lindsay; HT ♂ unique, CMNZ.

Hudson 1928, p. 101, pl. xlviii fig. 20.

***lithurga*** Meyrick, 1911b, pp. 71–72 (*Hydriomena*)

Makara WN, R.M. Sunley; HT ♂ unique, BMNH.

Hudson 1928, p. 103, pl. xii fig. 39, as *Hydriomena lithurga*.

***parora*** Meyrick, §1884c, p. 234; 1885d, p. 63 (*Harpaluce*) new combination

[Riccarton Bush], Christchurch MC, R.W. Fereday; LT ♂ selected by D.S. Fletcher, labelled "Harpaluce parora Meyr. ♂ Lectotype DSF 10/47", "parora Meyr.", "Christchurch New Zealand 18/2/82", "Lectotype", BMNH.

Hudson 1898, p. 56, not figured; 1928, pp. 108–109, pl. xiii fig. 24 and 25.

➤ Geometridae, *Austrocidaria*

***praerupta*** Philpott, 1918, p. 125 (*Hydriomena*)  
Mount Cleughearn, Hunter Mountains FD, A. Philpott;  
HT ♂ designated by Philpott, NZAC.  
Hudson 1928, p. 100, pl. xii fig. 48, as synonym of *callichlora*. Philpott 1928g, p. 484, fig. 1 and 2, as species.

***prionota*** Meyrick, §1883d, p. 528; 1884b, pp. 73–74 (*Arsinoe*)

Castle Hill MC, J.D. Enys; LT ♂ here designated, labelled “1878 Castle Hill from J.D. Enys”, “*Arsinoe prionota* Meyr. ♂ EM 1883” (Meyrick’s handwriting), “Lectotype ♂ *Arsinoe prionota* Meyrick teste J.S. Dugdale”, CMNZ.

Hudson 1898, p. 47, pl. vi fig. 47 (but lacking the external bend on the outer median vein seen in LT ♂); 1928, p. 102, pl. xlvi fig. 29 and 30; as *Hydriomena prionota*.

***similata*** Walker, 1862b, p. 1413 (*Cidaria*)

[Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂ unique, BM Geometridae genitalia slide no. 5350, BMNH.

Hudson 1898, p. 50, pl. vii fig. 14; 1928, pp. 99–100, pl. xii fig. 46, as *Hydriomena similata*. Dugdale 1964, pp. 610–611, fig. 1–8, as *Hydriomena similata*; 1971b, pp. 98–100, fig. 48–53, as *Austrocidaria similata*.

***timarata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 19 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 76).

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen incomplete, posterior part glued on and probably spurious, BMNH.

Hudson 1898, p. 50; 1928, p. 99; as synonym.

***nigrofasciata*** Prout, 1939, p. 264 (*Euphyia*; as aberration of *similata*). New synonymy.

Note. No specimens were found in BMNH.

***stricta*** Philpott, 1915, p. 195 (*Xanthorhoe*) new combination

Bold Peak OL, W.G. Howes; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 119, pl. xiv fig. 23 and 24, as *Xanthorhoe stricta*.

Note. The specimen labelled “♀ type” in NMNZ is a ♂ PT.

***umbrosa*** Philpott, 1917b, p. 241 (*Xanthorhoe*) new combination

Mount Cleughearn FD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 115, pl. xiii fig. 36, as *Xanthorhoe umbrosa*.

***venustatis*** Salmon, 1946, p. 4 (*Hydriomena*) new combination

Lake Gunn FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.

Hudson 1950, p. 85, pl. vii fig. 6, as *Hydriomena venustatis*.

● ***Cephalissa*** Meyrick, §1883d, p. 529; 1884b, p. 93. Type species *Cephalissa siria* Meyrick, by original monotypy.

***siria*** Meyrick, §1883d, p. 530; 1884b, p. 93 (*Cephalissa*)

Dunedin DN, (F.W. Hutton) R.W. Fereday; LT ♂ selected by D.S. Fletcher, labelled “Dunedin New Zealand RWF /81”, “*siria* Meyr.” (Meyrick’s handwriting), “*Cephalissa siria* Meyr. lectotype”, BMNH.

Hudson 1898, p. 51, pl. vi fig. 48; 1928, p. 98, pl. xi fig. 45, as *Hydriomena siria*.

● ***Chloroclystis*** of authors, sections A and C in the sense of Hudson (1928, pp. 89 and 95): ♂ with simple, pilose or evenly setulose antennae

***filata*** Guénée, 1857b, p. 353 (*Eupithecia*)

“Nouvelle Hollandie”; HT ♂ unique, MNHN. New Zealand: adventive, since 1960; throughout.

***albiplaga*** Prout, 1958, p. 409 (*Chloroclystis*; as aberration of *filata*).

Sydney N.S.W., Raynor; HT ♀ so labelled, BMNH. Note. The form with the forewing median area white is also present in New Zealand populations.

***impudicis*** Dugdale, 1964, p. 620, fig. 22, 23, 27, and 30 (*Chloroclystis*)

Beeman Camp, Campbell Island, J.L. Gressitt; HT ♂ designated by Dugdale, NZAC.

Dugdale 1971, p. 109, fig. 69–71 and 75, as *Pasiphila impudicis*.

***inductata*** Walker, 1862b, p. 1322 (*Coremia*)

Auckland AK, D. Bolton; HT ♂ unique, BM Geometridae genitalia slide no. 5466, BMNH.

Hudson 1898, p. 44, pl. vi fig. 17, as species, also as *Chloroclystis indicataria*; 1928, p. 89, pl. xi fig. 5 and 6, also as *Chloroclystis semialbata*. Dugdale 1971, p. 109, as *Pasiphila inductata*.

***subitata*** Walker, 1862b, p. 1362 (*Scotosia*). Synonymised by Meyrick (§1883d, p. 531), as synonym of *inductata*.

Auckland AK, D. Bolton; HT ♀ unique, BM Geometridae genitalia slide no. 5467, BMNH.

Hudson 1898, p. 44, as synonym.

***semialbata*** Walker, 1863a, p. 1708 (*Eupithecia*). Synonymised by Dugdale (1971, p. 109).

Nelson NN, T.R. Oxley; HT ♂ unique, BM Geometridae genitalia slide no. 5376, BMNH.

Hudson 1898, p. 41, as synonym of *bilineolata*; 1928, p. 89, pl. xi fig. 5 and 6, as species.

***indicataria*** Walker, 1863a, pp. 1708–1709 (*Eupithecia*). Synonymised by Meyrick (1913a, p. 23).

[?Hawkes Bay HB or Taupo TO]; W. Colenso; HT ♂ unique, abdomen missing, BMNH.

Hudson 1898, p. 44, pl. vi fig. 17, as species; 1928, p. 89, as synonym of *semialbata*.

**semilineata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 36 (*Cidaria*). Synonymised by Meyrick (1917b, p. 253).

[Nelson NN, T.R. Oxley]; HT ♀ unique, abdomen missing, BMNH.

Hudson 1928, p. 89, as synonym of *inductata*.

**"sp. near suffusa Hudson"** Dugdale, 1964, p. 618 (*Chloroclystis*). Synonymised by Dugdale (1971, p. 109).

Dugdale 1964, p. 618, fig. 21, 24–26, and 28–29; 1971, pp. 109–110, fig. 72–74 and 76; Campbell Island and Auckland Islands populations.

**lichenodes** Purdie, 1887, p. 70 (*Pasiphila*)

Dunedin DN, A. Purdie; ST series not located.

Hudson 1898, p. 44, pl. vi fig. 15 and 16; 1928, pp. 95–96, pl. xi fig. 29 and 30; as *Chloroclystis lichenodes*.

**nereis** Meyrick, 1888b, p. 51 (*Pasiphila*)

Mount Arthur NN, E. Meyrick; LT ♀ here designated, labelled "Mt Arthur New Zealand 4000 ft 16/1/86", "Pasiphila nereis Meyr. Type", abdomen missing, BMNH.

Hudson 1898, p. 43, pl. vi fig. 11; 1928, p. 96, pl. xi fig. 16, as *Chloroclystis nereis*.

**ida** Hudson, 1939, p. 405 (*Chloroclystis*), replacement name for *Chloroclystis minima* Hudson, 1905a, p. 356, but not Warren, 1897b, p. 227 (Queensland). Synonymised by Dugdale (1971, p. 114, as *minima*), following Prout (1927, p. 76). Mount Ida CO, J.H. Lewis; HT ♂ unique, labelled "594a", NMNZ.

Hudson 1928, p. 96, as *Chloroclystis minima*, as synonym; 1939, p. 405, as *Chloroclystis ida*.

**sphragitis** Meyrick, 1888b, pp. 51–52 (*Pasiphila*)

Christchurch MC, E. Meyrick; LT ♂ (labelled as "Type ♂" by Prout) here designated, labelled "Christchurch New Zealand 10/2/83", "Chloroclystis sphragitis Meyr. Type ♂", "Figured in Seitz vol. xii, pl.", BMNH.

Hudson 1898, p. 43, pl. vi fig. 13 and 14; 1928, p. 96, pl. xi fig. 31; as *Chloroclystis sphragitis*.

**testulata** Guenée, 1857b, p. 352 (*Eupithecia*)

"Nouvelle-Hollande"; HT ♀ unique, MNHP.

Hudson 1898, p. 45, pl. vi fig. 19, as *Phriessogonus denotatus*; 1928, p. 89, pl. xi fig. 7, as *Phriessogonus testulatus*.

Note. Prout (1958, p. 418) placed *testulata* in *Chloroclystis* without comment. Holloway (1977, p. 127) supported this affiliation, noting that genital structure indicated a closer relationship with *Chloroclystis* of authors than with *Phriessogonus* Butler, where Meyrick (1888b, p. 53; 1917b, p. 252) placed this species.

**denotata** Walker, 1862b, p. 1361 (*Scotosia*). Synonymised by Prout (1927, p. 75).

[Auckland AK], D. Bolton; LT ♀ here designated, labelled "Type" (by Prout?), "New Zeal. // 54.4", "-37 *Scotosia denotata*", BM Geometridae genitalia slide no. 2949, BMNH.

Hudson 1898, p. 45, as species; 1928, p. 89, as synonym.

**humeralata** Walker, 1862b, p. 1362 (*Scotosia*). Synonymised by Meyrick (1917b, p. 252).

[Auckland AK], D. Bolton; HT ♀ unique, BM Geometridae genitalia slide no. 2947, BMNH.

Not mentioned by Hudson.

**parvulata** Walker, 1863a, pp. 1721–1722 (*Phibalapteryx*). Synonymised by Meyrick (1888b, p. 53). [?Hawkes Bay HB or Taupo TO], W. Colenso; LT ♂ here designated, labelled "Holotype ♂" (by Prout?), "New Zeal. 53-19", "- Phibalapteryx parvulata", abdomen and head missing, BMNH.

Hudson 1898, p. 45; 1928, p. 89, as synonym.

**albiplaga** Prout, 1958, p. 418 (*Chloroclystis*; as aberration of *Chloroclystis testulata denotata*). New synonymy.

Dunedin DN, W.G. Howes; HT ♀ unique, designated by Prout, BMNH.

**irregularata** Prout, 1958, p. 418 (*Chloroclystis*; as aberration of *Chloroclystis testulata denotata*). New synonymy.

Sumner MC, J.W. Campbell; HT ♂ designated by Prout, BMNH.

Note. Prout's two names refer to colour morphs encountered in most populations.

Also 1 unidentified, vagrant species represented by 1 specimen, Kauaeranga Valley CL (NZAC).

● **Dasyuris** Guenée, 1868, p. 92. Type species *Dasyuris partheniata* Guenée, by original designation.

**Statira** Meyrick, §1883d, p. 529; 1884b, p. 90; preoccupied by *Statira* Berthold, 1827 (Coleoptera). Type species *Euclidia hectori* Butler, by subsequent designation (Meyrick 1917b, p. 263).

**Stathmonyma** Meyrick, §1885h, p. 539; 1886c, p. 184; replacement name for *Statira* Meyrick. Synonymised by Hudson (1898, p. 69).

**anceps anceps** Butler, 1877, p. 392, pl. 43 fig. 3 (*Fidonia*)

[?Castle Hill] MC, J.D. Enys; HT ♀ labelled by Butler as "type", wings glued to body, BMNH.

Hudson 1898, pp. 69–70, pl. viii fig. 29; 1928, p. 128, pl. xv fig. 55, as *Dasyuris anceps*.

**anceps grisescens** Prout, 1939, pp. 240–241 (*Dasyuris*; as aberration of *anceps*)

Mount Hector, Tararua Range WN, G.V. Hudson; HT ♀ labelled by Prout as "♀ type", BMNH.

Hudson (1939, p. 411) notes "a dark variety" collected by E.S. West in the Ruahine Range RI.

Note. A specimen collected by E.S. West ("Ruahine Range E S West, 4200 ft. 36") is in BMNH. As well as lacking yellow on the hindwings, *D. a. grisescens* has strong bands anterior to the median vein on the forewing underside.

» Geometridae, *Dasyuris*

**austrina** Philpott, 1928a, pp. 359–360 (*Dasyuris*)  
Bold Peak OL, A. Philpott; HT ♂ designated by Philpott,  
NZAC.

Hudson 1928, p. 128 (part), pl. xv fig. 20, as *Dasyuris*  
*hectori*, small variety; 1939, p. 411, as synonym, a  
“view shared by Mr Meyrick”.

Note. Philpott's species is distinct from *hectori* on genital  
characters, and no forms have been found that are  
intermediate in size, colour pattern, or structure.

**callicrena** Meyrick, §1883d, p. 529; 1884b, pp. 85  
(key) and 87 (*Pasithea*)

Kinloch OL, R.W. Fereday; HT ♀ unique, CMNZ.  
Hudson 1898, p. 73, pl. viii fig. 16, as *Notoreas callicrena*;  
1928, p. 130, pl. xv fig. 32, as *Dasyuris callicrena*.

**catadees** Prout, 1939, p. 241 (*Dasyuris*)  
Mount Peel NN, G.V. Hudson; HT ♂ labelled by Prout  
as “♂ type”, BMNH.  
Not mentioned by Hudson.

**enysii** Butler, 1877, pp. 391–392, pl. 42 fig. 9  
(*Fidonia*)

[Castle Hill MC], J.D. Enys; HT ♀ labelled by Butler as  
“type”, abdomen squashed, BMNH.

Hudson 1898, p. 69, pl. viii fig. 28; 1928, pp. 128–129,  
pl. xv fig. 46.

**homomorpha** Meyrick, §1883d, p. 529; 1884b,  
pp. 90 (key) and 91 (*Stathmonyma*; as *Statira*,  
preoccupied). Synonymised by Meyrick (§1884c,  
p. 234; 1885d, p. 65).

Mount Hutt MC, R.W. Fereday; LT ♂ here designated,  
labelled “Mt Hutt New Zealand RWF 1/81”, “Statira  
homomorpha Meyrick ♂ 1884”, body fungused, BMNH.  
Hudson 1898, p. 69; 1928, p. 128; as synonym.

**fulminea** Philpott, 1915, p. 195 (*Dasyuris*)?  
Bold Peak OL, W.G. Howes; HT ♂ designated by Phil-  
pott, NMNZ.  
Hudson 1928, p. 129, pl. xv fig. 44.

**hectori** Butler, 1877, p. 387, pl. 43 fig. 4 (*Euclidia*)  
[?OL or MC, J. Hector or J.D. Enys]; HT ♀ labelled by  
Butler as “type”, BMNH.

Hudson 1898, p. 70, pl. viii fig. 32; 1928, p. 128, pl. xv  
fig. 21.

**leucobathra** Meyrick, 1911b, p. 59 (*Dasyuris*, in  
error as *Notoreas* – see footnote, Meyrick (1911b)  
p. 59, and addendum, p. 68)

Arthur's Pass NC/WD, G.V. Hudson; LT ♀ here design-  
ated, labelled “Arthur's Pass New Zealand GVH  
12.08”, “This specimen may be one of the 2 syntypes.  
JSD 1981”, BMNH.

Hudson 1928, p. 130, pl. xv fig. 45.

**micropolis** Meyrick, 1929, p. 486 (*Dasyuris*)  
Arthur's Pass NC/WD, 5,000 ft, G.V. Hudson; LT ♂ here

designated, labelled “Arthur's Pass New Zealand GVH  
5000' 1.27”, “*Dasyuris micropolis* Meyr. type ♂”,  
BMNH.

Hudson 1939, p. 412, pl. lvi fig. 21.

**octans** Hudson, 1923c, p. 179 (*Dasyuris*)  
Hunter Mountains FD, S. Lindsay; LT ♂ here desig-  
nated, labelled “1071a”, NMNZ.  
Hudson 1928, p. 128, pl. li fig. 15.

**partheniata** Guénée, 1868, p. 93 (*Dasyuris*)  
[?Mount Hutt] MC, R.W. Fereday; HT ♂ unique, BMNH.  
Hudson 1898, p. 70, pl. viii fig. 30 and 31; 1928, p. 129,  
pl. xv fig. 47, as *Dasyuris partheniata*.

**pluvia** Hudson, 1928, p. 129 (*Dasyuris*)  
Field Peak, Tararua Range WN, Stella Hudson; LT ♂  
here designated, labelled “350a”, NMNZ.  
Hudson 1928, p. 129, pl. xv fig. 31.

**strategica** Meyrick, §1883d, p. 529; 1884b, pp. 85  
(key) and 87 (*Pasithea*)  
Lake Guyon MB/BR, W.T.L. Travers; HT ♂ unique,  
CMNZ.

Hudson 1898, p. 73, pl. viii fig. 15, as *Notoreas strategica*;  
1928, p. 130, pl. xv fig. 49, as *Dasyuris strategica*, ♀.

**transaurea** Howes, 1912, p. 203 (*Dasyuris*)  
Garvie Range, near Nevis CO, W.G. Howes; LT ♂  
labelled by L.B. Prout as “Type *transaureus* Howes”,  
BMNH.

Hudson 1928, p. 130, pl. xv fig. 40, as *D. transaurea*.

Also 2 undescribed species (NZAC).

• **Elvia** Walker, 1862b, pp. 1165 (key) and 1430.  
Type species *Elvia glaucata* Walker, by original  
monotypy.

**glaucata** Walker, 1862b, p. 1430 (*Elvia*)  
[Nelson NN], T.R. Oxley; HT ♀ unique, abdomen miss-  
ing, BMNH.

Hudson 1898, p. 46, pl. vi fig. 23 and 24; 1928, pp. 87–  
88, pl. xii fig. 14 and 15.

**dovonani** Felder & Rogenhofer, 1875, pl. cxxxii  
fig. 5 (*Elvia*). Synonymised by Meyrick (§1883d,  
p. 527; 1884b, p. 65).

[Nelson NN, T.R. Oxley]; HT ♂ unique, abdomen miss-  
ing, BMNH.

Hudson 1898, p. 46; 1928, p. 87; as synonym.

• **Epicyme** Meyrick, §1885h, p. 589; 1886c, p. 184;  
replacement name for *Hippolyte* Meyrick

**Hippolyte** Meyrick, §1883d, p. 526; 1884b, pp. 58  
(key) and 60; preoccupied by *Hippolyte* Leach,  
[1814] 1830 (Crustacea). Type species *Ptychopoda?*  
*rubropunctaria* Doubleday, by monotypy.

***rubropunctaria*** Doubleday, 1843, p. 287  
(*Ptychopoda?*)

[Auckland AK, A. Sinclair]; LT ♂ labelled as "Cotype" by A.G. Butler, BMNH.  
Hudson 1898, p. 51, pl. vi fig. 35; 1928, p. 104, pl. xi fig. 39 and 40; as *Euchoeca rubropunctaria*, after Meyrick (1917b, p. 256).

***risata*** Guenée, 1857a, p. 438 (*Asthena*). Synonymised by Meyrick (§1883d, p. 526; 1884b, p. 60). Tasmania, ?collector; HT with abdomen, antennae, and frenulum missing, BMNH.  
Hudson 1898, p. 51; 1928, p. 104; as synonym.

***mullata*** Guenée, 1868, p. 42 (*Asthena*). Synonymised by Meyrick (§1883d, p. 526; 1884b, p. 60). [Christchurch MC], R.W. Fereday; HT ♀ so labelled by L.B. Prout, antennae and abdomen missing, BMNH.  
Hudson 1898, p. 51; 1928, p. 104; as synonym.

• ***Epiphryne*** Meyrick, §1883d, p. 526; 1884b, pp. 58 (key) and 60. Type species *Cidaria undosata* Felder & Rogenhofer, by original monotypy.

***Hermione*** Meyrick, §1883d, p. 526; 1884b, p. 61; preoccupied by *Hermione* de Blainville, 1828 (Vermes). Type species *Hermione xanthaspis* Meyrick. Synonymised under *Venusia* of authors by Hudson (1898, p. 54).

***Panopaea*** Meyrick, §1883d, p. 527; 1884b, pp. 58 (key) and 62. Type species *Cidaria verriculata* Felder & Rogenhofer. Synonymised under *Venusia* of authors by Hudson (1898, p. 53).

***Aulopola*** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Hermione* Meyrick.

***Pancyma*** Meyrick, §1885h, p. 589; 1886c, p. 184; unnecessary replacement name for *Panopaea* Meyrick, not preoccupied by *Panopaea* Lamarck, 1818 nor *Panopaea* Adams, 1856 (both incorrect subsequent spellings of *Panopea* Ménard de la Groye, 1807 (Mollusca), nor by *Panopaea* Felder, 1861, incorrect subsequent spelling of *Panopea* Hübner, 1819 (Fletcher 1979, p. 152)).

***charidema*** ***charidema*** Meyrick, 1909b, p. 70  
(*Venusia*)

Auckland Island, G.V. Hudson; LT ♂ here designated, labelled "Auckland I. New Zealand GVH .08", "Paravicini Coll. BM 1937-383", BM Geometridae genitalia slide no. 5381, BMNH.

Hudson 1928, p. 105, pl. xiii fig. 6 and 7, as *Venusia charidema*.

***charidema autocharis*** Meyrick, 1924a, p. 202  
(*Venusia*)

Mount Ruapehu TO, 4,000 ft, G.V. Hudson; LT ♂ here designated, labelled "Mt Ruapehu New Zealand GVH 4000' 1.22", "Paravicini Coll. BM 1937-383", BMNH.  
Hudson 1928, p. 105, not figured; 1939, p. 407, pl. lxiii fig. 25 (larva); as *Venusia autocharis*.

***undosata*** Felder & Rogenhofer, 1875, pl. cxxviii fig. 2 (*Cidaria*)

[Nelson NN, T.R. Oxley]; ST ♀, abdomen eroded, BMNH [but Prout notes under ST "Felder's fig. seems to be from the 'Type' in Mus. Wien"].

Hudson 1898, p. 54, pl. vi fig. 33 and 34; 1928, pp. 105-106, pl. xiii fig. 1-4, as *Venusia undosata*, after Meyrick (1917b, p. 257).

***citrinata*** Warren, 1903, p. 265 (*Epiphryne*). Synonymised by Meyrick (1917b, p. 257).

Invercargill SL, [?W.G. Howes]; HT ♀ designated by Warren, BMNH.

Hudson 1928, p. 105, as synonym.

***verruculata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 20 (*Cidaria*)

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.

Hudson 1898, p. 53, pl. vi fig. 30 and 31; 1928, p. 104, pl. xiii fig. 9 and 10, as *Venusia verruculata*.

***xanthaspis*** Meyrick, §1883d, p. 526; 1884b, pp. 59 (key) and 61 (*Hermione*)

Lake Guyon MB/BR, R.W. Fereday; LT ♂ here designated, labelled "Lake Guyon New Zealand RWF /81", "Paravicini Coll. BM 1937-383", BMNH.

Hudson 1898, p. 54, pl. vi fig. 32; 1928, p. 106, pl. xiii fig. 5; as *Venusia xanthaspis*.

• ***Epyaxa*** Meyrick, §1883d, p. 527; 1884b, pp. 58 (key) and 71. Type species *Cidaria rosearia* Doubleday, by subsequent designation (Meyrick 1917b, p. 258).

Note. This genus was reinstated by Craw (1987), who included 6 Australian species.

***lucidata*** Walker, 1862a, p. 1200 (*Larentia*)

[Auckland AK], A. Sinclair; HT ♀ unique, BMNH.  
Hudson 1928, p. 112, not figured, as *Xanthorhoe lucidata*.

***robustaria*** Walker, 1862b, p. 1320 (*Coremia*). Synonymised by Prout (1927, p. 77).

[Auckland AK], D. Bolton; HT ♂ unique, BMNH.  
Hudson 1928, p. 112, as synonym; 1928, p. 111, pl. xi fig. 43, as *Xanthorhoe practica*.

***plurimata*** Walker, 1862b, p. 1321 (*Coremia*). Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 64).

[Auckland AK], D. Bolton; HT ♀ unique, BMNH.  
Hudson 1928, p. 112, as synonym.

***officiosa*** Meyrick, 1910b, p. 69 (*Hydriomena*). Synonymised by Dugdale (1973a, p. 200).

Raoul Island, Kermadec Islands, Wallace; HT ♀ unique, BM Geometridae genitalia slide no. 10588, BMNH.

Not mentioned by Hudson.

***practica*** Meyrick, 1911b, p. 72 (*Xanthorhoe*). Synonymised by Prout in Hudson (1939, p. 408).

Motueka NN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 111, pl. xi fig. 43, as *Xanthorhoe practica*.

» Geometridae, *Epyaxa*

**rosearia** Doubleday, 1843, p. 285 (*Cidaria*)  
[Auckland AK], A. Sinclair; 3 ST ♂♂, all lacking the abdomen, all labelled "Cidaria rosearia Doubleday det. D.S. Fletcher 1978, SYNTYPE", "42-55 // New Zealand", BMNH.

Hudson 1898, pp. 57–58, pl. vii fig. 22 and 23; 1928, pp. 110–111, pl. xiii fig. 41 and 42; as *Xanthorhoe rosearia*.

**subductata** Walker, 1862a, p. 1198 (*Larentia*).  
Synonymised by Prout (1927, p. 77).  
[Nelson NN], T.R. Oxley; HT ♀ unique, BM Geometridae genitalia slide no. 10581, BMNH.

Hudson 1898, p. 57, not figured; 1928, p. 112 (but pl. xiii fig. 34 does not represent a topotypic specimen); as species.

**ardularia** Guenée, 1868, p. 63 (*Coremia*).  
Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 71).  
[Christchurch MC], R.W. Fereday; ST ♂ in BMNH.  
Hudson 1898, p. 57; 1928, p. 110; as synonym.

**inamanearia** Guenée, 1863, pp. 63–64 (*Coremia*).  
Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 71).  
[Christchurch MC], R.W. Fereday; ST ♀ in BMNH.  
Hudson 1898, p. 57; 1928, p. 110; as synonym.

**homalocyma** Meyrick, 1902c, pp. 274–275 (*Xanthorhoe*).  
Synonymised by Dugdale (1971, p. 101).  
Chatham Islands, J. Fougeré; HT ♂ so labelled, CMNZ.  
Hudson 1928, p. 112, pl. xi fig. 36, as *Xanthorhoe homalocyma*.

**venipunctata** Walker, 1863a, p. 1666 (*Panagra*)  
[Auckland AK], D. Bolton; HT ♂ unique, BM Geometridae genitalia slide no. 10583, BMNH.

Hudson 1898, p. 64, pl. vii fig. 38, as *Xanthorhoe lucidata*; 1928, p. 112, pl. xiii fig. 35, as *Xanthorhoe venipunctata*.

**psamatodes** Meyrick, §1883d, p. 528; 1884b, pp. 77 (key) and 81 (*Larentia*).  
Synonymised by Meyrick (§1884, p. 234; 1885d, p. 64, as synonym of *lucidata* not of Walker; 1917b, p. 259, as synonym of *venipunctata* Walker).

Wellington WN or Dunedin DN, F.W. Hutton; ST series not found, OMNZ.

Hudson 1898, p. 64; 1928, p. 112; as synonym.

● **Gingidiobora** Craw, 1987, p. 291. Type species *Scotosia subobscurata* Walker, by original designation.

**nebulosa** Philpott, 1917b, p. 241 (*Xanthorhoe*)  
Coverham MB, H. Hamilton; HT ♂ designated by Philpott, NMNZ.  
Hudson 1928, p. 115, pl. xiii fig. 26.

**subobscurata** Walker, 1862b, p. 1358 (*Scotosia*)  
[Nelson NN], T.R. Oxley; HT ♀ unique, antennae missing, BMNH.

Hudson 1898, pp. 66–67, not figured; 1928, p. 115, pl. xiii fig. 30; as *Xanthorhoe subobscurata*.

**ascotata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 9 (*Cidaria*).  
Synonymised by Prout (1927, p. 77).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1928, p. 115, as synonym.

**petropolia** Meyrick, §1883d, p. 528; 1884b, pp. 82–83 (*Larentia*).  
Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 64).

Otira Gorge WD, E. Meyrick; LT ♀ here designated, labelled "Otira Gorge New Zealand 1800 ft 25/1/83", "Lectotype J.S.D. 1981", BMNH.  
Hudson 1898, p. 66, as synonym.

● **Helastia** Guenée, 1868, p. 94. Type species *Helastia eupitheciaria* Guenée, 1868, p. 95.  
Note. Craw (1987) redefined *Helastia*.

**alba** Craw, 1987, pp. 278–279 (*Helastia*)  
Queenstown OL, [M.O. Pasco]; HT ♂ designated by Craw, NZAC.

**angusta** Craw, 1987, p. 290 (*Helastia*)  
Lake Moke OL, J.S. Dugdale; HT ♂ designated by Craw, NZAC.

**christinae** Craw, 1987, p. 283 (*Helastia*)  
Roaring Meg Creek, Kawarau Gorge CO, J.S. Dugdale; HT ♂ designated by Craw, NZAC.

Note. The heading '*Helastia christinae* n. sp.' was omitted (printer's error) in the published version, but is implied by captions to fig. 24, 33, 57, 76, and 94, and by the section on etymology, p. 283.

**cinerearia** Doubleday, 1843, p. 286 (*Cidaria?*)  
[Auckland AK], A. Sinclair; ST ♂ so labelled by D.S. Fletcher, BM Geometridae genitalia slide no. 10292, BMNH.

Hudson 1898, p. 67, pl. viii fig. 2; 1928, pp. 112–113, pl. xi fig. 41, as *Xanthorhoe cinerearia*; Craw 1987, fig. 29, 49, 71, and 90.

Note. See *inoperata*, a junior objective synonym.

**infusata** Walker, 1862a, p. 1199 (*Larentia*).  
Synonymised by Meyrick (1917b, p. 259).  
[Auckland AK], A. Sinclair; HT ♀ unique, BM Geometridae genitalia slide no. 10293, BMNH.  
Hudson 1928, p. 112, as synonym.

**invexata** Walker, 1862a, p. 1199 (*Larentia*).  
Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 83).  
[Auckland AK], D. Bolton; ST ♂ so labelled by D.S. Fletcher, BM Geometridae genitalia slide no. 10294, BMNH.

Hudson 1898, p. 67; 1928, p. 112; as synonym.

**inoperata** Walker, 1862a, p. 1201 (*Larentia*).  
Synonymised by Walker (1863a, p. 1703). Objective synonym.

[Auckland AK], A. Sinclair; ST ♂ so labelled by D.S. Fletcher, labelled "Cidaria [sic] inopera Walker syntype ♂", "Cidaria cinerearia Doubleday syntype ♂", BM Geometridae genitalia slide no. 10292, BMNH. Hudson 1898, p. 67; 1928, p. 112, as synonym. Note. See *cinerearia*, above.

**diffusaria** Walker, 1862a, pp. 1201–1202 (*Larentia*). Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 64).

[Auckland AK], D. Bolton; HT ♀ unique, head and abdomen missing, BMNH. Hudson 1898, p. 67; 1928, p. 112; as synonym.

**eupitheciaria** Guenée, 1868, p. 95 (*Helastia*). Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 83).

[Christchurch MC], R.W. Fereday; LT ♂ designated by Craw (1987), abdomen missing, BMNH.

Hudson 1898, p. 67; 1928, p. 112; as synonym.

**adonata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 31 (*Cidaria*). Synonymised by Meyrick (1917b, p. 259).

[?Auckland AK or Waikato WO], F.R. Hochstetter; HT ♂ so labelled, antennae and abdomen missing, BMNH. Hudson 1898, p. 67; 1928, p. 112; as synonym.

Note. Felder & Rogenhofer (1875) state in the caption to their plate 131 "31 Cidaria adonata F & R ♂ Nova Zealandia (Hochstetter) antennae maris spathose breviter pectinato fimbriatae (♀ MC)", i.e., ♀ in Royal Imperial Museum.

**clandestina** Philpott, 1921, p. 338 (*Xanthorhoe*) Arthur's Pass NC/WD, E.S. Gourlay; HT ♂ unique, CMNZ. Hudson 1928, p. 113, pl. xii fig. 38, as *Xanthorhoe clandestina*; Craw 1987, p. 283, fig. 34, 58, 77, and 95.

**corcularia** Guenée, 1868, p. 61 (*Larentia*) [Christchurch MC], R.W. Fereday; HT ♂, BM Geometridae genitalia slide no. 10142, BMNH. Hudson 1898, p. 67, as synonym of *Xanthorhoe cinerearia*; 1928, p. 113, as synonym of *X. semisignata*; Craw 1987, p. 283, fig. 35, 59, 60, 78, 96, and 105.

**infantaria** Guenée, 1868, p. 62 (*Larentia*). Synonymised by Craw (1987).

[Christchurch MC], R.W. Fereday; HT ♀ unique, BM Geometridae genitalia slide no. 10296, BMNH.

Hudson 1898, p. 67; 1928, p. 113; as synonym of *Xanthorhoe cinerearia*.

**cryptica** Craw, 1987, pp. 288–289 (*Helastia*) River Jordan, Paradise OL, R.C. Craw; HT ♂ designated by Craw, NZAC.

**cymozeucta** Meyrick, 1913a, pp. 25–26 (*Xanthorhoe*) Ohakune RI/TO, G.V. Hudson; LT ♂ designated by Craw (1987), BM Geometridae genitalia slide no. 10299, BMNH.

Hudson 1928, p. 118, pl. xv fig. 9 and 10, as *Xanthorhoe cymozeucta*; Craw 1987, p. 289, fig. 42, 68, 85, and 100.

**maoriaria** Hudson, 1939, p. 410 (*Xanthorhoe*). Synonymised by Craw (1987). Wainuiomata WN, G.V. Hudson; LT ♂ designated by Craw (1987), NMNZ. Hudson 1928, p. 410, pl. lvi fig. 22.

**expolita** Philpott, 1917b, p. 240 (*Hydriomena*) Broken River MC, J.H. Lewis; HT ♂ designated by Philpott, NZAC. Hudson 1928, p. 98, pl. xii fig. 42, as *Hydriomena expolita*; Craw 1987, p. 289, fig. 43, 69, 86, and 101.

**farinata** Warren, 1896, p. 388 (*Xanthorhoe*) Wellington WN, [G.V. Hudson]; LT ♂ designated by Craw (1987), bearing a Hudson label "35n", BM Geometridae genitalia slide no. 10145, BMNH. Hudson 1928, p. 113, as *Xanthorhoe farinata*; Craw 1987, p. 288, fig. 25, 40, 66, 83, and 98.

**mutabilis** Craw, 1987, p. 282 (*Helastia*) Riwaka NN, A. Philpott; HT ♂ designated by Craw, NZAC.

**ohauensis** Craw, 1987, pp. 285–286 (*Helastia*) Freehold Range, Lake Ohau MK, S. Lindsay; HT ♂ designated by Craw, CMNZ.

**plumbea** Philpott, 1915, pp. 194–195 (*Xanthorhoe*) Queenstown OL, A. Philpott; HT ♂ designated by Philpott, NZAC. Hudson 1928, p. 113, as *Xanthorhoe plumbea*; Craw 1987, p. 282, fig. 32, 56, 75, and 93.

**salmoni** Craw, 1987, p. 285 (*Helastia*) Homer Tunnel FD, J.T. Salmon; HT ♂ designated by Craw, NMNZ.

**scissa** Craw, 1987, p. 287 (*Helastia*) Mt Tarndale MB, lower slopes, J.S. Dugdale; HT ♂ designated by Craw, NZAC.

**semisignata** Walker, 1862a, p. 1200 (*Larentia*) [Auckland AK], A. Sinclair; HT ♂ unique, head missing, BM Geometridae genitalia slide no. 10138, BMNH. Hudson 1898, p. 67, as synonym of *Xanthorhoe cinerearia*, after Meyrick (§1883d, p. 528; 1884b, p. 83); 1928, p. 113, pl. xiii fig. 37, as species, after Meyrick (1917b, p. 259); Craw 1987, p. 285, fig. 36, 61, 62, 79, and 97.

**punctilineata** Walker, 1862a, p. 1202 (*Larentia*). Synonymised by Meyrick (1917b, p. 259). [Auckland AK], D. Bolton; LT ♂ designated by Craw (1987), BM Geometridae genitalia slide no. 10140, BMNH.

Hudson 1898, p. 67, as synonym of *Xanthorhoe cinerearia*, after Meyrick (§1883d, p. 528; 1884b, p. 83); 1928, p. 113, as synonym of *X. semisignata*.

**dissociata** Walker, 1863a, p. 1734 (*Cidaria*). Synonymised by Meyrick (1917b, p. 259).

➤ Geometridae, *Helastia semisignata*

[Auckland AK], A. Sinclair; HT ♀ unique, BM Geometridae genitalia slide no. 10141, BMNH.  
Hudson entries as for *punctilineata*, above.

***similisata*** Walker, 1863a, p. 1735 (*Cidaria*). Synonymised by Meyrick (1917b, p. 259).  
[Auckland AK], D. Bolton; HT ♂ unique, BM Geometridae genitalia slide no. 10139, BMNH.  
Hudson entries as for *punctilineata*, above.

***siris*** Hawthorne, 1897, p. 283 (*Asaphodes*)  
Wellington WN, E.F. Hawthorne; HT ♂ unique, Hudson label "525c", NMNZ.  
Hudson 1898, p. 55, pl. vii fig. 15, as *Asaphodes siris*; 1928, p. 98, as synonym of *Hydriomena triphragma*; Craw 1987, p. 290, fig. 46, 89, and 103.

***triphragma*** Meyrick, §1883d, p. 528; 1884b, p. 74 (*Cidaria*)  
Blenheim MB, W. Skellon; LT ♂ designated by R.C. Craw (1987, p. 290), head mildewed, BMNH.  
Hudson 1898, p. 49, not figured; 1928, p. 98, pl. xii fig. 40; as *Hydriomena triphragma*; Craw 1987, p. 290, fig. 45, 70, 88, and 102.

● ***Homodotis*** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Eurydice* Meyrick.

***Eurydice*** Meyrick, §1883d, p. 527; 1884b, pp. 58 (key) and 62; preoccupied by *Eurydice* Leach, 1815 (Crustacea). Type species *Eurydice cymosema* Meyrick, by monotypy.

***Harpalyce*** Meyrick, §1883d, p. 527; 1884b, pp. 58 (key) and 63; preoccupied by *Harpalyce* Stephens, 1827 (Geometridae). Type species *Larentia megaspilata* Walker, by subsequent designation (Meyrick 1917b, p. 257).

***Probolaea*** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Harpalyce* Meyrick.

***amblyterma*** Meyrick, 1931a, p. 94 (*Asaphodes*)  
Whangarei ND, S.C. Patterson; HT ♀ labelled as ♂, abdomen missing, BMNH.  
Hudson 1939, p. 408, pl. lvi fig. 18, as *Asaphodes amblyterma*, ♂.

***falcata*** Butler, 1879a, p. 501 (*Larentia*?)  
[Dunedin DN], F.W. Hutton; HT ♂ unique, BMNH.  
Hudson 1928, p. 113, not figured, as *Xanthorhoe falcata*.

***rufescens*** Butler, 1879a, p. 502 (*Larentia*?). New synonymy.  
[Dunedin DN], F.W. Hutton; HT ♂ designated by Butler, BMNH.

Hudson 1898, p. 56, not figured; 1928, p. 108, pl. xiii fig. 17 and 18, as species.

Note. Prout (1927, p. 77) regarded HT *falcata* as a dark *rufescens*. NZAC holdings include specimens as dark as HT *falcata*, and others intermediate between that and HT *rufescens*.

***cymosema*** Meyrick, §1883d, p. 527; 1884b, pp. 59 (key) and 63 (*Eurydice*). Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 63).  
Dunedin DN, F.W. Hutton; LT ♂ here designated, labelled "Dunedin New Zealand FWH /77", "one of orig. 10 specs of *Eurydice cymosema*. FWH = Capt Hutton who coll. originals for Fereday. ?should this be lectotype. See WHTT, DSF 10/47", abdomen missing, BMNH.  
Hudson 1898, p. 56; 1928, p. 108; as synonym.

***megaspilata*** Walker, 1862a, p. 1198 (*Larentia*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, head mildewed, BMNH.

Hudson 1898, pp. 55–56, pl. vii fig. 17–20; 1928, p. 108, pl. xiii fig. 14–16.

***assata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 4 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 63).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.

Hudson 1898, p. 55; 1928, p. 108; as synonym.

***nehata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 6 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 63).  
[Nelson NN, T.R. Oxley]; HT ♀ not located.

Hudson 1898, p. 55; 1928, p. 108, as synonym.  
Note. Felder & Rogenhofer indicate in the caption to their plate 131 fig. 6 that the specimen is in MC, (?)now NHMW.

● ***Horisme*** of authors (Holloway 1979, p. 312) (in part)

***suppressaria*** Walker, 1863a, p. 1721 (*Phibalapteryx*)  
[Nelson NN], T.R. Oxley; LT ♀ here designated, labelled "Phibalapteryx suppressaria", "Figured in Seitz vol. xii", "Auckland N. Zeal. 60.73", "Type" (circular, green-margined label), abdomen missing, BMNH.  
Hudson 1928, p. 112, not figured.

***paucita*** Howes, 1942, p. 277 (*Hydriomena*?). New synonymy.  
Portobello DN, W.G. Howes; HT ♂ designated by Howes, NMNZ.

Hudson 1946, pp. 86–87, pl. iii fig. 7, as *Hydriomena paucita*.

Note. See the discussion by Holloway (1979, pp. 312–314, text-fig. 56(2), pl. 60 fig. 3 and 4). New Zealand *suppressaria* has shorter labial palpi than the Australian *Cidaria clandestinata* Walker (= *Horisme mortuata* (Guenée)), and is almost identical with *Horisme grisearia* Holloway as regards the genitalia and abdominal colour pattern.

● ***Hydriomena*** of authors (Meyrick 1917b, p. 255) (in part)

Note. Species included here are those not removed to *Astrocidaria*, *Cephalissa*, or *Anachloris*.

***arida*** Butler, 1879a, p. 505 (*Melanthisia*)  
[Dunedin DN], F.W. Hutton; HT ♂ unique, BMNH.

- Hudson 1898, pp. 50–51, pl. vii fig. 15; 1928, p. 100, pl. xii fig. 33 and 34.
- chaotica** Meyrick, §1883d, p. 528; 1884b, pp. 76–77 (*Cidaria*). Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 64).
- Arthur's Pass NC/WD, E. Meyrick; LT ♀ so labelled by D.S. Fletcher, BMNH.
- Hudson 1898, p. 50; 1928, p. 100; as synonym.
- canescens** Philpott, 1918, p. 125 (*Hydriomena*) Queenstown OL, M.O. Pasco; HT ♂ unique, not located in Pasco Collection, SMNZ.
- Hudson 1928, p. 101, pl. xii fig. 41.
- clarkei** Howes, 1917, p. 274 (*Chloroclystis*) new combination
- Flagstaff Hill DN, C.E. Clarke; HT ♀ designated by Howes, AMNZ.
- Hudson 1928, pp. 96–97, pl. xii fig. 18, as *Chlorochytis clarkei*.
- Note. Venation and wing pattern exclude *clarkei* from *Chloroclystis*, and it is here placed close to *canescens*, the description of which it fits well.
- deltoidata** Walker, 1862b, pp. 1321–1322 (*Coremia*) [Nelson NN], T.R. Oxley; LT ♂ here designated, labelled “42. *Coremia deltoidata*”, “Auckland New Zeal. 60.73”, BM Geometridae genitalia slide no. 5349, BMNH.
- Hudson 1898, pp. 47–48, pl. vii fig. 1–9 (fig. 8 is close to the LT wing pattern); 1928, pp. 101–102, pl. xii fig. 24–28 (fig. 24 is close to the LT wing pattern).
- inclarata** Walker, 1862b, p. 1411 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Nelson NN], T.R. Oxley; LT ♂ here designated, labelled “75. *Cidaria inclarata*”, “Auckland N. Zeal. 60.73”, “Type” (circular, green-margined label), BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- perductata** Walker, 1862b, p. 1412 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [?Wellington WN], Major Parry; HT ♂ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- congressata** Walker, 1862b, pp. 1412–1413 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Auckland AK], D. Bolton; HT ♀ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- conversata** Walker, 1862b, pp. 1413–1414 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- descriptata** Walker, 1862b, p. 1414 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- bisignata** Walker, 1862b, p. 1415 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Nelson NN], T.R. Oxley; HT ♀ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- congregata** Walker, 1862b, p. 1415 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Nelson NN], T.R. Oxley; HT ♀ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- aggregata** Walker, 1862b, pp. 1415–1416 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♀ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- plagifurcata** Walker, 1862b, p. 1416 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- pastinaria** Guenée, 1868, pp. 64–65 (*Coremia*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Christchurch MC], R.W. Fereday; ST ♂ in BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- inopiatata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 3 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [?Nelson NN], F.R. von Hochstetter; HT ♂ unique, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- monoliata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 8 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.
- perversata** Felder & Rogenhofer, 1875, pl. cxxxii fig. 14 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 70).
- [Nelson NN, T.R. Oxley]; HT ♂ labelled as “Type”, BMNH.
- Hudson 1898, p. 47; 1928, p. 101; as synonym.
- hawthornei** Thierry-Mieg, 1915, p. 63 (*Hydriomena*; as variety of *deltoidata*). New synonymy.
- Springfield MC, E.F. Hawthorne; HT ♂ unique, NMNZ.
- Hudson 1898, p. 48, pl. vii fig. 9, as “a development of another striking variety”.
- Note. Hudson's illustration is of Hawthorne's specimen, listed in Hudson's Register under “42 m var beaten out of flax bush Springfield Canterbury 1200 ft 14 Jan -93 (ex coll. Hawthorne)”.
- hemizona** Meyrick, 1897b, p. 385 (*Hydriomena*) Cape Terawhiti WN, [1,500 ft], G.V. Hudson; HT ♂ unique, BMNH.
- Hudson 1898, p. 48, pl. vii fig. 10; 1928, p. 100, pl. xii fig. 35 and 36.

***purpurifera*** Fereday, §1883d, p. 531; 1884, pp. 119–120 (*Cidaria*)

Mount Hutt MC, R.W. Fereday; LT ♂ here designated, labelled “Fereday collection”, “LECTOTYPE *Cidaria purpurifera* Fereday teste J.S. Dugdale”, CMNZ.

Hudson 1898, pp. 49–50, pl. vii fig. 12; 1928, p. 99, pl. xii fig. 45.

***ochreifera*** Prout, 1939, p. 290 (*Euphyia*; as subspecies of *purpurifera*). **New synonymy.**

Flagstaff Hill DN, W.G. Howes; HT ♂ designated by Prout, BMNH.

Not mentioned by Hudson.

Note. Prout distinguished his subspecies on hindwing colour. NZAC has specimens intermediate for this character.

***rixata*** Felder & Rogenhofer, 1875, pl. cxxii fig. 1 (*Cidaria*)

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Hudson 1898, p. 49, pl. vii fig. 11; 1928, p. 99, pl. xii fig. 43 and 44.

***squalida*** Butler, 1879a, p. 505 (*Cidaria*). Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 75).

Dunedin DN, F.W. Hutton; HT ♂ labelled as “type”, BMNH.

Hudson 1898, p. 49; 1928, p. 99; as synonym.

***liara*** Prout, 1939, p. 290 (*Euphyia*; as subspecies of *rixata*). **New synonymy.**

Titahi Bay WN, W.G. Howes; HT ♂ designated as “Type” by Prout, BMNH.

Not mentioned by Hudson.

Note. Prout distinguished his subspecies on hindwing pattern, and distinguished between North Island (ochreous) and South Island (grey). NZAC has specimens from both North and South islands with ochreous hindwings, and others of intermediate colour.

***subrectaria*** Guenée, 1857b, p. 411 (*Coremia*)

Tasmania, ?collector; HT ♂ unique, MNHP.

Hudson 1928, p. 102, pl. xiii fig. 31.

***casta*** Butler, 1880, pp. 553–554 (*Coremia*). Synonymised by Meyrick (1917b, p. 255).

Blenheim MB, W. Skellon; HT ♀ unique, BMNH.

Hudson 1928, p. 102, as synonym.

Note. The name *subrectaria* might not apply to New Zealand populations, which on colour pattern agree better with Brisbane Qld specimens in BMNH. The other Meyrick synonym listed by Hudson (*Cidaria responsata* Walker) is from South Australia.

● ***Microdes*** Guenée, 1857b, pp. 296–297. Type species *Microdes villosata* Guenée, by subsequent designation (Meyrick 1917b, p. 252); Tasmania.

***epicryptis*** Meyrick, 1897b, p. 384 (*Microdes*) Wellington WN, G.V. Hudson; LT ♂ selected by D.S.

Fletcher, labelled “*Microdes epicryptis* Meyrick lectotype ♂ DSF 1948”, “Wellington New Zealand GVH .96”, “Lectotype”, BMNH.

Hudson 1928, p. 88, pl. xi fig. 2.

Note. The specimens identified as “*Scoparia elaphra* Meyr.” from Antipodes Is by Salmon in Salmon & Bradley (1956, p. 64) are both *M. epicryptis*.

***quadristrigata*** Walker, 1862a, p. 1200 (*Microdes*) [Auckland AK], D. Bolton; HT ♀ unique, BMNH.

Hudson 1928, p. 88, pl. xi fig. 1.

***interclusa*** Walker, 1862a, p. 1202 (*Larentia*). Synonymised by Meyrick (1884b, p. 109).

[?Auckland AK], J.F. Churton; HT ♀ unique, BMNH. Hudson 1928, p. 88, as synonym.

***toriata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 34 (*Microdes*). Synonymised by Meyrick (1917b, p. 252).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH. Hudson 1928, p. 88, as synonym.

***rectilineata*** Hudson, 1898, p. 45 (*Chloroclystis*). Synonymised by Meyrick (1913a, p. 23).

Wellington WN, W.P. Cohen; HT ♀ unique, NMNZ. Hudson 1898, p. 45, pl. vi fig. 22, as species; 1928, p. 88, as synonym.

● ***Notoreas*** Meyrick, §1885h, p. 589; 1886c, p. 184; replacement name for *Pasithea* Meyrick.

***Pasithea*** Meyrick, §1883d, p. 529; 1884b, p. 84; preoccupied by *Pasithea* Oken, 1807 (*Vermes*). Type species *Fidonia perornata* Walker, by subsequent designation (Meyrick 1917b, p. 261).

***Lythria*** of authors, but not Hübner (1823, p. 300) (Meyrick 1917b, p. 263; Prout 1927, p. 78).

Note. The indicated arrangement is that of Craw (1986).

***arcuata*** Philpott, 1921, pp. 338–339 (*Notoreas*)

St Arnaud Range NN/MB, R.R. Grimmett; HT ♀ lacking locality label, but designated by Philpott, NZAC. Hudson 1928, p. 125, but not pl. xiv fig. 39.

***atmogramma*** Meyrick, 1911b, pp. 59–60 (*Notoreas*)

Mount Holdsworth WN, G.V. Hudson; LT ♀ labelled as “♂ type”, BMNH.

Hudson 1928, p. 124, pl. xiv fig. 36 and 37.

***blax*** Prout, 1939, p. 244 (*Notoreas*)

Bold Peak, Humboldt Range OL, W.G. Howes; HT ♂ labelled by Prout as “♂ Type”, BMNH.

Not mentioned by Hudson. Prout (in Seitz, vol. xii) 1939, p. 244, pl. 24 line f.

***chioneres*** Prout, 1939, p. 244 (*Notoreas*)

Obelisk, Old Man Range CO, W.G. Howes; HT ♂ designated by Prout, BMNH.

Not mentioned by Hudson. Prout (in Seitz, vol. xii) 1939, pl. 24 line f.

**chrysopeda** Meyrick, 1888b, p. 48 (*Arctesthes*; as *Arcteuthes*, misspelling)

Mount Arthur NN, 4,000 ft, E. Meyrick; LT ♂ here designated, labelled "Mt Arthur New Zealand 4000 ft 15/1/86", "Arctesthes chrysopeda Meyr. Type ♂", BMNH.

Hudson 1898, p. 68, pl. vii fig. 33 and 34; 1928, p. 132, pl. xv fig. 2 and 3; as *Lythria chrysopeda*.

**galaxias** Hudson, 1928, p. 125 (*Notoreas*)

Old Man Range CO, 4,000 ft, J.H. Lewis; HT ♀ unique, NMNZ.

Hudson 1928, p. 125, pl. xv fig. 29.

**hexaleuca** Meyrick, 1914a, p. 103 (*Dasyuris*)

Ben Lomond OL, A. Philpott; HT ♀ unique, BMNH.

Hudson 1928, p. 125, not figured; 1939, p. 411, pl. lvi fig. 30 (according to Meyrick, from Mount Cook MK).

**ischnocypha** Meyrick, 1905, pp. 221–222 (*Notoreas*)

Craigieburn Range MC, 5,600 ft, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1898, p. 72, pl. viii fig. 27, as *Notoreas isoleuca* (not of Meyrick, q.v.); 1928, p. 126, pl. xv fig. 30, as *Notoreas ischnocypha*.

Note. Meyrick proposed the name *ischnocypha* for the species depicted by Hudson.

**isoleuca** Meyrick, 1897b, p. 386 (*Notoreas*)

Castle Hill MC, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 125, pl. xv fig. 1, as *Notoreas isoleuca*.

**isomoera** Prout, 1939, p. 244 (*Notoreas*)

[Ben Lomond], Queenstown OL, W.G. Howes; HT ♂ designated by Prout, BMNH.

Not mentioned by Hudson. Prout 1939 (in Seitz, vol. xii), pl. 24 line f.

**mechanitis** Meyrick, §1883d, p. 529; 1884b, pp. 85 (key) and 86 (*Pasithea*)

Arthur's Pass NC/WD, 3,100 ft, E. Meyrick; LT ♀ labelled as "Type" by L.B. Prout, BMNH.

Hudson 1898, p. 72, pl. viii fig. 9–11; 1928, p. 124, pl. xiv fig. 38; as *Notoreas mechanitis*, after Meyrick (§1885h, p. 589; 1886c, p. 184).

**niphocrena** Meyrick, §1883d, p. 529; 1884b, pp. 85 (key) and 88 (*Pasithea*)

Arthur's Pass NC/WD, 4,500 ft, E. Meyrick; LT ♀ labelled as "Type" by L.B. Prout, BMNH.

Hudson 1898, p. 74, not figured; 1928, p. 126, pl. xiv fig. 42 and 43; as *Notoreas niphocrena*, after Meyrick (§1885h, p. 589; 1886c, p. 184).

**ortholeuca** Hudson, 1923b, p. 129 (*Notoreas*)

Stoney Peak, Glenorchy OL, F.S. Oliver; HT in F.S. Oliver Collection, lost.

Hudson 1928, p. 125, pl. li fig. 17.

**paradelpha** Meyrick, §1883d, p. 529; 1884b, pp. 85 (key) and 86–87 (*Pasithea*)

Ben Lomond OL, 5,000 ft, E. Meyrick; LT ♂ labelled as "type ♂" by L.B. Prout, BMNH.

Hudson 1898, p. 72, pl. viii fig. 12–14; 1928, pp. 124–125, pl. xiv fig. 40; as *Notoreas paradelpha*, after Meyrick (§1885h, p. 589; 1886c, p. 184).

**perornata** Walker, 1863a, p. 1672 (*Fidonia*)

[?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂ unique, BMNH.

Hudson 1898, p. 72 (but pl. viii fig. 4–8 not a good match with HT), as *Notoreas perornata*, after Meyrick (§1885h, p. 589; 1886c, p. 184); 1928, p. 131, pl. xv fig. 42 (a reasonable match), as *Lythria perornata*, after Prout (1927, p. 78).

Note. Not to be confused with *Lythria perornata* Walker, 1862a, p. 1056, from Tasmania.

**regilla** Philpott, 1928a, p. 360 (*Lythria*)

Dun Mountain NN/MB, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 412, as synonym of *Lythria perornata*.

Prout 1939, p. 247, pl. 24 line h, as *Arctesthes regilla*.

**simplex** Hudson, 1898, p. 74 (*Notoreas*)

Mount Arthur NN, G.V. Hudson; HT ♀ unique, labelled "402a" ["Tableland of Mt Arthur, Jan 7 [18]91, 3400 ft – Hudson's Register"], NMNZ.

Hudson 1898, p. 74, pl. viii fig. 26; 1928, p. 126, pl. xiv fig. 44.

Also 1 undescribed species (NZAC).

• **Orthoclydon** Warren, 1894, p. 393. Type species *Acidalia praefactata* Walker, by original designation.

**chlorias** Meyrick, §1883d, p. 528; 1884b, pp. 77 (key) and 80 (*Larentia*)

Castle Hill MC, E. Meyrick; HT ♂ unique, BMNH.

Hudson 1898, pp. 63 and 64, not figured; 1928, p. 107, pl. xl ix fig. 1 and 2, pl. xiv fig. 46.

**princeps** Hudson, 1903, p. 244, pl. xxx fig. 1 (*Venusia*). Synonymised by Meyrick (1905, p. 220).

[MC], R.W. Fereday; HT ♂ unique, CMNZ.

Hudson 1928, p. 107, as synonym.

**praefactata** Walker, 1861, p. 781 (*Acidalia*)

[Nelson NN], T.R. Oxley; HT ♀ unique, BMNH.

Hudson 1898, pp. 60–61, pl. vii fig. 30, as *Xanthorhoe praefactata*; 1928, pp. 106–107, pl. xiv fig. 21 and 22, as *Orthoclydon praefactata*, following Meyrick (1917b, p. 261).

**subtentaria** Walker, 1863a, pp. 1610–1611 (*Acidalia*). Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 78).

[Auckland AK], A. Sinclair; HT ♀ unique, BMNH.

Hudson 1898, p. 60; 1928, p. 106; as synonym.

➤ Geometridae, *Orthoclydon praefactata*

**absconditaria** Walker, 1863a, p. 1611 (*Acidalia*).  
Synonymised by Meyrick (§1883d, p. 528; 1884b, p. 78).

[?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♀ unique, BMNH.

Hudson 1898, p. 60; 1928, p. 106; as synonym.

**pseudostinaria** Hudson, 1918, p. 61 (*Xanthorhoe*)  
Otira Valley WD, G.V. Hudson; HT ♂ unique, Hudson  
label “767a”, NMNZ.

Hudson 1928, p. 107, pl. xiv fig. 12, as *Orthoclydon pseudostinaria*.

● **Paradetis** Meyrick, §1885h, p. 589; 1886c, p. 184;  
replacement name for *Parysatis* Meyrick.

**Parysatis** Meyrick, §1883d, p. 526; 1884b, pp. 58–  
59; preoccupied by *Parysatis* Thomson, 1868  
(Coleoptera). Type species *Parysatis porphyrias*  
Meyrick, by original monotypy.

**porphyrias** Meyrick, §1883d, p. 526; 1884b, p. 59  
(*Parysatis*)

Otira Gorge WD, E. Meyrick; HT ♀ unique, BMNH.

Hudson 1898, p. 41, pl. vi fig. 36; 1928, p. 109, pl. xi fig.  
37 and 38.

● **Paranotoreas** Craw, 1986, p. 136. Type species  
*Fidonia*(?) *brepbosata* Walker, by original  
designation.

**brepbosata** Walker, 1862, p. 1037 (*Fidonia*?)  
Waikouaiti DN, P. Earl; LT ♀ here designated, labelled  
“New Zeal. 45-30” (circular), “12. Fidonia brephosata”  
(strip cut from Walker’s proof), “Type” (circular,  
green margin), BMNH.

Hudson 1898, p. 75, pl. vii fig. 20–23; 1928, pp. 126–127,  
pl. xv fig. 32 and 33; as *Notoreas brephos*, an unjustified  
emendation by Meyrick (1884b, p. 90 – “I have  
corrected Walker’s barbarously-formed name”). Philpott  
1928g, p. 484, fig. 8, ♂ genitalia.

**catocalaria** Guenée, 1868, p. 62 (*Larentia*).  
Synonymised by Meyrick (§1883d, p. 529; 1884b, p. 89).  
“Canterbury” [?Mount Hutt MC], R.W. Fereday; HT ♂  
unique, BMNH.

Hudson 1898, p. 75; 1928, p. 126; as synonym.

**brepbos** Felder & Rogenhofer, 1875, pl. cxxix  
fig. 5 (*Fidonia*). Synonymised by Meyrick (§1883d,  
p. 529; 1884b, p. 89).

[Nelson NN, T.R. Oxley]; LT ♂ here designated, labelled  
“Novara cxxix f5 Fidonia brephos n ♂ N. Seeld” (purple  
ink), BMNH.

**vulcanica** Meyrick, §1883d, p. 529; 1884b, pp. 85  
(key) and 89 (*Pasithea*). Synonymised by Craw  
(1986, p. 136).

Makatoku HB, E. Meyrick; LT ♀ here designated, labelled

“Makatoku New Zealand 8/3/83”, “Type” (circular, red  
margin), BMNH.

Hudson 1898, p. 75, pl. viii fig. 24; 1928, p. 127, pl. xv  
fig. 35; as *Notoreas vulcanica*, after Meyrick (§1885h, p.  
589; 1886c, p. 184).

Note. There is no sign of the Kaweka Range HB STs at  
BMNH. Prout (1939, p. 243, not figured) stated: “Possibly  
*vulcanica* is only an extreme local form of [*brepbosata*  
Walker?]”.

**ferox** Butler, 1877, p. 392, pl. 42 fig. 8 (*Fidonia*)  
[Castle Hill MC], J.D. Enys; HT ♀ labelled by Butler as  
“type”, BMNH.

Hudson 1898, p. 74, pl. viii fig. 17 (forewings wrong  
colour); 1928, p. 126, pl. xv fig. 34; as *Notoreas ferox*.

**fulva** Hudson, 1905a, p. 357 (*Lythria*)

Wedderburn CO, J.H. Lewis; LT ♂ here designated,  
labelled “546a” [“Wedderburn, summer 1899–1900  
J.H. Lewis” – Hudson’s Register], NMNZ.

Hudson 1928, p. 127, pl. xv fig. 7, as *Notoreas fulva*.

**opipara** Philpott, 1915, p. 196 (*Notoreas*)

Mount Rakeahua SI, A. Philpott; HT ♂ designated by  
Philpott, NZAC.

Hudson 1928, pp. 123–124, pl. xv fig. 16, as *Notoreas  
opipara*.

**zopyra** Meyrick, §1883d, p. 529; 1884b, pp. 85 (key)  
and 89 (*Pasithea*)

Mount Hutt MC, in gully, R.W. Fereday; LT ♀ here desig-  
nated, labelled “Holotype ♀ *Pasithea zopyra* Meyr-  
ick”, “Jan. 1882 Mt Hutt in shingle bed of gully”,  
CMNZ.

Hudson 1898, pp. 74–75, pl. viii fig. 18 and 19, as *Noto-  
reas zopyra*, after Meyrick (§1885h, p. 589; 1886c, p.  
184); 1928, pp. 126–127, as synonym of *Notoreas bre-  
phos*, after Meyrick (1917b, p. 262). Prout 1939, p.  
243, pl. 24 line d, as species, after Philpott (1928g, p.  
484, fig. 7, ♂ genitalia).

● **Pasiphila** Meyrick, §1883d, p. 527; 1884b, pp. 58  
(key) and 66. Type species *Eupithecia?* *bilineolata*  
Walker, by original monotypy.

**Helastiodes** Warren, 1895, p. 110. Type species  
*Eupithecia?* *bilineolata* Walker, by original  
designation.

Note. Gathered here are those *Chloroclystis* species  
with fasciculate ♂ antennae; cf. *Chloroclystis*  
(above) and Dugdale 1971, p. 106.

**acompsa** Prout, 1927, p. 76 (*Chloroclystis*), replace-  
ment name for *Chloroclystis modesta* Philpott,  
1915, p. 193, preoccupied (in Prout’s view) by  
*Chloroclystis modesta* (Warren, 1893, p. 383;  
Sikkim) (as *Calluga modesta*)

Bold Peak OL, C.C. Fenwick; HT ♂ designated by Phil-  
pott (as *Chloroclystis modesta*), NMNZ.

Hudson 1928, p. 94, pl. xv fig. 19, as *Chloroclystis  
acompsa*.

**aristias** Meyrick, 1897b, p. 385 (*Chloroclystis*)  
Mount Peel, Mount Arthur Tableland NN, G.V. Hudson;  
HT ♂ unique, worn, BM Geometridae genitalia slide  
no. 5368, BMNH.  
Hudson 1898, p. 42, pl. vi fig. 21 and 22; 1928, p. 94, pl.  
xi fig. 26 and 27; as *Chloroclystis aristias*.

**bilineolata** Walker, 1862a, p. 1246 (*Eupithecia*?)  
Nelson NN, T.R. Oxley; HT ♀ unique, BM Geometridae  
genitalia slide no. 7274, BMNH.  
Hudson 1898, pp. 41–42 (in part), pl. vi fig. 9 and 10;  
1928, p. 93, pl. xliv fig. 4 (but not pl. xi fig. 8), as  
*Chloroclystis bilineolata*.

Note. Gathered under this name are the intricately patterned green, *Hebe*-defoliating 'species' with two spine-like, unequal cornuti on the aedeagal vesica.

**paralodes** Meyrick, 1913, p. 23 (*Chloroclystis*).  
New synonymy.

[Head of] Lake Wakatipu OL, G.V. Hudson; LT ♂ here  
designated, labelled "Chloroclystis paralodes Meyrick  
Type ♂" (Prout's handwriting), "L. Wakatipu New  
Zealand GVH .07", BM Geometridae genitalia slide no.  
11155, BMNH.

Hudson 1928, p. 92, pl. xi fig. 20 and 21, as species.

**zatricta** Meyrick, 1913a, p. 24 (*Chloroclystis*). New  
synonymy.

Wellington WN, G.V. Hudson; HT ♂ unique, BM Geo-  
metridae genitalia slide no. 11164, BMNH.

Hudson 1928, p. 92, not figured, as species.

Note. Although Hudson (1928) states "I am unable to  
identify this form", Meyrick had pointed out (1913a, p.  
24) that Hudson regarded many of his specimens as forms  
of *antarctica*, which Meyrick split into 3 species in 1913.

**lacustris** Meyrick, 1913a, p. 24 (*Chloroclystis*).  
New synonymy.

Lake Harris, head of Lake Wakatipu OL, G.V. Hudson;  
LT ♂ here designated, labelled "Figured in Seitz vol. xii,  
pl.", "L. Wakatipu New Zealand GVH .06", "Type", head  
missing, BMNH.

Hudson 1928, pp. 92–93, pl. xi fig. 19, as species.

**hudsoni** Prout, 1958, p. 410 (*Chloroclystis*; as  
aberration of *antarctica*). New synonymy.

Arthur's Pass NC/WD, G.V. Hudson; HT ♂ depicted by  
Hudson, NMNZ.

Hudson 1928, pp. 92–93, pl. xi fig. 4, as *Chloroclystis  
bilineolata*.

Note: Arthurs Pass ♂♂ have 2 cornuti on the aedeagal  
vesica.

[Dunedin DN], F.W. Hutton; HT ♂ unique, worn,  
BMNH.

Hudson 1898, p. 41, as synonym of *Chloroclystis bilineo-  
lata*, after Meyrick (§1883c, p. 527; 1884b, p. 66); 1928,  
p. 93, as synonym of *Chloroclystis charybdis*.

**antarctica** Hudson, 1898, p. 42 (*Chloroclystis*).  
West Plains SL, A. Philpott; specimen matching Hud-  
son's illustration not found in NMNZ.  
Hudson 1898, p. 42, pl. vi fig. 20, as *Chloroclystis antarctica*;  
1928, p. 93, pl. xi fig. 8, as synonym of *Chloro-  
chystis bilineolata*, after Meyrick (1913a, p. 24).  
Note. Males from West Plains collected by Philpott agree  
well with Hudson's (1898) illustration, and with HT *char-  
ybris* genitalia.

**cotinaea** Meyrick, 1913a, pp. 24–25 (*Chloroclystis*)  
Masterton WA, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1928, p. 93, not figured; as *Chloroclystis cotinaea*.

**tornospila** Meyrick, 1931a, p. 94 (*Chloroclystis*).  
New synonymy.

Waimarino [National Park] TO, G.V. Hudson; HT ♂  
unique, bearing Hudson label "1171a", BMNH.

Hudson 1939, pp. 403–404, pl. lvi fig. 13, as *Chloroclystis  
tornospila*.

Note. HT ♂♂ of both nominal species share a character-  
istic hindwing shape (rectangular anal angle) and fea-  
tures of the colour pattern on head, patagia, abdomen,  
forewing, and hindwing. Most conspicuous are the E-  
shaped lunule at the apex of the discal cell, and the prom-  
inent pre-tornal blotch between veins *CuA2* and *CuP*.

**dryas** Meyrick, 1891, p. 97 (*Pasiphila*)

Wellington WN, G.V. Hudson; HT ♂ unique, BM Geo-  
metridae genitalia slide no. 5379, BMNH.

Hudson 1898, p. 43, pl. vi fig. 12; 1928, pp. 93–94, pl. xi  
fig. 17, as *Chloroclystis dryas*.

**erratica** Philpott, 1916, pp. 420–421 (*Chloroclystis*)

Mount Cleughearn, Hunter Mountains FD, A. Philpott;  
HT ♂ designated by Philpott, NZAC.

Hudson 1928, pp. 94–95, pl. xi fig. 35, as *Chloroclystis  
erratica*.

**fumipalpata** Felder & Rogenhofer, 1875, pl. cxxxii  
fig. 33 (*Eupithecia*)

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.

Hudson 1928, p. 96, pl. xi fig. 28, as *Chloroclystis  
fumipalpata*.

**maculata** Hudson, 1898, pp. 44–45 (*Chlorocly-  
stis*). Synonymised by Philpott (1926a, p. 388).

[Manners Street], Wellington WN, W.P. Cohen; HT ♀  
unique, NMNZ.

Hudson 1898, pp. 44–45, pl. vi fig. 18, as species; 1928,  
p. 96, as synonym.

**furva** Philpott, 1917b, pp. 239–240 (*Chloroclystis*)  
Mount Cleughearn, Hunter Mountains FD, A. Philpott;

HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 94, pl. xi fig. 34, as *Chloroclystis furva*.

➤ Geometridae, *Pasiphila*

**halianthes** Meyrick, 1907c, p. 107 (*Chloroclystis*)  
[head of] Lake Wakatipu OL, G.V. Hudson; LT ♂ here  
designated, labelled "Chloroclystis halianthes Meyr.  
Type ♂" and "Figured in Seitz vol. xii, pl. ..." (in  
Prout's writing), "L. Wakatipu New Zealand G.V.H.  
.06", BMNH.

Hudson 1928, p. 95, pl. xii fig. 17, as *Chloroclystis*  
*halianthes*.

**rufulitincta** Prout, 1914, p. 123 (*Chloroclystis*).  
Synonymised by Meyrick (1917b, p. 274).

Ben Lomond OL, W.G. Howes; HT ♂ designated as "♂  
type" by Prout, BMNH.

Hudson 1928, p. 95, as synonym.

**heighwayi** Philpott, 1927a, p. 704 (*Chloroclystis*)  
Pukeatua Bush, Banks Peninsula MC, W. Heighway; HT  
♂ designated by Philpott, NZAC.

Hudson 1928, p. 93, not figured, as *Chloroclystis*  
*heighwayi*.

Note. Locality label on HT reads "Kiwi Bush".

**humilis** Philpott, 1917b, p. 240 (*Chloroclystis*)  
Queenstown OL, M.O. Pasco; HT ♂ designated by Phil-  
pott [Pasco Collection, SMNZ, lost], AT ♀ designated  
by Philpott, NZAC.

Hudson 1928, p. 97, pl. xi fig. 9.

**melanocentra** Meyrick, 1934, p. 151 (*Chloroclystis*).  
New synonymy.

Ben Lomond OL, Lawford White; HT ♂ and AT ♀ *in  
copulo*, pinned on a piece of cork, CMNZ.

Hudson 1939, p. 405, pl. lixii fig. 26, as species.

Note. Hudson (1939) stated: "This species seems close to  
*Chloroclystis humilis* Philpott". Examination of type  
specimens and specimens from several localities dis-  
closed no clear differences, hence the synonymy.

**lunata** Philpott, 1912, p. 115 (*Chloroclystis*)  
Wallacetown SL, A. Philpott; HT ♂ designated by Phil-  
pott, NZAC.

Hudson 1928, p. 93, pl. xi fig. 22–25, as *Chloroclystis*  
*lunata*.

Note. *C. vieta* (below) may prove to be synonymous.

**magnimaculata** Philpott, 1915, p. 193 (*Chloroclystis*)  
Queenstown OL, M.O. Pasco; HT ♂ designated by Phil-  
pott, NZAC.

Hudson 1928, p. 95, pl. xi fig. 33, as *Chloroclystis*  
*magnimaculata*.

**rufipellis** Meyrick, 1927b, pp. 313–314 (*Chloro-  
clystis*). New synonymy.

Gollan's Valley WN, G.V. Hudson; HT ♂ unique,  
BMNH.

Hudson 1939, p. 405, pl. lvi fig. 11, as species.

**irabunda** Prout, 1958, p. 410 (*Chloroclystis*; as  
subspecies of *magnimaculata*). New synonymy.

Flagstaff, Dunedin DN, ?collector; HT ♂ designated by  
Prout as "♂ Type", BMNH.

Note. Populations of *P. magnimaculata* are variable for  
intensity of colour pattern; intergrades of all three nomi-  
nal conditions are met with.

**malachita** Meyrick, 1913a, p. 25 (*Chloroclystis*)  
Lake Harris OL, G.V. Hudson; LT ♂ here designated,  
labelled "Chloroclystis malachita Meyr. Type ♂" by  
Prout, "Lake Harris New Zealand G.V.H. .11", head  
loose, BMNH.

Hudson 1928, p. 95, pl. xi fig. 32, as *Chloroclystis*  
*malachita*.

Note. The ♂ antenna has fascicles of cilia, and is not (as  
Hudson states) evenly ciliated.

**luminosa** Philpott, 1915, pp. 192–193 (*Chloro-  
clystis*). Synonymised by Meyrick (1917b, p. 254).  
Ben Lomond OL, A. Philpott, HT ♂ and AT ♀ not  
located in NZAC.

Hudson 1928, p. 95, as synonym.

**melochlora** Meyrick, 1911b, p. 58 (*Chloroclystis*)  
Otira Valley WD, G.V. Hudson; LT ♂ here designated,  
labelled "Chloroclystis melochlora Meyr. Type ♂" by  
Prout, "Otira River New Zealand G.V.H. 12.08",  
BMNH.

Hudson 1928, p. 91, pl. xi fig. 14, as *Chloroclystis*  
*melochlora*.

**muscosata** Walker, 1862a, p. 1246 (*Eupithecia?*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.

Hudson 1898, p. 41, as synonym of *Chloroclystis bilineo-  
lata*; 1928, p. 91, pl. xi fig. 15, as *Chloroclystis*  
*muscosata*.

**cidariaria** Guenée, 1868, p. 62 (*Eupithecia*). Syn-  
onymised by Meyrick (1888b, p. 50).

[Christchurch] MC, R.W. Fereday; type series probably  
in MNHN.

Hudson 1928, p. 91, as synonym.

**aquosata** Felder & Rogenhofer, 1875, pl. cxxxii  
fig. 38 (*Cidaria*). Synonymised by Meyrick (1888b,  
p. 50).

[Nelson NN, T.R. Oxley]; HT ♀ unique, ♂ genitalia glued  
on to base of ♀ abdomen, BMNH.

Hudson 1928, p. 91, as synonym.

**nebulosa** Dugdale, 1971b, pp. 112–114, fig. 77–83  
(*Pasiphila*)

Adams Island, Auckland Islands, K.A.J. Wise; HT ♂ des-  
ignated by Dugdale, NZAC.

**plinthina** Meyrick, 1888b, pp. 49–50 (*Pasiphila*)  
Wellington WN, A. Purdie; HT ♂ unique, BMNH.

Hudson 1928, pp. 90–91, pl. xi fig. 10, as *Chloroclystis*  
*plinthina*.

**punicea** Philpott, 1923, pp. 148–149 (*Chloroclystis*)  
Rowallan FD, A. Philpott; HT ♂ designated by Philpott,  
NZAC.

Hudson 1928, p. 91, pl. xlvi fig. 12, as *Chloroclystis*  
*punicea*.

Note. Both this HT and the ST series of *semochlora* (below) have very short palpi. The whole species-complex includes *bilineolata* and *suffusa*, and requires critical appraisal.

**rivalis** Philpott, 1916, p. 421 (*Chloroclystis*)

Mount Cleughearn, Hunter Mountains FD, A. Philpott; HT ♂ designated by Philpott, head missing, NZAC.

Hudson 1928, p. 90, pl. xv fig. 18, as *Chloroclystis rivalis*.

Note. Hudson depicted a North Island specimen (Ruapehu TO), and noted that "specimens from Mount Egmont [TK] and Mount Ruapehu are smaller and darker than those from the south".

**rubella** Philpott, 1915, pp. 193–194 (*Chloroclystis*)

Bold Peak OL, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 94, pl. xii fig. 19, as *Chloroclystis rubella*.

**sandycias** Meyrick, 1905, pp. 219–220 (*Chloroclystis*)

Wellington WN, G.V. Hudson; LT ♂ here designated, labelled "Chloroclystis sandycias Meyr. Type ♂" by Prout, "Wellington New Zealand G.V.H. /98", BM Geometridae genitalia slide no. 5391, BMNH.

Hudson 1898, p. 41, pl. vi fig. 8, as *Chloroclystis plinthina*; 1928, p. 90, pl. xi fig. 11–13, as *Chloroclystis sandycias*.

**semochlora** Meyrick, 1919, pp. 349–350 (*Chloroclystis*)

Mount Egmont TK, 3,000 ft, G.V. Hudson; LT ♂ here designated, labelled "Chloroclystis semochlora Meyr. Type ♂" by Prout, "Mt Egmont New Zealand G.V.H. 3000' 2.18", BMNH.

Hudson 1928, p. 92, pl. xlvi fig. 3 and 4, as *Chloroclystis semochlora*.

**suffusa** Hudson, 1928, p. 97 (*Chloroclystis*)

Mount Egmont TK, 3,000 ft, M.N. Watt; LT ♂ here designated, labelled "Holotype ♂" by J.T. Salmon, "Egmont 26.1.16 M.N. Watt", NMNZ.

Hudson 1928, p. 97, pl. xii fig. 16, as *Chloroclystis suffusa*.

**urticæ** Hudson, 1939, p. 404 (*Chloroclystis*)

South Karori WN, G.V. Hudson; LT ♂ here designated, labelled "1089h" ["South Karori bred from larva taken on nettle, bred Nov. 7–21 1937" – Hudson's Register], NMNZ.

Hudson 1939, p. 404, pl. liv fig. 1 (larva).

**vieta** Hudson, 1950, p. 85 (*Chloroclystis*)

Mount Egmont TK, 3,200 ft, A. Castle; HT ♀ unique, NMNZ.

Hudson 1950, p. 85, pl. vi fig. 9.

Note. This species and *lunata* Philpott should be critically compared. Although Hudson reports 3,500 ft as the altitude, the label on the HT reads "3200 ft".

Also 6 undescribed species (NZAC).

● **Phriissogonus** Butler, 1882a, p. 94, in the sense of Holloway (1979, p. 317). Type species *Scotosia canata* Walker, 1862b, p. 1357 (= *Larentia latilcostata* Walker, 1862a, p. 1196).

**laticostatus** Walker, 1862a, p. 1196 (*Larentia*)  
Sydney N.S.W., Lambert; ST ♂ labelled as "Type", BM Geometridae genitalia slide no. 8080, BMNH.

New Zealand: first recorded from Nelson NN (Meyrick 1913a, pp. 22–23).

Hudson 1928, pp. 88–89, pl. xi fig. 3 and 4.

● **Poecilasthena** Warren, 1894, p. 394. Type species *Acidalia pulchraria* Doubleday, by original designation.

**Astheniodes** Hampson, 1903b, p. 647. Type species *Astheniodes polycymaria* Hampson, 1903b, p. 648 (= *Astheña subpurpureata* Walker), by original designation; India. Synonymised with *Astheña* of authors by Prout (1927, p. 76).

**pulchraria** Doubleday, 1843, p. 286 (*Acidalia*)  
"New Zealand", E. Dieffenbach; HT ♀ unique, BMNH.  
Hudson 1898, p. 52, pl. vi fig. 37 and 38; 1928, p. 103, pl. xii fig. 29 and 30; as *Astheña pulchraria*.

**ondinata** Guenée, 1857a, p. 438 (*Astheña*). Synonymised by Meyrick (†1883d, p. 527; 1884b, p. 69).

"Tasmanie", ?collector; HT ♂ labelled "ex typicalibus specimenibus", "ex Musaco Ach. Guenée", BMNH.  
Hudson 1898, p. 52; 1928, p. 103; as synonym.

**plurilineata** Walker, 1861, p. 563 (*Chlorochroma*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 69).

"Tasmania", A.J. Smith; HT ♂ unique, BMNH.  
Hudson 1898, p. 52; 1928, p. 103; as synonym.

**schistaria** Walker, 1861, p. 782 (*Acidalia*)  
[Nelson NN], T.R. Oxley; HT ♀ unique, BMNH.  
Hudson 1898, pp. 52–53, pl. vi fig. 40 and 42; 1928, p. 104, pl. xlvi fig. 1 and 2; as *Astheña schistaria*.

**subpurpureata** Walker, 1863a, p. 1588 (*Astheña*)  
[Nelson NN], T.R. Oxley; HT ♀ unique, abdomen missing, BMNH.

Hudson 1898, pp. 52–53 (part), pl. vi fig. 39, 40a, and 41, as synonym of *Astheña schistaria*, after Meyrick (§1883d, p. 527; 1884b, p. 69).

**tuhuata** Felder & Rogenhofer, 1875, pl. cxxviii fig. 5 (*Acidalia*). Synonymised by Prout (1927, p. 76).

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.  
Hudson 1898, p. 52, as synonym of *Astheña schistaria*; 1928, p. 103, pl. xii fig. 31 and 32, as *Astheña subpurpureata*.

**polycymaria** Hampson, 1903b, p. 648 (*Astheniodes*). Synonymised by Prout (1927, p. 76).

» Geometridae, *Poecilasthena subpurpureata*

"India", F. Moore Collection; HT ♂ labelled as "Astheniodes polycymaria Hampson type ♂", BMNH. Hudson 1928, p. 103, as synonym. Note. Prout (1927, p. 76) regards the type locality as erroneous.

• ***Tatosoma*** Butler, 1874, p. 43. Type species *Cidaria tipulata* Walker (as *Cidaria agrionata* var. *tipulata* Walker), by original designation.

***agrionata*** Walker, 1862b, p. 1417 (*Cidaria*) [?Hawkes Bay HB or Taupo TO], W. Colenso; LT ♂ here designated, labelled "86. *Cidaria agrionata*", "53-19 New Zeal.", "Type", BM Geometridae genitalia slide no. 6451, BMNH. Hudson 1928, pp. 85-86, pl. xii fig. 6 and 7.

***collectaria*** Walker, 1862b, p. 1419 (*Cidaria*). New synonymy. [?Hawkes Bay HB or Taupo TO], W. Colenso; LT ♀ here designated, labelled "89. *Cidaria collectaria*", "53-19 New Zeal.", "25", "Type", BM Geometridae genitalia slide no. 6452, BMNH. Hudson 1928, p. 85, as synonym of *Tatosoma tipulata*.

***alta*** Philpott, 1913, p. 76 (*Tatosoma*) Humboldt Range FD, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ. Hudson 1928, p. 86, pl. xii fig. 10 and 11.

***nigra*** Hudson, 1922, p. 196 (*Tatosoma*). New synonymy. Mount Ruapehu TO, 4,000 ft, G.V. Hudson; HT ♀ unique, labelled "1050a", NMNZ. Hudson 1928, p. 87, pl. 1 fig. 22, as species.

***apicipallida*** Prout, 1914, pp. 122-123 (*Tatosoma*) Ben Lomond OL, W.G. Howes; HT ♀ labelled as "*Tatosoma apicipallida* Prout ♀ type", BMNH. Hudson 1928, p. 86, pl. xlii fig. 28 and 29.

***fasciata*** Philpott, 1914, p. 118 (*Tatosoma*) Lake McKenzie, Hollyford Valley OL, M.O. Pasco and J. Speden; HT ♂ designated by Philpott, lost (originally in Pasco Collection, SMNZ). Hudson 1928, p. 86, pl. xii fig. 1, pl. xliv fig. 31.

***lestevata*** Walker, 1862b, pp. 1416-1417 (*Cidaria*) [Nelson NN], T.R. Oxley; LT ♂ here designated, labelled "85. *Cidaria lestevata*", "60-73 Auckland N. Zeal.", "Type", abdomen in gelatin capsule, BMNH. Hudson 1898, pp. 39-40, pl. vi fig. 25; 1928, p. 85, pl. xii fig. 4 and 5.

***ranata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 11 (*Sauris*). Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 67). [Nelson NN, T.R. Oxley]; HT ♂ unique, apex of abdomen missing, BMNH. Hudson 1898, p. 39; 1928, p. 85; as synonym.

***monoviridisata*** Clarke, 1920, p. 35 (*Tatosoma*) Waitati DN, C.E. Clarke; HT ♂ designated by Clarke, AMNZ. Hudson 1928, p. 86, pl. xlii fig. 6 and 7.

***transitaria*** Walker, 1862b, p. 1419 (*Cidaria*) [?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♀ unique, BM Geometridae genitalia slide no. 8086, BMNH.

Hudson 1898, p. 40, as synonym of *Tatosoma agrionata*; 1928, p. 86, as species, after Meyrick (1911b, p. 71).

***timora*** Meyrick, §1884c, p. 234 (listed); 1885d, p. 65, replacement name for *Tatosoma agrionata* in the sense of Meyrick (1884b, p. 68). New synonymy. Christchurch MC, R.W. Fereday; HT ♂ so labelled by Prout, BMNH.

Hudson 1898, p. 40, pl. vi fig. 28 and 29, as species.

***semifasciata*** Prout, 1958, p. 455 (*Tatosoma*; as aberration of *transitaria*). New synonymy.

Dunedin DN, W.G. Howes; HT ♀ labelled as "*Tatosoma transitaria* ab. *semifasciata* ♀ type" by Prout, BM Geometridae genitalia slide no. 3087, BMNH.

***tipulata*** Walker, 1862b, pp. 1417-1418 (*Cidaria*) [?Hawkes Bay HB or Taupo TO], W. Colenso; LT ♂ here designated, labelled "87. *Cidaria tipulata*", "53-19 New Zeal.", "5", "Type", BMNH.

Hudson 1898, p. 40, pl. vi fig. 26 and 27, as *Tatosoma agrionata*; 1928, p. 85, pl. xii fig. 8 and 9, as species.

***inclinataria*** Walker, 1862b, p. 1418 (*Cidaria*). New synonymy.

[Auckland AK]. A. Sinclair; LT ♂ (described as ♀) here designated, labelled "*Cidaria inclinataria* Walk., det. D.S. Fletcher 1972 syntype", "New Zeal. 47-104", abdomen missing, BMNH.

Hudson 1898, p. 40; 1928, p. 85; as synonym of *Tatosoma agrionata*.

***mistata*** Felder & Rogenhofer, 1875, pl. cxxxii fig. 12 (*Sauris*). Synonymised by Prout (1927, p. 75). [Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH. Hudson 1898, p. 40, as synonym of *Tatosoma agrionata*.

***topia*** Philpott, 1903, p. 247, pl. xxxii fig. 3 and 4 (*Tatosoma*; as *topea*)

West Plains, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 87, pl. xii fig. 12 and 13.

Note. The NZAC copy of *Trans. N.Z. Inst.* vol. 35 has "topea" corrected to "topia" in Philpott's handwriting.

• ***Xanthorhoe*** of authors, in part of Meyrick (1917b, p. 258)

Note. Most of the following species were placed in *Helastia* by Dugdale (1971, pp. 101-102), but that genus has been restricted by Craw (1987).

***bulbulata*** Guenée, 1868, p. 94 (*Cidaria*)

[Christchurch] MC, R.W. Fereday; HT ♀ unique, BMNH. Hudson 1898, p. 68, pl. viii fig. 1; 1928, p. 111, pl. xiii fig. 11.

**frigida** Howes, 1946, pp. 145–146, fig. 2  
(*Xanthorhoe*)

Homer FD, T.M. Smith; HT ♂ unique, NMNZ.

**lophogramma** Meyrick, 1897b, p. 386 (*Xanthorhoe*)

Castle Hill MC, G.V. Hudson; HT ♂ unique, head missing, BMNH.

Hudson 1898, p. 59, pl. vii fig. 47 and 48; 1928, p. 110, pl. xiii fig. 45 and 46.

**occulta** Philpott, 1903, p. 248, pl. xxxii fig. 5  
(*Xanthorrhoe*)

West Plains, Invercargill SL, A. Philpott; HT ♂ designated by Philpott, lost; 2 PT ♂♂ designated by Philpott, NZAC.

Hudson 1928, p. 122, pl. xiv fig. 14 and 15.

Note. Meyrick (1934, pp. 151–152) describes the female.

**orophylla** Meyrick, §1883d, p. 527; 1884b, p. 71  
(*Epyaxa*)

Lake Wakatipu OL, 4,000 ft, E. Meyrick; LT ♂ here designated, labelled “Epyaxa orophylla Meyr. ♂ type”, “Lake Wakatipu New Zealand 4000 ft 16/12/82”, BMNH.

Hudson 1898, p. 58, pl. vii fig. 24 and 25; 1928, p. 110, pl. xiii fig. 22 and 23; as *Xanthorhoe orophylla*, after Meyrick (1917b, p. 258, as *X. orophyla*).

**orophylloides** Hudson, 1909, p. 68, pl. ii fig. 12  
(*Xanthorhoe*)

North Arm, Carnley Harbour, Auckland Island, A.A. Dorrien-Smith; HT ♂ labelled “14” [“Carnley Harbour 21.xi.07 Dorrien-Smith” – Hudson’s Register], genitalia preparation no. 124, NMNZ.

Hudson 1928, p. 110, pl. xiii fig. 38.

Note. Dugdale (1971b, p. 103) placed this species in *Helastia*, but Craw (1987) has shown this to be inappropriate.

**subantarctica** Salmon, 1956, p. 80 (*Xanthorhoe*).

Synonymised by Dugdale (1964, p. 618).

“Campbell Island, J.H. Sorensen” [Auckland I. 21 Apr. 1944, Hoskin]; HT ♂ designated by Salmon, NMNZ.

Note. Dugdale (1964, p. 618; 1971, p. 103) also gave erroneous HT locality data. The AT ♀ is from Campbell I.

**semifissata** Walker, 1862b, pp. 1320–1321  
(*Coremia*)

[Nelson NN], T.R. Oxley; LT ♂ here designated, labelled “40. Coremia semifissata”, “Auckland N. Zeal. 60–73”, “Coremia semifissata Walker syntype ♂ det. D.S. Fletcher 1979”, BM Geometridae genitalia slide no. 10596, BMNH.

Hudson 1898, p. 59, pl. vii fig. 26 and 27; 1928, p. 110, pl. xiii fig. 47 and 48; as *Xanthorhoe semifissata*.

**ypsilonaria** Guenée, 1868, p. 64 (*Coremia*). Syn-

onymised by Meyrick (§1883d, p. 527; 1884b, p. 72).

[Christchurch MC], R.W. Fereday; HT ♂ unique, BMNH. Hudson 1898, p. 59; 1928, p. 110; as synonym.

**delicatulata** Guenée, 1868, p. 94 (*Xanthorhoe*).

Synonymised by Meyrick (§1883d, p. 527; 1884b, p. 72).

[Christchurch MC], R.W. Fereday; HT ♀ designated by Guenée as “Typicum specimen”, BMNH.

Hudson 1898, p. 59; 1928, p. 110; as synonym.

—◎—

## Subfamily OENOCHROMINAE (in the sense of Forbes 1948)

● **Adeixis** Warren, 1897a, p. 27. Type species *Adeixis insignata* Warren, 1897a, p. 27, by original designation; Australia.

**Paragyrtis** Meyrick, 1905, p. 222. Type species *Panagra inostentata* Walker, 1861, pp. 1012–1013, by original monotypy. Synonymised by Prout (1910, p. 22).

**griseata** Hudson, 1903, pp. 244–245 (*Dichromodes*) Seaward Moss [E of Invercargill] SL, A. Philpott; 2 STs labelled “543” [“543a,b Seaward Moss, Invercargill Jan. 4–1900 (A. Philpott)” – Hudson Register], not in NMNZ.

Hudson 1928, pp. 133–134, pl. xv fig. 37, as *Adeixis griseata*, after Prout (1910, p. 22, genus) and Philpott (1921, p. 339, species).

Note. Meyrick (1905, p. 223) states: “Seaward Moss, Invercargill, described from one New Zealand and twelve Australian examples ...”. Hudson (Hudson – Meyrick correspondence, p. 183; NZAC) sent Meyrick a specimen labelled “543” and listed it as “Dichromodes griseata Hdsn, poor one”, over which entry Meyrick wrote: “This is a common Australian insect, *Paragyrtis inostentata* Walk. I have not seen it before from New Zealand; it occurs all over Australia”. He repeated these remarks in his covering letter (p. 180). It is likely that 1 ST survives under *inostentata* in BMNH. NZAC has 21 topotypic specimens, collected in 1919 and 1920. Prout (1927, p. 79) also recognised *griseata* and *inostentata* as distinct entities.

● **Dichromodes** Guenée, 1857, p. 320. Type species *Dichromodes ainaria* Guenée, 1857, p. 321, by subsequent designation (Prout 1910, p. 23); Tasmania.

**Cacopsodos** Butler, 1877, p. 395. Type species *Cacopsodos niger* Butler, by monotypy. Synonymised by Meyrick (§1885h, p. 589, as *Cacopsodes*).

**cynica** Meyrick, 1911b, p. 60 (*Dichromodes*) Lyttelton MC, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, p. 134, pl. xv fig. 11.

**gypsotis** Meyrick, 1888b, p. 60 (*Dichromodes*)

Lake Wakatipu OL, E. Meyrick; HT ♀ unique, BMNH.

➤ Geometridae, *Dichromodes gypsotis*

Hudson 1898, p. 78, not figured; 1928, p. 135, pl. xv fig. 15.

*ida* Hudson, 1905a, p. 356 (*Dichromodes*)  
Ida Valley CO, J.H. Lewis; LT ♂ here designated, labelled  
“595a”, NMNZ.  
Hudson 1928, p. 134, pl. xv fig. 17.

*niger* Butler, 1877, p. 395, pl. xlili fig. 4 (*Cacopsodos*)  
Castle Hill MC, J.D. Enys; HT ♂ labelled “Cacopsodos  
niger Butler Type”, tip of abdomen missing, BMNH.  
Hudson 1898, p. 78, pl. viii fig. 40; 1928, pp. 134–135,  
pl. xv fig. 13 and 36; as *Dichromodes nigra*.

*simulans* Hudson, 1905, p. 356 (*Dichromodes*)  
[Ida Valley CO], J.H. Lewis; 2 ST ♂♂ labelled “729a”  
and “729b” [“Ida Valley, 1905 (J.H. Lewis)”—  
Hudson’s Register], NMNZ.  
Hudson 1928, p. 134, pl. xv fig. 14, as from Old Man  
Range CO, 4000 ft.

*sphaeriata* Felder & Rogenhofer, 1875, pl. cxxxii fig.  
14 (*Cidaria*)  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1898, p. 67, as synonym of *Xanthorhoe cineraria*;  
1928, p. 134, pl. xv fig. 12, as species.

*petrina* Meyrick, 1892, p. 216 (*Dichromodes*).  
Synonymised by Prout (1912, p. 53; 1927, p. 79).  
Wellington WN, G.V. Hudson; LT ♂ here designated,  
labelled “Dichromodes petrina Meyr. ♂ type” by Prout,  
“Wellington New Zealand G.V.H. /90”, BMNH.  
Hudson 1898, pp. 78–79, pl. viii fig. 39, as species; 1928,  
p. 134, as synonym.

● *Samana* Walker, 1863b, p. 197. Type species  
*Samana falcatella* Walker, by original monotypy.

*acutata* Butler, 1877, p. 401 (*Samana*)  
[Christchurch MC], J.D. Enys; HT ♂ unique, head and  
abdomen missing, BMNH.  
Hudson 1898, p. 76, not figured; 1928, p. 133, pl. xv fig.  
39.

*falcatella* Walker, 1863b, p. 197 (*Samana*)  
[?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♀  
unique, antennae missing, BMNH.  
Hudson 1898, p. 76, pl. viii fig. 36; 1928, p. 133, pl. xlvi  
fig. 25.

● *Theoxena* Meyrick, §1883d, p. 526; 1884b, p. 56.  
Type species *Panagra scissaria* Guénée, by original  
monotypy.

*scissaria* Guénée, 1868, p. 43 (*Panagra*)  
[Christchurch MC], R.W. Fereday; LT ♂ here designated,  
labelled “ex Typicalibus Specimenibus”, “Schissaria  
Gn Nlle Zel.”, BM genitalia slide no. 9891, BMNH.

Hudson 1898, p. 79, pl. viii fig. 41; 1928, p. 133, pl. xv  
fig. 38.

● *Xyridacma* Meyrick, 1888b, p. 60. Type species  
*Hemerophila hemipteraria* in the sense of Meyrick  
(1888b), not of Guénée (1857) (= *Xyridacma*  
*veronicae* Prout, below), by original monotypy. See  
Fletcher (1979, p. 216).

*Lyrcea* Walker, 1860, p. 259, preoccupied by  
*Lyrcea* Adams, 1854 (Mollusca). Type species *Lyrcea*  
*alectoraria* Walker, by original monotypy. Synonymised  
by Meyrick (1917b, p. 266, under *Eprrhanthis* of authors, but not Hübner, 1823,  
p. 296).

*Xynonia* Prout, 1910, pp. 8 (key) and 65,  
replacement name for *Lyrcea* Walker.  
Note. The name *Xyridacma* involves misidentification  
of a type species, and should be referred,  
under Article 70(a) of the Code, to the Commission  
on Zoological Nomenclature.

*alectoraria* Walker, 1860a, p. 259 (*Lyrcea*)  
[Auckland AK], D. Bolton; HT ♀ unique, BMNH.  
Hudson 1898, p. 80, pl. viii fig. 44–47; 1928, p. 136, pl.  
xvi fig. 5–8; as *Eprrhanthis alectorara*, apparently the  
first use of this combination.  
Note. Meyrick (§1883d, p. 530; 1884b, p. 95) synonymised  
?Aspilates primaria Walker, 1862a, p. 1076, and  
*Endropia primaria* Walker, [1863], p. 1506. Both species  
are of unknown provenance, have pectinate ♂  
antennae (simple in *Xyridacma*), and are irrelevant to  
*Xyridacma* and the New Zealand fauna. Meyrick  
(§1884c, p. 234) later regarded them as unidentifiable.

*octomaculata* Thierry-Mieg, 1915, p. 63 (*Eprrhanthis*;  
as variety of *alectoraria*). Synonymised by  
Hudson (1928, p. 136).

Wellington WN, G.V. Hudson; LT ♀ here designated,  
labelled “1117e” [“Seivers Hill, Terawhiti Dec. 26 1891”  
— Hudson Register], NMNZ.  
Hudson 1898, p. 80, pl. viii fig. 47, as “one very well  
marked variety”.

Note. The type entity is in one sense Hudson’s illustration,  
but one specimen in his collection (NMNZ) agrees  
well with it, and is here chosen as LT.

*ustaria* Walker, 1863a, pp. 1519–1520 (*Ennomos*)  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1898, p. 80, pl. viii fig. 42 and 43, as synonym  
of *Eprrhanthis alectoraria*; 1928, pp. 135–136, pl. xvi  
fig. 3 and 4, as *Eprrhanthis ustaria*, after Prout (1912,  
p. 53).

*achroiaria* Felder & Rogenhofer, 1875, pl. cxxiii  
fig. 6 (*Amilapis*?). Synonymised by Hudson (1928,  
p. 135).

[Nelson NN, T.R. Oxley]; HT ♂ unique, incomplete, only  
part of thorax, wings, and base of abdomen remaining,  
glued to the head, body, and wing stubs of another  
(?ennomine) geometrid, BMNH.

Hudson 1898, p. 80, as synonym of *Epirrhanthis alectoaria*; 1928, p. 135, as synonym of *Epirrhanthis ustaria*.

**varians** Butler, 1879a, p. 496 (*Lyrcea*). Synonymised by Hudson (1928, p. 135).

Wairarapa WA, F.W. Hutton; LT ♀ here designated, labelled "Lyrcea varians Butler. Type New Zealand 79.19", "Wairarapa", BMNH.

Hudson 1898, p. 80, as synonym of *Epirrhanthis alectoaria*; 1928, p. 35, as synonym of *Epirrhanthis ustaria*.

**hudsoni** Prout, 1920, p. 34 (*Xyridacma*; as aberration of *ustaria*). **New synonymy.**

Dunedin DN, W.G. Howes; HT ♂ designated by Prout, BMNH.

**unilinea** Prout, 1920, p. 34 (*Xyridacma*; as aberration of *ustaria*; as illustration).

**veronicae** Prout, 1934, p. 136, pl. 3b (*Xyridacma*), replacement name for *Xyridacma hemipteraria* in the sense of Meyrick (1888b) but not Guenée (1857)

Auckland AK, E. Meyrick; HT ♂ designated by Prout, BMNH.

Hudson 1898, p. 80, pl. vii fig. 48 and 49; 1928, p. 135, pl. xv fig. 26–28, as *Epirrhanthis hemipteraria*; 1946, p. 90, as *Epirrhanthis* (*Xyridacma*) *veronicae*.

—♂—

### Subfamily STERRHINAE

(in the sense of McGuffin 1967, p. 8)

● **Scopula** Schrank, 1802, p. 162. Type species *Phalaena paludata* Linnaeus, in the sense of Holloway (1979, pp. 293–297); Portugal.

**rubraria** Doubleday, 1843, p. 286 (*Ptychopoda*?). [Auckland AK, A. Sinclair]; ST ♂ so labelled by D.S. Fletcher, abdomen missing. BMNH.

Hudson 1898, p. 77, pl. viii fig. 37 and 38; 1928, p. 132, pl. xv fig. 8, as *Leptomeris rubraria*, apparently the first use of that combination.

**repletaria** Walker, 1861, p. 778 (*Acidalia*). Synonymised by Meyrick (§1884c, p. 234; 1885d, p. 63).

Sydney N.S.W., Lambert; HT ♂ unique. BMNH. Not mentioned by Hudson.

**attributa** Walker, 1861, p. 779 (*Acidalia*). Synonymised by Meyrick (§1884c, p. 234; 1885b, p. 63).

Sydney N.S.W., Lambert; HT ♀ unique. BMNH. Hudson 1898, p. 77; 1928, p. 132; as synonym.

**figlinaria** Guenée, 1857, p. 454, pl. 12 fig. 8 (*Acidalia*). Synonymised by Meyrick (§1883d, p. 526; 1884b, p. 57).

Tasmania, ?collector; HT ♂, BM Geometridae genitalia slide no. 8078, BMNH.

Hudson 1898, p. 77; 1928, p. 132; as synonym.

**acidaliaria** Walker, 1862a, p. 1037 (*Fidonia*?). Synonymised by Meyrick (§1883d, p. 526; 1884b, p. 57).

[Auckland AK]. A. Sinclair; LT ♂ here designated, labelled "13. Fidonia acidaliaria", "45.61 New Zealand", "Type". BMNH.

Hudson 1898, p. 77; 1928, p. 132; as synonym.

### Rejected species of Geometridae

(1) *Campogramma correlata* Walker, 1862b, pp. 1330–1331. "New Zealand. Presented by Dr Sinclair". Does not resemble any New Zealand species, but does resemble specimens from Bathurst, Vict., Australia. Dr Sinclair could have collected it in Australia.

(2) *Panagra promelanaria* Walker, 1863a, p. 1666. Rejected by Meyrick (§1884b, p. 109); Australian. (3) *Eupithecia inexpiata* Walker, 1863a, p. 1708. Wrongly labelled; from Patagonia (BMNH).

—♂—


### Superfamily DREPANOIDEA

#### Family THYATIRIDAE

**Thyatira batis** Linnaeus was introduced to New Zealand for control of blackberry (bramble), but failed to establish. NZAC has specimens reared from the introductory batch.

—♂—

**Superfamily NOCTUOIDEA**  
**Family ARCTIIDAE**



**Arctiidae**  
(179) *Metacrias erichrysa* Meyrick

● ***Metacrias*** Meyrick, 1886e, pp. 83 (key) and 749. Type species *Metacrias erichrysa* Meyrick, by subsequent designation (Kirby 1892, p. 263).

***erichrysa*** Meyrick, 1886e, p. 749 (*Metacrias*)  
Mount Arthur NN, E. Meyrick (reared); LT ♂ here designated, labelled "Mt Arthur New Zealand 4000 ft bred 15/1/86", "Metacrias erichrysa Meyrick, E. Meyrick det. in Meyrick Colln 3/4", BMNH.  
Hudson 1898, p. 4, pl. iv fig. 5; 1928, pp. 43–44, pl. vi fig. 12.

***huttoni*** Butler, 1879a, pp. 487–488 (*Phaos*)  
Queenstown OL, F.W. Hutton; HT ♂ labelled as "Type" by Butler, BMNH.  
Hudson 1898, p. 5, pl. iv fig. 6; 1928, p. 43, pl. vi fig. 1; as *Metacrias huttoni*, after Meyrick (1886e, p. 750).

***strategica*** Hudson, 1889, p. 53 (*Arctia*)  
Richardson Range SC, W.W. Smith; HT ♂ labelled "116a" ["Richardson Range W.W. Smith 9 Feb 1881, beaten from bed of native broom (Carm. flagelliformis), Sth Canterbury" – Hudson's Register], NMNZ.  
Hudson 1898, p. 4, pl. iv fig. 4; 1928, p. 44, pl. vi fig. 9 and 10; as *Metacrias strategica*, after Meyrick (1890, p. 216).  
Note. The Richardson Range ("near Albury" – Meyrick 1890, p. 216) may be part of the Hunters Hills. W.W. Smith would have been searching there for the laughing owl, *Sceloglaux albifacies*.

***hudsoni*** Rothschild, 1914, p. 259, pl. 24g (*Metacrias*; as subspecies of *strategica*). **New synonymy.**  
Invercargill SL, [?W.G. Howes]; 21 STs in BMNH.  
Note. Gibbs (1962, p. 162) includes Invercargill specimens in his concept of *strategica*, and does not record any genital differences as coinciding with colour pattern differences between MC-SC and DN-SL populations.

● ***Nyctemera*** Hübner, [1820], p. 178. Type species *Phalaena lacticinia* Cramer, 1777, pp. 147 and 149, pl. 128 fig. E, by subsequent designation (Hampson 1894, p. 46); India.

***Leptosoma*** Boisduval, 1832, p. 197. Type species *Leptosoma annulatum* Boisduval, by subsequent designation (Meyrick 1912b, p. 93). Junior homonym of *Leptosoma* Desmarest, 1823 (Crustacea). Synonymised by Walker (1854, p. 391).

***amica*** White, 1841, p. 482 (*Agagles*)  
Australia.

New Zealand: sporadic immigrant, interbreeding freely with local *annulata* populations, and giving rise to individuals showing the same range of variation in colour pattern as do R.J. Tillyard's laboratory F1 hybrid specimens in NZAC (Kay 1980, pp. 154–158). Note. Meyrick (1886d, p. 15) discussed differences between *amica* and *annulata*.

***annulata*** Boisduval, 1832, p. 197, pl. 5 fig. 9 (*Leptosoma*)

[New Zealand], Dumont d'Urville's collectors; HT ♂ unique, BMNH.

Hudson 1898, pp. 2–3, pl. iv fig. 1 and 2; 1928, p. 45, pl. vi fig. 3.

Note. Boisduval reported this species from New Guinea (Meyrick 1886e, p. 761), and Walker (1854, p. 392) confused *amica* with it.

***doubledayi*** Walker (Wing MSS.), 1854, p. 392 (*Nyctemera*). Synonymised by Meyrick (1886e, p. 760).

[?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂ labelled "Walker's type of *Nyctemera Doubledayi*", BMNH.

Hudson 1898, p. 2; 1928, p. 45; as synonym.

***tripunctaria*** in the sense of Swinhoe (1916, p. 213; *Deilemerra*), not of Linnaeus (1758, p. 523). Synonymised by Bryk (1937, p. 54).

***antipodea*** Salmon, 1956, p. 64, fig. 5 and 6 (*Nyctemera*; as subspecies of *annulata*). Synonymised by Dugdale (1971b, p. 71).

Antipodes Island, E.G. Turbott; HT ♂ designated by Salmon, AMNZ.

● ***Tyria*** Hübner, [1819], p. 166. Type species *Phalaena Noctua jacobaeae* Linnaeus, by subsequent designation (Hampson 1901, p. 185).

***jacobaeae*** Linnaeus, 1758, p. 511 (*Phalaena Noctua*) Europe.

New Zealand: introduced by man for control of ragwort, *Senecio jacobaea*; after 40 years (1980+) populations are increasing in area, particularly in WN-WA.

● ***Utetheisa*** Hübner, [1819], p. 168. Type species *Phalaena ornatrix* Linnaeus, 1758, p. 511, by subsequent designation (Kirby 1892, p. 345); America.

***lotrix lotrix*** Cramer, 1777, pp. 20 and 149, pl. 109 fig. E and F (*Phalaena*)

Northern Australia and the Pacific.

New Zealand: occasional immigrant; a fresh specimen collected by R.A. Cumber at Paihia ND may have fed locally as a larva, on adventive leguminous weeds (e.g., *Psoralea*) that are abundant there.

***pulchelloides vaga*** Jordan, 1939, p. 284, fig. 228 and 252–254 (*Utetheisa*)

Semplak, Western Java; HT ♂ designated by Jordan, BMNH.

New Zealand: sporadic immigrant, occasionally establishing for a summer but not known to survive the winter; recorded from ND, AK, CL, TK, WN, NN, BR, DN.

Hudson 1898, pp. 3–4, pl. iv fig. 3; 1928, pp. 44–45, pl. vi fig. 3; as *Utetheisa pulchelloides*, after Meyrick (1912b, p. 92).

Note. Meyrick (1890, p. 217) records this as *Deiopeia pulchella* Linnaeus.

—◎—

### Family CTENUCHIDAE (as in Watson *et al.* 1980)

● ***Antichloris*** Hübner, 1818, pp. 9 and 24. Type species *Zygaena eriphia* Fabricius, 1777, p. 276, by subsequent designation (Kirby 1892, p. 158); Surinam.

***viridis*** Druce, 1884, p. 68 (*Antichloris*)

Panama, Volcan de Chiriquí; HT ♂ unique, BMNH. Indigenous to Central America (Mexico – Ecuador), on *Musa*.

New Zealand: cocoons and adults arrive in shipments of Ecuadorian bananas. The possibility of it establishing where ornamental *Musa* cultivars are grown cannot be ruled out.

Note. Field (1975, pp. 2–4 and 14–16) gave life history details and resurrected Druce's generic placement. Previously most authors had referred this species to *Ceramidia* Butler.

—◎—

### Family LYMANTRIIDAE

● ***Teia*** Walker, 1855, pp. 703 (key) and 803–804. Type species *Teia anartoides* Walker, by original monotypy.

***anartoides*** Walker, 1855, p. 804 (*Teia*)

South-eastern Australia.

New Zealand: live adults, eggs, and larvae were landed at Dunedin DN in 1983.

—◎—

## Family NOCTUIDAE

### Subfamily ACRONICTINAE

(as in Nye 1975)

● ***Athetis*** Hübner, [1821], p. 209. Type species *Noctua dasychira* Hübner, by subsequent designation (Hampson 1909, p. 299, cited as “*furvula* Hübner [1808]”); Europe.

***Radinogoes*** Butler, 1886, pp. 393–394. Type species *Radinogoes tenuis* Butler, by subsequent designation (Hampson, 1909, p. 434); Queensland. Synonymised by Holloway (1977, p. 66).

***tenuis*** Butler, 1886, p. 394 (*Radinogoes*)

“Throughout mainland Australia” (Holloway 1977, p. 67). New Zealand: vagrant, sporadic, non-establishing; specimens in NZAC are from ND, AK, and TK. Each time I have collected in ND (in late November 1981 and late January 1985) I have taken *A. tenuis*. Note. Fox (1975) gives records and an illustration.

● ***Bityla*** Walker, 1865b, p. 869. Type species *Bityla thoracica* Walker, 1865, by original monotypy.

***defigurata*** Walker, 1865b, p. 756 (*Xylina*)

[Nelson NN], T.R. Oxley; HT ♀ unique, BMNH. Hudson 1898, p. 29, pl. v fig. 33; 1928, p. 76, pl. x fig. 12.

***thoracica*** Walker, 1865b, pp. 869–870 (*Bityla*). Synonymised by Meyrick (1887, p. 31).

Auckland AK, D. Bolton; LT ♂ with green-rimmed type label, BMNH.

Hudson 1898, p. 29; 1928, p. 76; as synonym.

***sericea*** Butler, 1877, p. 387 (*Bityla*)

[?MC], J.D. Enys; HT ♀ unique, BMNH. Hudson 1898, pp. 29–30, pl. v fig. 34; 1928, p. 76, pl. x fig. 11.

***pallida*** Hudson, 1905a, p. 355 (*Orthosia*). New synonymy.

Napier HB, H.W. Simmonds; HT ♂ unique, NMNZ.

Hudson 1928, p. 76, pl. x fig. 10.

● ***Cosmodes*** Guenée, 1852b, p. 289. Type species *Phalaena elegans* Donovan, by original monotypy; New South Wales.

***elegans*** Donovan, 1805, pl. 36 fig. 5, text overleaf (*Phalaena*)

New South Wales.

New Zealand: regular migrant, possibly summer-establishing.

Hudson 1898, p. 33, pl. vi fig. 2; 1928, p. 77, pl. x fig. 17.

Note. I swept 2 ♀♀ from amongst *Lobelia* growing between rushes at Mangawhai Heads ND/AK in Jan-

➤ Noctuidae, *Cosmodes elegans*

uary 1985. Walker (1858, p. 86) records 4 specimens from New Zealand.

- **Platysenta** Grote, 1874, p. 28. Type species *Platysenta atriciliata* Grote, 1874, by original monotypy; U.S.A.

**illetta** Walker, 1875, p. 684 (*Perigea*)

Northern Hindustan; India through South-east Asia to Australia and the Pacific to Rapa Island.

New Zealand: occasional immigrant.

Not mentioned by Hudson. Fox 1975, p. 372.

- **Rictonis** Nye, 1975, p. 433, objective replacement name for *Nitocris* Guenée

*Nitocris* Guenée, 1868, p. 4; preoccupied by *Nitocris* Rafinesque, 1815 (Hymenoptera). Type species *Mamestra comma* Walker, by subsequent designation (Hampson 1909, p. 383).

**comma** Walker, 1856c, p. 239 (*Mamestra*)

[?Auckland AK], J.F. Churton; HT ♀ unique, head missing, BMNH.

Hudson 1898, p. 7, pl. v fig. 27 and 28, as *Orthosia comma*; 1928, p. 76, pl. x fig. 19 and 20, as *Ariathisa comma*, after Meyrick (1912b, p. 103).

- implexa** Walker, 1857a, p. 405 (*Graphiphora*; as *Graphiphoga*). Synonymised by Meyrick (1887, p. 30).

[?Hawkes Bay HB or Taupo TO], W. Colenso; LT ♀ labelled as "Holotype ♀", "N. Zealand W. Colenso 53-19", "42. Graphiphora implexa", BMNH.

Hudson 1898, p. 7; 1928, p. 76; as synonym.

- plusiata** Walker, 1865b, p. 472 (*Hadena*). Synonymised by Meyrick (1887, p. 80).

[Nelson NN], T.R. Oxley; LT ♂ labelled as "Type", "Hadena plusiata", BMNH.

Hudson 1898, p. 7; 1928, p. 76; as synonym.

- bicomma** Guenée, 1868, pp. 4-5 (*Nitocris*), replacement name for *comma* Walker. Synonymised by Meyrick (1887, p. 30).

No type material or locality given.

Hudson 1898, p. 7; 1928, p. 76; as synonym.

- **Spodoptera** Guenée, 1852a, p. 153. Type species *Hadena mauritia* Boisduval, 1833, by subsequent designation (Hampson 1894, p. 248).

**exempta** Walker, 1857a, p. 355 (*Agrotis*)

Type locality unknown. Zimmerman (1958, p. 336) states "nearly cosmopolitan; widespread in Pacific".

New Zealand: 1 ♂, Nelson NN, 1970.

Not mentioned by Hudson.

Note. The specimen agrees with diagnostic characters given by Zimmerman (1958, pp. 333-334, fig. 274-276, 278, and 280).

**litura** Fabricius, 1775, p. 601 (*Phalaena Noctua*) India; Afghanistan through South-east Asia to Australia and eastern Pacific.

New Zealand: immigrant from Australian populations, intermittently establishing.

Not mentioned by Hudson.

Note. New Zealand specimens agree better in adult and larval colour pattern with Australian and Indian specimens than with those from east of Fiji.

**mauritia acronyctoides** Guenée, 1852, pp. 154-155 (*Spodoptera*; as species)

Tahiti; "Red Sea to India, Burma, Ceylon, Malaya to Australia" (Zimmerman 1958, p. 345), and throughout the Pacific.

New Zealand: reported once, in Wellington WN, 1896. Not recorded by Fox (1978).

Hudson 1898, p. 6, pl. v fig. 31, as *Orthosia margarita*; 1928, p. 77, pl. x fig. 21, as *Spodoptera mauritia*.

**margarita** Hawthrone, 1897, p. 283 (*Orthosia*). Synonymised by Meyrick (1912b, p. 103).

Wellington WN, E.F. Hawthrone; LT ♂ here designated, labelled "522a" ["at light in sitting room, Perceval Terrace, April 11, 1896" - Hudson's Register], NMNZ.

Hudson 1898, p. 6, pl. v fig. 31.

Note. A PLT ♂ labelled "522b" was collected "amongst papers in office, 16 April 1896" (Hudson Register entry, NMNZ).

—♂—

**Subfamily AGARISTINAE**

- **Phalaenodes** Lewin, 1805, p. 2. Type species *Phalaenodes glycinae* Lewin, by original monotypy.

**glycinae** Lewin, 1805, p. 2, pl. I (*Phalaenodes*)

New South Wales.

New Zealand: introduced by man, established ND, AK, TO.

Hudson 1946, p. 72, pl. iv fig. 3.

—♂—

**Subfamily CATOCALINAE**

(as in Kitching 1984;

includes Ophiderinae of authors)

- **Achaea** Hübner, [1823], p. 269. Type species *Phalaena melicerta* Drury, by subsequent designation (Moore 1885, p. 163); India (Bombay).

**janata** Linnaeus, 1758, p. 527 (*Phalaena Geometra*)

India; India to Australia and eastern Pacific.

New Zealand: sporadically regular immigrant, usually in late summer or autumn; larvae not persisting over winter.

Hudson 1928, p. 78, pl. x fig. 8, as *Ophiusa melicerte*.

Note. Buller (1905, p. 333) gives details of occurrence in 1903 (as *Achaea melicerte*); Philpott (1926a, p. 388) gives a synopsis of records (as *Ophiusa melicerte*).

***melicerte*** Drury, 1773, p. 46, pl. 23 fig. 1 (*Phalaena*; as *melicerte*). Synonymised in Nye (1975, p. 18).

Hudson 1928, p. 78, as species.

***traversii*** Fereday, 1877a, pp. 457–458 (*Catocala*). Synonymised by Hudson (1928, p. 78). Wellington WN, W.L. Travers; HT ♂ unique, CMNZ. Hudson 1928, p. 78, as synonym.

● ***Anomis*** Hübner, [1821], p. 249. Type species *Anomis exacta* Hübner, 1882, by original monotypy; Mexico – Peru.

***flava*** Fabricius, 1775, p. 601 (*Phalaena Noctua*) East Indies, Koenig; HT ?lost. New Zealand: occasional non-establishing immigrant (Fox 1978).

Hudson 1928, p. 83, pl. ix fig. 28.

***involuta*** Walker, 1858b, p. 1003 (*Gonitis*) Sri Lanka; HT in BMNH. New Zealand: occasional non-establishing immigrant (Fox 1978).

Hudson 1928, p. 82, pl. x fig. 27, as *Anomis sabulifera* (not of Guenée, 1852, p. 404, “Abyssinia”), after Philpott (1927a, p. 704).

● ***Arcte*** of authors, but not Kollar (1844, p. 477)

***coerulea*** Guenée, 1852c, pp. 41–42, pl. xiii fig. 10 (*Cocytodes*)

East Indies; Ceylon and India through to Samoa. New Zealand: 1 record from Auckland AK; specimen in AMNZ.

Not mentioned by Hudson.

Note. This is the type species of genus *Cocytodes*.

● ***Artigisa*** Walker, 1863, p. 160. Type species *Artigisa nigrosignata* Walker, by original monotypy; Borneo.

***melanephele*** Hampson, 1914, p. 206 (*Artigisa*) Australia, Tasmania, R.M. Green; ST series in BMNH. Hudson 1928, p. 78, pl. x fig. 18, as *Catada lignicoloria* in the sense of Philpott (1927d, p. 81) but not of Walker (1866, p. 1579), with *C. impropria* in the sense of Meyrick (1917a, p. 246) listed as a synonym.

Note. The ST series of *A. melanephele* agrees in wing pattern exactly with New Zealand specimens in NMNZ and NZAC. From *A. melanephele*, HT *Hemerophila lignicoloria* Walker (also from Tasmania) differs in having pectinate antennae, and HT *Thermesia impropria* Walker, 1865c, p. 1064, has the hindwing differently patterned ventrally, and the ♂ valva (Philpott 1927d, p. 81, fig. 2) different. *A. melanephele* is estab-

lished in the Auckland area; larvae feed on dead leaves and rotting tree stumps.

● ***Dasypodia*** Guenée, 1852c, p. 174. Type species *Dasypodia selenophora*, by subsequent designation (Meyrick 1912b, p. 105).

***cymatodes*** Guenée, 1852c, p. 175 (*Dasypodia*) Tasmania; HT ♂ in BMNH. New Zealand: immigrant, now established on (adventive) *Acacia* and *Albizia*, commoner in northern localities. Hudson 1928, p. 81; 1939, p. 401, pl. xii fig. 11. Note. Philpott (1928a, p. 359) gives early records, all from northern localities.

***selenophora*** Guenée, 1852c, p. 175 (*Dasypodia*) Tasmania; HT ♀ in MNHN. New Zealand: immigrant, now established on (adventive) *Acacia* and *Albizia*, commoner in southern localities. Hudson 1898, p. 35, pl. vi fig. 4; 1928, p. 80, pl. x fig. 13.

● ***Elygea*** Billberg, 1820, p. 85. Type species *Phalaena materna* Linnaeus, by subsequent designation (Berio 1966, as reported by Nye 1975, p. 175); India.

***materna*** Linnaeus, 1767, p. 840 (*Phalaena Noctua*) India; type material in Linnaean Society Collection, London. New Zealand: fairly regular immigrant, non-establishing. Philpott (1926a, p. 388) gives early records. Hudson 1928, p. 80, as *Ophideres materna*.

● ***Grammodes*** Guenée, 1852c, p. 275. Type species *Noctua geometrica* Fabricius, 1775, p. 599, by original designation; eastern India.

***pulcherrima*** Lucas, 1892, pp. 258–259 (*Grammodes*) Brisbane; north-eastern Australia. New Zealand: 1 record, Titahi Bay WN; specimen in NMNZ. Hudson 1928, p. 79, pl. xliv fig. 30, as *Ophiusa pulcherrima*; Hudson 1905a, p. 356, in *Grammodes*.

● ***Hypocala*** Guenée, 1852c, p. 73. Type species *Hyblaea deflorata* Fabricius, by subsequent designation (Hampson 1894, p. 452); India.

***deflorata australiae*** Butler, 1892, p. 21 (*Hypocala*) Australia; HT (?gender) in BMNH. New Zealand: sporadic immigrant, non-establishing (Fox 1978).

● ***Mocis*** Hübner, [1823], p. 267. Type species *Phalaena virbia* Cramer, 1780, by subsequent designation (Hampson 1913a, p. 76, as “*undata* Fabr.”; see Nye 1975, p. 319).

➤ Noctuidae, *Mocis*

***alterna*** Walker, 1858c, pp. 1833–1834 (*Euclidia*)

Northern Australia.

New Zealand: 1 record, Nelson NN.

Hudson 1928, p. 79, pl. x fig. 26, following Philpott (1927a, p. 704).

● ***Othreis*** Hübner, [1823], p. 264. Type species *Phalaena pomona* Cramer, 1775 (junior subjective synonym of *Phalaena fullonia* Clerck, 1764), by subsequent designation (Moore 1885, p. 131, as *fullonica* Linnaeus); India.

***fullonia*** Clerck, 1764, pl. 48 (*Phalaena*)

India – South-east Asia – Pacific.

New Zealand: occasional non-establishing immigrant; throughout. First reported by Fereday (1883a), as “*Ophideres archon* of Felder”. Hudson 1928, p. 80, as *Ophideres fullonica*.

● ***Speiredonia*** Hübner, [1823], p. 272. Type species *Phalaena feducia* Stoll in Cramer, 1790, p. 160, pl. 36 fig. 3, by subsequent designation (Desmarest 1857, p. 130; as *Spiredonia*).

***Sericia*** Guenée, 1852c, p. 172. Type species *Sericia spectans* Guenée, by subsequent designation (Desmarest 1857, p. 130), as in Holloway (1977, p. 88).

***spectans*** Guenée, 1852c, p. 173 (*Sericia*)

Australia, Tasmania; LT ♂ designated by Viette (1951, p. 162), MNHN.

New Zealand: recorded twice, in WI and NN.

Hudson 1928, p. 81, pl. li fig. 12; 1939, p. 401; as *Sericia spectans*.

● ***Tathorhynchus*** Hampson, 1894, p. 268. Type species *Scopula vinctalis* Walker (junior subjective synonym of *Spintherops exsiccata* Lederer), by original designation.

***exsiccata fallax*** Swinhoe, 1902, p. 423 (*Tathorhynchus*; as species)

Western Australia.

New Zealand: very occasional immigrant (Fox 1978).

—◎—

Subfamily CHLOEOPHORINAE

(as in Nye 1975)

● ***Pseudoips*** Hübner, 1822, p. 63. Type species *Pyralis sagana* Fabricius, 1781, p. 276, as *Pseudoips prasinana* Linnaeus in the sense of Hübner, 1822, by subsequent designation (Fletcher, 1966, p. 16, but cited as *prasinana*; see Nye 1975, p. 414).

***sagana*** Fabricius, 1781, p. 276 (*Pyralis*)

Europe.

New Zealand: 1 record in 1932, Orua River WI, in NZAC. Not mentioned by Hudson.

Note. This species – clearly man-adventive – has not been seen since 1932 despite the widespread plantings of *Quercus*, the larval host genus.

—◎—

Subfamily CUCULLIINAE

(as in Nye 1975)

● ***Andesia*** Hampson, 1906, pp. 6 (key) and 142. Type species *Andesia oenistis* Hampson, 1906, p. 142, by original designation; Argentina.

***pessota*** Meyrick, 1887, p. 29 (*Miselia*)

[Riccarton Bush], Christchurch MC, R.W. Fereday; LT ♀ here designated, labelled “Dec. 1873 Riccarton Bush”, CMNZ.

Hudson 1898, p. 6, pl. v fig. 26, as *Miselia pessota*; 1928, p. 49, pl. vi fig. 17, as *Andesia pessota*, after Meyrick (1914a, p. 102).

Note. There are no relevant specimens (currently under the genus *Sympistis* Hübner) in BMNH. At CMNZ, of the 2 relevant specimens only the ♀ is labelled as from a Christchurch locality and as caught in December; the ♂ is labelled “Feb'y/73 Dunedin”. While I could have inferred that Meyrick's citing of both sex and locality was mistaken, I saw no need, as the ♀ would have been seen by him, in Fereday's collection, in 1880.

● ***Austramathes*** Hampson, 1906, pp. xiv (key) and 492. Type species *Graphiphora purpurea* Butler, by original designation.

***purpurea*** Butler, 1879a, p. 490 (*Graphiphora*)

Dunedin DN, F.W. Hutton; HT ♂ unique, BMNH.

Hudson 1898, p. 8, pl. v fig. 32, as *Xanthia purpurea*, after Meyrick (1887, p. 81); 1928, p. 49, pl. vii fig. 3, as *Austramathes purpurea*, after Meyrick (1912b, p. 95).

***ceramodes*** Meyrick, 1887, p. 31 (*Xanthia*). Synonymised by Meyrick (1888a, p. 46).

“North Island”, J.D. Enys; LT ♂ here designated, labelled “1873-15 from Enys., North Island”, “Fereday Collection”, “31”, CMNZ.

Hudson 1898, p. 8; 1928, p. 49; as synonym.

• ***Homohadena*** Grote, 1873b, p. 180. Type species *Hadena badistriga* Grote, 1872, by monotypy; U.S.A. (New York).

***fortis*** Butler, 1880, pp. 549–550 (*Toxocampa*)

Blenheim MB, W. Skellon; HT ♂ unique, BMNH. Hudson 1928, pp. 49–50, pl. vi fig. 11.

***iota*** Hudson, 1903, p. 243 (*Miselia*). Synonymised by Meyrick (1912b, p. 95).

Karori WN, G.V. Hudson; HT ♂ unique, NMNZ. Hudson 1928, p. 49, as synonym.

—◎—

### Subfamily HADENINAE

(as in Nye 1975)

• ***Aletia*** in the sense of Meyrick (1912b, p. 97), not strictly of Hübner ([1821], p. 239). Type species *Noctua vitellina* Hübner, 1808, pl. 81 fig. 379, by subsequent designation (Moore 1881, p. 333); Europe.

***argentaria*** Howes, 1945, p. 66, pl. 7 fig. 2 (*Aletia*)  
The Wilderness SL, W.G. Howes; HT ♂ designated by Howes, NMNZ.

Hudson 1950, p. 77, pl. vii fig. 15 (a rather inaccurate representation).

***cucullina*** Guenée, 1868, p. 40 (*Xylocampa*)  
[Rakaia MC], H.G. Knaggs; HT ♂ unique, BMNH. Hudson 1898, pp. 27–28, pl. v fig. 23, as *Melanchra cucullina*; 1928, p. 58, pl. vii fig. 19, as *Aletia cucullina*, after Meyrick (1912b, p. 98).

***parmata*** Philpott, 1926a, p. 387 (*Aletia*). New synonymy.

Mount Grey NC, S. Lindsay; HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 58, pl. x fig. 25, as species.

Note. Antennal and colour pattern characters agree with those of HT *cucullina*.

***funerea*** Philpott, 1927a, pp. 703–704 (*Aletia*).  
Synonymised by Hudson (1928, p. 58).

Mount Arthur Tableland NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 58, as synonym.


***cuneata*** Philpott, 1916, p. 420 (*Aletia*)

Ben Lomond OL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, pp. 56–57, pl. ix fig. 19.

***cyanopetra*** Meyrick, 1927b, p. 313 (*Melanchra*) new combination

Waiho Gorge WD, A. Castle; LT ♂ here designated, labelled "Melanchra cyanopetra Meyr. Type ♂ Waiho Gorge 18.2.27 A. Castle", NMNZ.



### Noctuidae: Hadeninae (180) *Meterana meyricki* Hampson

Hudson 1939, p. 400, pl. lv fig. 29.

Note. The LT and the 3 other Waiho Gorge specimens in NMNZ and NZAC lack the abdominal crests, a characteristic of *Aletia* in Meyrick's sense.

***dentata*** Philpott, 1923, p. 148 (*Aletia*)

Tongariro TO, 5,000 ft, J.G. Myers; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 56, as synonym of *Aletia cuneata*.

***empyrea*** Hudson, 1918, p. 61 (*Aletia*)

Routeburn Valley OL, C.E. Clarke; LT ♀ here designated, marked with a red disc, labelled "Routeburn 25.12.14", "Aletia empyrea Huds. Type ♂" (in Clarke's handwriting), "C.E. Clarke Collection", AMNZ.

Hudson 1928, p. 59, pl. x fig. 14 and 15.

Note. Another specimen (also ♀) is labelled "Aletia empyrea Huds. Type ♀". The other 3 specimens are also ♀♀.

***falsidica falsidica*** Meyrick, 1911b, p. 70 (*Hyssia*; as species)

Mount Arthur Tableland NN, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 56, pl. vii fig. 17, as *Aletia falsidica*, after Meyrick (1912b, p. 98).

***falsidica hamiltoni*** Hampson, 1913b, p. 594 (*Hyssia*; as species)

Tararua Range WN, A. Hamilton; HT ♂ designated by Hampson, BMNH.

Hudson 1928, p. 56, as synonym.

Note. Hudson (1928, p. 56) points out that the true type locality is the Tararua Range, not Wellington as the HT specimen label has it. Hampson's subspecies is here regarded as representing the North Island populations of *Aletia falsidica*.

***fibriata*** Meyrick, 1913a, p. 22 (*Aletia*; as *fibrata*, a misprint – Meyrick 1914a, p. 101).

Mount Richmond MB, F.G. Gibbs; LT ♂ here designated, labelled "Aletia fibrata Meyr. 5/1 E. Meyrick det. in Meyrick Coll.", "Mt Richmond New Zealand F.G.G. 4500' 12.05", "Lectotype" (yellow-rimmed circular label), BMNH.

> Noctuidae, *Aletia fibrata*

Hudson 1928, p. 55, pl. vii fig. 25, as *Aletia fibrata*.

**inconstans** Butler, 1880, p. 545 (*Spaelotis*)

Blenheim MB, W. Skellon: HT ♂ with antennae missing, genitalia on "Agrotidae genitalia slide 347". BMNH. Hudson 1898, p. 9, as synonym of *Leucania griseipennis*; 1928, p. 57, pl. viii fig. 29, as *Aletia inconstans*.

**lacustris** Meyrick, 1934, p. 151 (*Aletia*)

Lake Rotoroa BR, Lawford White; LT ♂ here designated, labelled "Lake Rotoroa 14/2/31", "Aletia lacustris ver. by E. Meyrick" S. Lindsay's handwriting). CMNZ. Hudson 1939, p. 394, pl. lv fig. 27.

Note. Meyrick was sent 4 specimens by Lindsay: he retained one and sent the others back. Lindsay placed a HT label on the ♂ here regarded as a LT. The rest of the series remained at CMNZ. The specimen Meyrick retained superficially resembles a dark *A. sistens*, and is not conspecific with the rest of the series, which represents a dark, grey-green "*Aletia*" not uncommon in collections from the South Island.

**longstaffi** Howes, 1911, p. 128 (*Morrisonia*)

Ben Lomond OL, W.G. Howes and G.B. Longstaff: LT ♂ here designated, labelled "Ben Lomond 10.iii.1910", BMNH.

Hudson 1928, p. 58, pl. ix fig. 25.

**mitis** Butler, 1877, pp. 383–384 (*Agrotis*)

[Castle Hill MC], J.D. Enys: HT ♂ unique, BMNH. Hudson 1898, p. 27, as synonym of *Melanchna cucullina*; 1928, p. 55, as synonym of *Aletia moderata*; 1939, p. 394, as species, after Philpott (1928a, p. 359).

**munda** Philpott, 1917b, p. 239 (*Aletia*). New synonymy.

Waikouaiti RI/TO, H.W. Simmonds: HT ♂ designated by Philpott as "Type ♂", NMNZ.

Hudson 1928, pp. 57–58, pl. xliv fig. 17, as species.

Note. No differences are apparent in antennal structure, thoracic vestiture, or wing pattern between topotypic *mitis* and *munda*, hence the proposed synonymy.

**gourlayi** Philpott, 1921, p. 337 (*Aletia*). Synonymised by Philpott (1928g, p. 483).

Arthur's Pass NC/WD, E.S. Gourlay: HT ♂ designated by Philpott, CMNZ.

Hudson 1928, p. 57, pl. xlix fig. 31, as species; 1939, p. 394, as synonym.

**moderata** Walker, 1865a, pp. 705–706 (*Agrotis*)

[Nelson NN], T.R. Oxley: LT ♀ here designated, labelled "N. Zealand Auckland Oxley 60-73", "Agrotis? moderata", "Lectotype ♀ J.S.D. 1980", "Agrotidae genitalia slide 352", BMNH.

Hudson 1898, p. 9 as *Leucania moderata*, p. 9 and pl. iv fig. 8 as *Leucania griseipennis*; 1928, p. 55, pl. vii fig. 15, as *Aletia moderata*.

**griseipennis** Felder & Rogenhofer, 1875, pl. cix fig. 22 (*Mamestra*)

[Nelson NN, T.R. Oxley]: LT ♀ here designated, labelled

"griseipennis n.", "cix f. 22 Mamestra griseipennis n. New Zealand", "606", circular red type label, antennae missing, abdomen broken, BMNH.

Hudson 1898, p. 9, pl. iv fig. 8, as *Leucania griseipennis*; 1928, p. 55, as synonym of *Aletia moderata*.

Note. The synonymy is revised from that given by Meyrick (1912b, p. 97) because of Meyrick's confusion between *virescens* Butler (below) and *griseipennis* F. & R. Meyrick's action submerged Butler's *virescens* in what he (Meyrick) thought Felder & Rogenhofer's *griseipennis* to be. A comparision of the LTs involved in the present synonymy – both collected by Oxley around Nelson – showed their obvious conspecificity. With its conspicuously banded hindwings, *A. moderata* is quite distinct from all other New Zealand '*Aletia*' species. The reader's attention is drawn also to comments by Meyrick (1888a, p. 45).

**nobilis** Howes, 1946, pp. 144–145 (*Aletia*)

Homer FD, W.G. Howes: HT ♂ designated by Howes, NMNZ.

Hudson 1950, p. 78, pl. viii fig. 4.

**obsecrata** Meyrick, 1914a, p. 101 (*Aletia*)

Ben Lomond OL, A. Philpott: HT ♀ unique, antennae missing, thorax compressed, BMNH.

Hudson 1928, p. 58, pl. ix fig. 17 and 18.

**panda** Philpott, 1920, p. 42 (*Aletia*)

Mount Earnslaw OL, G.V. Hudson: HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 57, pl. ix fig. 9.

**probenota** Howes, 1945, p. 65, pl. 7 fig. 1 (*Aletia*)

Homer FD, W.G. Howes: HT ♂ designated by Howes, NMNZ.

Hudson 1950, pp. 77–78, pl. vii fig. 8.

**sistens** Guenée, 1868, p. 39 (*Eumichtis*)

[Rakaia MC], H.G. Knaggs: HT ♂ unique, BMNH.

Hudson 1898, p. 9, not figured, as synonym of *Leucania moderata*, after Meyrick (1887, p. 19); 1928, p. 55, as synonym of *Aletia moderata*, and p. 56, pl. vii fig. 18, as *Aletia temenaula*.

**temenaula** Meyrick, 1907c, p. 106 (*Leucania*). New synonymy.

Dunedin DN, G.V. Hudson: LT ♂ by unknown designator, labelled "Leucania temenaula Meyr. 5/2 E. Meyrick det. in Meyrick Coll.", "Dunedin New Zealand GVH '05", yellow LT label, BMNH.

Hudson 1928, p. 56, pl. vii fig. 18, as species.

**pachyscia** Meyrick, 1907c, pp. 106–107 (*Leucania*). New synonymy.

Lake Wakatipu OL, G.V. Hudson: LT ♀ designated by I.W.B. Nye, labelled "Leucania pachyscia Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", "L. Wakatipu New Zealand GVH /03", yellow LT label, BMNH.

Hudson 1928, p. 56, not figured, as species.

Note. No consistent differences in colour pattern, antennae, or ♂ external genitalia were found. Examples resem-

bling all 3 type specimens were collected on one night at Cromwell CO, hence the proposed synonymy. Revisers' attention is drawn to comments by Meyrick (1888a, p. 45).

**sollenis** Meyrick, 1914a, pp. 101–102 (*Aletia*)

Waipori DN, A. Philpott; LT ♂ by unknown designator, labelled "Aletia sollenis 1/1 E. Meyrick det. in Meyrick Coll.", "Waipori New Zealand AP 28.4.89 [28.11.09]", yellow LT label, antennae and legs missing, BMNH.

Hudson 1928, pp. 58–59, pl. x fig. 1.

Note. One ♂ labelled "Waipori 28.11.09" is in NZAC.

**hollandiae** not of Guenée, 1852, but in the sense of Meyrick (1911b, p. 69 and 1912b, p. 102) (*Dasygaster*); misidentification by Meyrick (1914a, p. 102)

**temperata** Walker, 1858c, pp. 1648–1649 (*Bryophila*)

[?Auckland AK], J.F. Churton; HT ♀ unique, Agrotidae genitalia slide no. 340, BMNH.

Hudson 1898, p. 9, as *Leucania temperata*; 1928, p. 75, pl. x fig. 28, as *Melanchra temperata*.

**inceptura** Walker, 1858c, pp. 1736–1737 (*Xylina*). Synonymised by Meyrick (1887, p. 27), and later validated (1888a, p. 45).

[?Auckland AK], J.F. Churton; HT ♂ unique, BMNH.

Hudson 1898, p. 9; 1928, p. 75; as synonym.

**deceptura** Walker, 1858c, p. 1737 (*Xylina*). Synonymised by Meyrick (1887, p. 27), and later validated (1888a, p. 45).

[?Auckland AK], J.F. Churton; HT ♀ unique, BMNH.

Hudson 1898, p. 9; 1928, p. 75; as synonym.

**accurata** Philpott, 1917b, p. 239 (*Aletia*). New synonymy.

Titahi Bay WN, M.O. Pasco; HT ♂ unique, lost, formerly in Pasco Collection, SMNZ.

Hudson 1928, p. 57, pl. xliv fig. 16, as species.

Note. Philpott's description notes that the veins are marked with white interrupted by black dots. This feature, the habitus, and the coastal type locality are characteristic of *A. temperata*, hence the proposed synonymy.

**eucrossa** Meyrick, 1927b, p. 313 (*Aletia*). New synonymy.

Waiuku AK, per J.C. Andersen; HT ♀ unique, bearing Hudson label "1101a" and "Waiuku N.Z. summer 1925–26 P. Shepherd", NMNZ.

Hudson 1938, p. 394, pl. Iv fig. 28, as species.

Note. The illustration of the unique specimen matches HT *temperata* and HT *deceptura* in colour pattern and habitus, hence the proposed synonymy.

**virescens** Butler, 1879a, p. 478 (*Chera*)

[?OL] "Otago", F.W. Hutton; HT ♀ unique, BMNH.

Hudson 1898, p. 9, as synonym of *Leucania griseipennis* (error); 1928, pp. 55–56, pl. vii fig. 16, as *Aletia grisipennis* in the sense of Meyrick (1912b, p. 34), not of Felder & Rogenhofer (1875).

Also 1 undescribed species (NZAC).

● **Dipaustica** Meyrick, 1912b, p. 98. Type species *Leucania epiastra* Meyrick, by original monotypy.

**epiastra** Meyrick, 1911b, p. 58 (*Leucania*)

Makara WN, reared by R.M. Sunley; LT ♂ here designated, labelled "Leucania epiastra Meyr. 8/3 E. Meyrick det. in Meyrick Coll.", "Makara New Zealand RMS bred .09", ST label, BMNH.

Hudson 1928, pp. 59–60, pl. x fig. 4.

Also 2 undescribed species (NZAC, and K.J. Fox Collection).

● **Feredayia** Kirkaldy, 1910, p. 8, objective replacement name for *Erana* Walker

**Erana** Walker, 1857b, pp. 495 and 605, preoccupied by *Erana* Gray, 1840 (Aves). Type species *Erana graminosa* Walker, by original monotypy.

**graminosa** Walker, 1857b, p. 605 (*Erana*)

[?Auckland AK], J.F. Churton; HT ♂ unique, BMNH. Hudson 1898, p. 28, pl. v fig. 24 and 25; 1928, pp. 62–63, pl. vii fig. 30 and 31.

**vigens** Walker, 1865b, pp. 743–744 (*Erana*). Synonymised by Meyrick (1887, p. 28).

[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.

Hudson 1898, p. 28; 1928, p. 62; as synonym.

**sphagnea** Felder & Rogenhofer, 1875, pl. cix fig. 17 (*Mamestræ*). New synonymy.

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.

Hudson 1898, p. 17; 1928, p. 65, as synonym of *Melanchra plena*.

Note. There is no doubt as to the synonymy of *sphagnea* F. & R. with Walker's *graminosa*.

● **Graphania** Hampson, 1905, pp. xiv (key) and 468 (description). Type species *Heliothobus disjungens* Walker, by original designation.

**Alysina** Cockerell, 1913, p. 15, objective replacement name for *Alysia* Guenée. Synonymised by Dugdale (1971b, p. 118).

**Alysia** Guenée, 1868, p. 3, preoccupied by *Alysia* Latreille, 1804 (Hymenoptera). Type species *Alysia specifica* Guenée, 1868, by original monotypy. *A. specifica* is a junior subjective synonym of *Agrotis nullifera* Walker (p. 204).

**Maoria** Warren, 1912, p. 76, preoccupied by *Maoria* Laporte, 1868 (Coleoptera); see Cockerell, 1913, p. 15. Type species *Erana plena* Walker, by original designation. Synonymised by Dugdale (1971b, p. 118).

Note. Nye (1975, p. 223) states that as *plena* and *disjungens* are congeneric, *Graphania* is available as a subjective replacement name.

» Noctuidae, *Graphania*

**agorastis** Meyrick, 1887, p. 18 (*Mamestraa*)

Lake Guyon MB/BR, R.W. Fereday; LT ♂ by unknown designator, labelled "Mamestra agarastis Meyr. 5/1 E. Meyrick det. in Meyrick Coll.", "Lectotype ♂", "L. Guyon New Zealand R.W.F. 3/71", yellow LT label, legs missing, BMNH.

Hudson 1898, p. 18, pl. v fig. 30; 1928, p. 73, pl. viii fig. 31, as *Melanchra agarastis*.

**averilla** Hudson, 1921, p. 255 (*Melanchra*)

Mount Egmont TK, 3,000 ft, Averil Lysaght; ST series (2 ♀♀) not located in NMNZ.

Hudson 1928, p. 68, pl. xlvi fig. 18.

**furtiva** Philpott, 1924c, pp. 663–664, fig. 1B (*Melanchra*). **New synonymy.**

Mount Arthur NN, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 66, pl. viii fig. 14, in text under *Melanchra mutans*; 1939, p. 396, pl. lv fig. 4, as species.

Note. In both nominal species ♀♀ tend to be silvery, and no clear difference in ♂ valval or antennal structures were seen, hence the proposed synonymy. Both are montane, and abundantly distinct from sympatric *mutans*. Meyrick (1929, p. 484) distinguished *furtiva* from *mutans*.

**beata** Howes, 1906, pp. 510–511 (*Melanchra*)

Otataru SL, W.G. Howes; HT ♀ unique, not located.

Hudson 1928, p. 67, pl. x fig. 2, as *Melanchra beata*; the ♂ figured is in NZAC, collected by Philpott at Kapuka SL.

Note. Large series reared on several occasions by Mr B. Patrick consistently differed in colour pattern elements from other species.

**bromias** Meyrick, 1902c, p. 273 (*Melanchra*)

Chatham Islands, J. Fougeret; LT ♂ by unknown designator, labelled "Melanchra bromias Meyr. 4/3 E. Meyrick det. in Meyrick Coll.", "bromias Meyr." (in Meyrick's handwriting), "Chatham Is. F. /00", yellow LT label, BMNH.

Hudson 1928, p. 67, pl. viii fig. 15, as *Melanchra bromias*.

**brunneosa** Fox, 1970, pp. 22–24, fig. 1–5 (*Melanchra*)

Mount Egmont TK, K.J. Fox; HT ♂ designated by Fox, NMNZ.

**chlorodonta** Hampson, 1911, pp. 423–424 (*Morrisonia*)

Ngaruawahia WO, G.B. Longstaff; HT ♀ designated by Hampson, abdomen missing, BMNH.

Hudson 1928, p. 66, pl. ix fig. 10, as *Melanchra chlorodonta*.

Note. Discovered at Ngaruawahia, north of Hamilton, not – as Hudson states (1928, p. 66) – at Cape Egmont lighthouse TK.

**chryserythra** Hampson, 1905, p. 452 (*Morrisonia*)

[?Orepuki or other locality in SL], Dunlop; HT ♂ des-

ignated by Hampson, BMNH.

Hudson 1928, pp. 75–76, pl. ix fig. 6 and 7, as *Melanchra chryserythra*.

**disjungens** Walker, 1858c, p. 1681 (*Heliothobus*) [Waikouaiti DN], P. Earl; HT ♂ unique, wings sooty along termen, BMNH.

Hudson 1898, p. 15, pl. v fig. 43, as *Melanchra disjungens*; 1928, p. 60, pl. vii fig. 23, as *Persectania disjungens*, after Meyrick (1912b, p. 98).

Note. Earl's specimen looks as though it had been damaged by heat from a paraffin lamp.

**nervata** Guenée, 1868, p. 40 (*Hadena*). **Synonymised** by Meyrick (1887, p. 15).

Rakaia MC, R.W. Fereday; LT ♂ here designated, labelled "H. nervata Gn ... [description on folded paper]", "Ex Musaeo Ach. Guenée", "Lectotype JSD 1980", BMNH. Hudson 1898, p. 15; 1928, p. 60; as synonym.

**erebia** Hudson, 1909, p. 68 (*Melanchra*)

Erebus Cove, Port Ross, Auckland Island, R. Browne; HT ♀ unique, NMNZ.

Hudson 1928, p. 67, pl. x fig. 3, as *Melanchra erebia*.

Dugdale 1971, pp. 124–126, fig. 104–112, as subspecies of *Graphania mutans*.

**oceania** Salmon, 1956, pp. 78–79 (*Melanchra*)

**Synonymised** by Dugdale (1971b, p. 124).

Ocean Island, Auckland Islands, ?J.H. Sorensen; HT ♂ designated by Salmon, NMNZ.

**fenwicki** Philpott, 1921, p. 337 (*Melanchra*)

Dunedin DN, C.C. Fenwick; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 73, pl. xlvi fig. 17.

**dives** Philpott, 1930b, p. 1 (*Melanchra*). **New synonymy.**

Flagstaff Hill DN, C.E. Clarke; HT ♂ designated by Philpott, AMNZ.

Hudson 1939, p. 395, pl. lv fig. 5 and 6, as species.

Note. Large series taken by Mr B. Patrick around Dunedin included many specimens intermediate in colour pattern between the two nominal species. No differences were seen in colour pattern or antennal structure, hence the proposed synonymy.

**homoscia** Meyrick, 1887, pp. 21–22 (*Mamestraa*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1898, p. 21, pl. v fig. 7; 1928, pp. 74–75, pl. ix fig. 23; as *Melanchra homoscia*.

**sminthistis** Hampson, 1905, p. 280, pl. 86 fig. 17 (*Hyssia*). **Synonymised** by Meyrick (1912b, p. 102).

"New Zealand", G.F. Mathew; HT ♂ designated by Hampson, BM genitalia slide no. 353, BMNH.

Hudson 1928, p. 74, as synonym.

**infensa** Walker, 1857b, p. 748 (*Orthosia*)

[Waikouaiti DN], P. Earl; HT ♀ unique, BMNH.

Hudson 1898, p. 23, pl. v fig. 12; 1928, p. 70, pl. viii fig. 19; as *Melanchra infensa*.

**arachnias** Meyrick, 1887, p. 23 (*Mamestra*).  
Synonymised by Meyrick (1888a, p. 45).  
Blenheim MB, W. Skellon; HT ♀ unique, CMNZ.  
Hudson 1898, p. 23; 1928, p. 70; as synonym.

**insignis** Walker, 1865b, p. 724 (*Eplexia*)  
[Nelson NN], T.R. Oxley; LT ♀ here designated, labelled  
“Eplexia insignis Walker”, “New Zealand Auckland  
Oxley 60-73”, green circular type label, BMNH.  
Hudson 1898, p. 16, pl. iv fig. 29 and 30; 1928, p. 65, pl.  
viii fig. 10-12 (pl. viii fig. 11 is a good match for LT  
♀); as *Melanchra insignis*.

**turbida** Walker, 1865b, pp. 754-755 (*Xylina*).  
Synonymised by Meyrick (1888a, p. 45).  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1898, p. 16; 1928, p. 65, pl. viii fig. 10 (a fair  
match for HT ♂); as synonym.

**angusta** Felder & Rogenhofer, 1875, pl. cix fig.  
18 (*Mamestra*). **New synonymy.**  
[Nelson NN, T.R. Oxley]; HT ♀ unique, abdomen missing,  
BMNH.  
Hudson 1898, p. 18; 1928, p. 66; as synonym of *Melan-*  
*chra mutans*, after Meyrick (1887, p. 17).

**skelloni** Butler, 1880, pp. 547-548 (*Hadena*).  
Synonymised by Meyrick (1912b, p. 100).  
Blenheim MB, W. Skellon; HT ♂ unique, antennae missing,  
BMNH.  
Hudson 1928, p. 65, as synonym.

**polychroa** Meyrick, 1887, pp. 16-17 (*Mames-  
tra*). **Synonymised by Meyrick (1888a, p. 45).**  
Christchurch MC, R.W. Fereday; LT ♂ designated by  
D.S. Fletcher, labelled “Melanchra insignis Walker 10/5  
E. Meyrick det. in Meyrick Coll.”, “Mamestra polychroa  
Meyr. det. D.S. Fletcher in Meyrick Coll.”, “Christchurch  
New Zealand RWF /85”, BMNH.  
Hudson 1898, p. 16; 1928, p. 65; as synonym.

**xanthogramma** Meyrick, 1912c, p. 117 (*Melan-  
chra*). **New synonymy.**  
Wellington WN, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1928, p. 65, not figured; 1939, p. 65; as species.  
Note. NZAC has many specimens with colour patterns  
intermediate between those of HT *insignis* and HT *xan-*  
*thogramma*. These HTs have the same wing shape, and  
there are no obvious differences in genitalia, hence the  
proposed synonymy. What is usually called *xan-*  
*thogramma* in collections is a broader-winged, previously  
unrecognised species in the *insignis* group.

**lignana** Walker, 1857b, p. 758 (*Hadena*)  
[Waikouaiti DN], P. Earl; HT ♂ unique, BMNH.  
Hudson 1898, p. 26, pl. v fig. 19; 1928, p. 71, pl. viii fig.  
21; as *Melanchra lignana*.

**lindsayi** Dugdale (*Graphania*) new species. Replacement name for *Melanchra olivea* not of Watt (1916, p. 413), but in the sense of Hudson (1928, p. 67 (part), pl. xliv fig. 32♀; 1939, p. 396).  
Dunedin DN, W.G. Howes; HT ♀ labelled “954g” (Hudson's label), “Dunedin”, NZAC.  
Hudson 1928, p. 65, pl. xliv fig. 32; 1939, p. 396.

**Note.** This commonly collected species is sympatric with *olivea* Watt, but differs from it as detailed by Lindsay in Hudson (1939, p. 396). Lindsay's concept of Philpott's *lata* is an excellent description of Watt's *olivea*. As the species with jagged subterminal line and oblique, curved reniform has been adequately characterised and illustrated, but is separated as either *olivea* or *lata* in collections, it needs a name. I propose to name it after Mr Stuart Lindsay, who first appreciated its distinctiveness.

**lithias** Meyrick, 1887, p. 17 (*Mamestra*)  
Castle Hill MC, J.D. Enys; ?ST ♂ labelled “Fereday Col-  
lection”, CMNZ.  
Hudson 1898, pp. 17-18, pl. iv fig. 33; 1928, p. 74, pl. ix  
fig. 24; as *Melanchra lithias*.  
Note. Hudson (1898) reported that the 2 STs were in the  
Fereday collection.

**maya** Hudson, 1898, p. 17 (*Melanchra*)  
Tableland Track, Mount Arthur NN, 3,400 ft, G.V. Hud-  
son; HT ♀ unique, NMNZ.  
Hudson 1898, p. 17, pl. iv fig. 31; 1928, p. 64, pl. viii fig.  
8, as *Melanchra maya*.

**mollis** Howes, 1907, p. 533 (*Melanchra*; as *molis*)  
Dunedin DN, W.G. Howes; LT ♂ here designated,  
labelled “coll. G. Howes. Museum Collection”,  
NMNZ.

Hudson 1928, p. 68, pl. iv fig. 27, as *Melanchra mollis*.  
Note. In Philpott's copy of *Trans. N.Z. Inst.* vol. 40, p.  
533, is the marginal note “Laps cal.” beside Howes's  
“*molis*”. Howes (1912, p. 204) twice referred to the  
species as *mollis*, and gave a new illustration.

**morosa** Butler, 1880, p. 543 (*Xylophasia*)  
[Blenheim MB], W. Skellon; HT ♂ labelled as “Type”,  
antennae missing, BMNH.

Hudson 1898, pp. 19-20, pl. v fig. 3 and 4, as *Melanchra*  
*pelistis*, and p. 26, as synonym of *M. lignana*, after  
Meyrick (1887, p. 26); 1928, p. 74, pl. viii fig. 22, as  
*Melanchra morosa*.

**pelistis** Meyrick, 1887, p. 20 (*Mamestra*). **Syn-**  
**onymised by Meyrick (1912b, p. 101).**  
Akaroa MC, R.W. Fereday; LT ♂ by unknown designator,  
labelled “*Melanchra morosa* Meyr. 5/1 E. Meyrick  
det. in Meyrick Coll.”, “Akaroa New Zealand R.W.F. /82”,  
“*pelistis* Meyr. lectotype ♂”, BMNH.  
Hudson 1898, pp. 19-20, pl. v fig. 3 and 4, as species;  
1928, p. 74, as synonym.

**mutans** Walker, 1857b, p. 602 (*Hadena*)  
[Auckland AK], D. Bolton; LT ♂ here designated, labelled  
“100. *Hadena mutans*”, “N. Zealand Bolton 54-4”,  
green circular type label, BMNH.  
Hudson 1898, p. 18, pl. iv fig. 13-16; 1928, p. 66, pl. viii  
fig. 13 and 14; as *Melanchra mutans*.

**lignifusca** Walker, 1857b, p. 603 (*Hadena*). **Syn-**  
**onymised by Meyrick (1887, p. 17).**  
[Auckland AK], D. Bolton; LT ♂ here designated, labelled  
“101 *Hadena lignifusca*”, “New Zealand Bolton 54-4”,  
green type label, BMNH.

» Noctuidae, *Graphania mutans*

---

Hudson 1898, p. 18; 1928, p. 66; as synonym.

**spurcata** Walker, 1857b, p. 631 (*Xylina*). Synonymised by Meyrick (1912b, p. 100).

[Auckland AK], D. Bolton; LT ♀ here designated, labelled "21. *Xylina spurcata*", "New Zealand Bolton 54-4", green type label, BMNH.

Hudson 1928, p. 66, as synonym.

**vexata** Walker, 1865b, p. 755 (*Xylina*). Synonymised by Meyrick (1887, p. 17).

[Nelson NN], T.R. Oxley; HT ♂ unique, genitalia missing, BMNH.

Hudson 1898, p. 18; 1928, p. 66; as synonym.

**acceptrix** Felder & Rogenhofer, 1875, pl. cix fig. 19 (*Mamestra*). Synonymised by Meyrick (1887, p. 17).

[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.

Hudson 1898, p. 18; 1928, p. 66; as synonym.

**passa** Morrison, 1874b, p. 139 (*Mamestra*). Synonymised by Franclemont (1981, p. 133).

"California" U.S.A. (error), H.K. Morrison; HT ♀ unique, MSUZ.

Not mentioned by Hudson.

**debilis** Butler, 1877, p. 385 (*Hadena*). Synonymised by Meyrick (1887, p. 17).

[Hawkes Bay HB], J.D. Enys; HT ♀ unique, BMNH.

Hudson 1898, p. 18; 1928, p. 66; as synonym.

**pallescens** Warren, 1912, p. 77 (*Maoria*; as aberration of *mutans*). **New synonymy.**

Dunedin DN, A. Purdie; HT ♀ designated by Warren, BMNH.

Not mentioned by Hudson.

**nullifera** Walker, 1857b, p. 742 (*Agrotis*)

[Waikouaiti DN], P. Earl; HT ♂ unique, BMNH.

Hudson 1898, pp. 9–10, pl. iv fig. 9, as *Leucania nullifera*, after Meyrick (1887, p. 7); 1928, p. 55, pl. vi fig. 14, as *Aletia nullifera*, after Meyrick (1912b, p. 97).

**specifica** Guenée, 1868, p. 3 (*Alysia*). Synonymised by Meyrick (1887, p. 7).

"Canterbury" MC, R.W. Fereday; HT ♂ designated by Guenée, BMNH.

Hudson 1898, p. 9; 1928, p. 55; as synonym.

**olivea** Watt, 1916, p. 413 (*Melanchra*)

Mount Egmont TK, M.N. Watt; HT ♂ not located; 9 ♀ 9 ♂ PTs same locality, Jan 1916, NMNZ.

Hudson 1928, p. 67, pl. ix fig. 30 and 31, as *Melanchra olivea*.

**lata** Philpott, 1927d, p. 81 (*Melanchra*). **New synonymy.**

Arthur's Pass [village] NC/WD, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1939, p. 396, not figured, as species.

Note. Hudson (1939, p. 396, in part), refers to *olivea* Watt from Egmont, not the Howes specimens referred to by Watt (1916, p. 413, footnote) as "the southern form of

*M. olivea*". Watt's type material and Philpott's are indistinguishable. See also *lindsayi*, above.

**oliveri** Hampson, 1911, p. 424 (*Morrisonia*)

Bold Peak OL, F.S. Oliver; HT ♀ designated by Hampson, Agrotidae genitalia slide no. 298, BMNH.

Hudson 1928, p. 69, pl. viii fig. 18, as *Melanchra oliveri*.

**omicron** Hudson, 1898, p. 22, pl. v fig. 42 (*Melanchra*)

[Karori] WN, A. Norris; ST series with Hudson labels "143a", "143b", "143c", "143d" not located.

Hudson 1898, p. 22, pl. v fig. 42, as species; 1928, p. 57, as synonym of *Aletia inconstans*; 1939, p. 400, as species.

**omoplaca** Meyrick, 1887, p. 24 (*Mamestra*)

Rakaia MC, R.W. Fereday; LT ♂ selected by Hampson (1905, p. 382), BMNH.

Hudson 1898, pp. 23–24, pl. v fig. 13; 1928, p. 70, pl. viii fig. 26 and 27, as *Melanchra omoplaca*.

**umbra** Hudson, 1903, p. 243 (*Melanchra*). Synonymised by Meyrick (1912b, p. 101).

Invercargill SL, W.G. Howes; LT ♂ here designated, labelled "527a" ["Invercargill, Howes and Philpott" – Hudson Register], NMNZ.

Hudson 1928, p. 70, as synonym.

**micropis** Hampson, 1918a, p. 121 (*Morrisonia*). **New synonymy.**

[?Wellington WN], G.V. Hudson; HT ♀ designated by Hampson, Agrotidae genitalia slide no. 330, BMNH.

Not mentioned by Meyrick or Hudson.

**pagaia** Hudson, 1909, p. 67, pl. II fig. 9 (*Leucania*)

The Snares islands, W. Benham; HT ♂ unique, NMNZ.

Hudson 1928, p. 52, pl. ix fig. 20, as *Leucania pagai*. Dugdale 1971, pp. 123–124, fig. 97–103, as subspecies of *Graphania insignis*.

Note. The status of various entities in the *insignis* complex is uncertain; while *pagaia* is a veritable *Graphania*, its relationship to *scutata*, *fenwicki*, and *insignis* requires further study.

**paracausta** Meyrick, 1887, pp. 15–16 (*Mamestra*)

Castle Hill MC, J.D. Enys; LT ♂ here designated, labelled "Mamestra paracausta Meyr." (in Meyrick's handwriting, "Fereday Collection", "72", CMNZ).

Hudson 1898, p. 15, pl. iv fig. 28 and 28A; 1928, p. 68, pl. viii fig. 23 and 24; as *Melanchra paracausta*.

**pelanodes** Meyrick, 1931a, pp. 92–93 (*Melanchra*) **new combination**

Waimarino [National Park] TO, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1939, p. 400, pl. iv fig. 21, as *Melanchra pelanodes*.

**petrograpta** Meyrick, 1929, p. 484 (*Melanchra*)

Kinloch, head of Lake Wakatipu OL, G.V. Hudson; HT ♀ unique, Agrotidae genitalia slide no. 360, BMNH.

Hudson 1928, pl. ix fig. 21, as *Melanchra mutans*, variety;

1939, p. 396, pl. lv fig. 14; 1950, pp. 81–82, pl. vi fig. 2; as *Melanchra petrograpta*.

***phricias*** Meyrick, 1888a, p. 46 (*Mamestra*)

Christchurch MC, R.W. Fereday; LT ♂ by unknown designator, labelled "Mamestra phricias Meyr. 5/1 E. Meyrick det. in Meyrick Coll.", "Christchurch New Zealand RWF 7/2/65", yellow LT label, BMNH. Hudson 1898, p. 27, pl. v fig. 22; 1928, p. 75, pl. ix fig. 11; as *Melanchra phricias*.

Note. Meyrick proposed *phricias* as a name for *temperata* Meyrick, 1887, p. 27 (not of Walker, 1858, p. 1648). A ♀ ex Felder Collection in BMNH is labelled as "Spiti", "Cucullia cellulata Warr. type ♀ [symbol inverted] Warr.", and "Spiti – err. loc. Hadeninae = Morrisonia phricias Meyr. fr. N. Zealand". I cannot trace publication of Warren's epithet.

***plena*** Walker, 1865b, p. 744 (*Erana*)

[Nelson NN], T.R. Oxley; HT ♂ unique, photographed (BM negative no. 47546), Noctuidae slide no. 82330 ♂, BMNH.

Hudson 1898, p. 17, pl. iv fig. 32; 1928, p. 65, pl. viii fig. 3 and 4; as *Melanchra plena*.

***viridis*** Butler, 1880, p. 547 (*Dianthroecia*). Synonymised by Meyrick (1887, p. 17).

Blenheim MB, W. Skellon; HT ♀ designated by Butler, photographed (BM negative no. 47547), BMNH. Hudson 1898, p. 17; 1928, p. 65; as synonym.

***prionistis*** Meyrick, 1887, p. 27 (*Mamestra*)

Rakaia MC, R.W. Fereday; LT ♂ by unknown designator, labelled "Mamestra prionistis Meyr. 3/1 E. Meyrick det. in Meyrick Coll.", "Rakaia New Zealand RWF 2/76", BMNH.

Hudson 1898, p. 27, pl. v fig. 21; 1928, p. 75, pl. ix fig. 22; as *Melanchra prionistis*.

***rubescens*** Butler, 1879a, p. 489 (*Xylophasia*)

"Otago", F.W. Hutton; HT ♂ labelled as "Xylophasia rubescens Butler (Type)", BMNH.

Hudson 1898, pp. 25–26, pl. v fig. 18; 1928, p. 71, pl. ix fig. 4 and 5; as *Melanchra rubescens*. Dugdale 1971, pp. 119 and 129–130, fig. 118–120, as *Graphania rubescens*.

***scutata*** Meyrick, 1929, p. 485 (*Melanchra*)

Wellington WN, G.V. Hudson; HT ♂ unique, BMNH. Hudson 1928, pl. viii fig. 12, as *Melanchra insignis*, ♀; 1939, p. 395, pl. iv fig. 12, as *Melanchra scutata*.

***sericata*** Howes, 1945, pp. 66–67 (*Melanchra*)

Homer Tunnel FD, W.G. Howes; HT ♂ designated by Howes, NMNZ.

Hudson 1950, p. 80, pl. vii fig. 12.

***sequens*** Howes, 1912, pp. 204–205 (*Morrisonia*)

Whakarewarewa BP, G.B. Longstaff; HT ♂ designated by Howes, no longer so labelled; series of possible STs, BMNH.

Hudson 1928, p. 75, pl. ix fig. 26, as *Melanchra sequens*.

***distracta*** Meyrick, 1924a, p. 202 (*Melanchra*).

**New synonymy.**

Mount Ruapehu TO, 4,000 ft, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 76, as species.

***tetrachroa*** Meyrick, 1931a, p. 92 (*Melanchra*)

Waimarino [National Park] TO, G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1939, pp. 399–400, pl. iv fig. 20, as *Melanchra tetrachroa*.

***ustistriga*** Walker, 1857b, pp. 630–631 (*Xylina*)

[?Hawkes Bay HB or Taupo TO], W. Colenso; LT ♀ here designated, labelled "N. Zealand Colenso 53-19", "19. Xylina ustistriga", "Lectotype JSD 1980", BMNH.

Hudson 1898, pp. 26–27, pl. v fig. 20 and 20A; 1928, p. 68, pl. viii fig. 16 and 17; as *Melanchra ustistriga*.

***lignisecta*** Walker, 1857b, p. 631 (*Xylina*). Synonymised by Meyrick (1887, p. 26).

Auckland AK, D. Bolton; HT ♀ unique, labelled in error as LT by Dugdale, BMNH.

Hudson 1898, p. 26; 1928, p. 68, as synonym.

Note. Of two candidate specimens under *ustistriga* in BMNH, both from Auckland and collected by Bolton, only the ♀ can be Walker's unique holotype of *lignisecta*. G.F. Hampson had chosen an Oxley specimen collected in 1860, after Walker's description was published.

Also 4 undescribed species (NZAC).

• ***Ichneutica*** Meyrick, 1887, p. 13. Type species *Ichneutica ceraunias* Meyrick, by original monotypy.

***cana*** Howes, 1914, p. 96 (*Ichneutica*)

Hector Mountains OL, W.G. Howes; HT ♂ unique, not found in NMNZ.

Hudson 1928, pp. 50–51, pl. ix fig. 15.

***lata*** Philpott, 1915, p. 192 (?*Aletia*). Synonymised by Salmon (1945, pp. 1–3).

Vanguard Peak OL, H. Hamilton and F.S. Oliver; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 51, pl. vi fig. 24, as species; 1950, p. 75, as synonym of *Ichneutica cana*.

Note. This synonymy was arrived at without examination of the type material.

***ceraunias*** Meyrick, 1887, p. 13 (*Ichneutica*)

Mount Arthur NN, 4,700 ft, E. Meyrick; HT ♂ unique, BMNH.

Hudson 1898, pp. 14–15, pl. iv fig. 25 and 26; 1928, p. 50, pl. vi fig. 4–6.

***dione*** Hudson, 1898, p. 14, pl. iv fig. 27 (*Ichneutica*)

Mount Arthur NN, 4,400 ft, C.W. Palmer; HT ♂ unique, NMNZ.

Hudson 1928, p. 50, pl. vi fig. 22; 1950, p. 73, pl. v fig. 9.

» Noctuidae, *Ichneutica dione*

Note. Hudson (1950, p. 73) presents a clear view of his opinion of type specimens.

**homERICA** Howes, 1943, pp. 371–372 (*Ichneutica*)  
Homer Tunnel FD, W.G. Howes; HT ♂ designated by  
Howes, NMNZ.  
Hudson 1950, p. 74, pl. v fig. 10 and 11.

**lindsayi** Philpott, 1926a, p. 387 (*Ichneutica*)  
Hunter Mountains FD, 4,000 ft, S. Lindsay; HT ♂ des-  
ignated by Philpott, CMNZ.  
Hudson 1928, p. 50, pl. vi fig. 8.

**marmorata** Hudson, 1924, pp. 7-8 (*Persectania*)  
Arthur's Pass NC/WD, J.W. Campbell; HT ♀ unique,  
CMNZ.  
Hudson 1928, p. 51, pl. 1 fig. 10 and 17, as *Ichneutia*  
*marmorata*.

**dives** Philpott, 1924a, p. 207 (*Ichneutica*). Syn-  
onymised by Philpott (1927d, p. 80).  
Mount Arthur NN, A. Philpott; HT ♂ designated by  
Philpott, NZAC.  
Hudson 1928, p. 51, as synonym; ♂ figured pl. 1 fig. 10.

**nervosa** Hudson, 1922, p. 196 (*Ichneutica*)  
Bold Peak OL, F.S. Oliver; ST series not located.  
Hudson 1928, p. 51, pl. 1 fig. 1.

**notata** Salmon, 1946, pp. 1–3, pl. 1 fig. 2 and 3  
(*Ichneutica*)  
Mount Peel NN, H. Hamilton; HT ♀ designated by  
Salmon, NMNZ.  
Hudson 1950, pp. 75–76, pl. viii fig. 5; also 1928, pl. vi  
fig. 23 (as *Ichneutica lata* ♂).

Also 1 undescribed species (NZAC).

● **Meterana** Butler, 1877, pp. 385–386. Type spe-  
cies *Dianthezia pictula* not of White (1855), but in  
the sense of Butler (1877, pp. 385–386) [= *Miselia*  
*meyricci* Hampson, 1911], by original designation.  
Note. This genus is resurrected for those New  
Zealand "*Melanchra*" species excluded from *Graphania*  
by Dugdale (1971b, pp. 118–119).

**alcyone** Hudson, 1898, p. 24, pl. v fig. 14 (*Melan-  
chra*) new combination  
Botanic Gardens, Wellington WN, A. Norris; ST series  
not located.  
Hudson 1928, p. 70, pl. viii fig. 25, as *Melanchra alcyone*.  
Dugdale 1971, p. 119, in "Erana" group.

**asterope** Hudson, 1898, pp. 24–25, pl. v fig. 15  
(*Melanchra*) new combination  
Mt Arthur Tableland NN, G. V. Hudson; HT ♀ unique,  
labelled "76a", NMNZ.  
Hudson 1928, p. 72, pl. viii fig. 30, as *Melanchra*  
*asterope*. Dugdale 1971, p. 119, in "Erana" group.

**badia** Philpott, 1927d, p. 80 (*Melanchra*) new  
combination

Leslie Valley, Mount Arthur NN, A. Philpott; HT ♂ des-  
ignated by Philpott, NZAC.

Hudson 1939, p. 398, pl. lv fig. 11, as *Melanchra badia*.  
**meridiana** Salmon, 1956, p. 573, pl. 21 fig. 1 and  
2 (*Melanchra*). New synonymy.

Portobello DN, W.G. Howes; HT ♂ designated by  
Salmon, NMNZ.

Note. Salmon (1956, pl. 21 fig. 3) illustrates *Graphania*  
*morosa*. The ♀ "M. coeleno" with genitalia on BMNH  
Agrotidae slide no. 300 is a ♀ *M. badia*.

**coeleno** Hudson, 1898, p. 26, pl. iv fig. 39 (*Melan-  
chra*) new combination

Wellington WN, G.V. Hudson; candidate NT ♂, labelled  
"456a", "on roof of observatory, Oct. '97", NMNZ.

Hudson 1928, p. 69, pl. viii fig. 20, as *Melanchra coeleno*.  
Dugdale 1971, p. 119, in "Erana" group.

Note. The original specimens collected by Norris and  
Hawthorne could not be located. Specimen "456a" was  
probably collected after Hudson had sent his MS. to  
London to be typeset, as he mentioned no specimens  
other than those captured by Norris and Hawthorne.

**coctilis** Meyrick, 1931a, p. 93 (*Melanchra*) new  
combination

Flora River, Mount Arthur NN, G.V. Hudson (reared);  
LT ♂ here designated, labelled "Melanchra coctilis  
Meyr. 1/1 E. Meyrick det. in Meyrick Coll.", "Flora  
R. New Zealand GVH bred .30", "Lectotype", BMNH.  
Hudson 1939, p. 399, pl. lv fig. 19.

**decorata** Philpott, 1905, pp. 328–329 (*Melanchra*)  
new combination

?West Plains SL, A. Philpott; HT ♂ designated by Phil-  
pott, no other label, NZAC.

Hudson 1928, p. 70, pl. viii fig. 5, as *Melanchra decorata*.

**diatmeta** Hudson, 1898, p. 21, pl. v fig. 5 (*Melan-  
chra*) new combination

Wellington WN, A. Norris; HT ["described from a spec-  
imen coll. Norris" – Hudson Register, NMNZ] not  
located.

Hudson 1928, pp. 69–70, pl. viii fig. 7, as *Melanchra*  
*diatmeta*.

**dotata** Walker, 1857b, pp. 522–523 (*Dasypolia*) new  
combination

[?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♂  
unique, Agrotidae genitalia slide no. 323, BMNH.

Hudson 1898, p. 24, pl. v fig. 16; 1928, p. 72, pl. viii fig.  
34; as *Melanchra dotata*. Dugdale 1971, p. 119, in  
"Erana" group.

**boldensis** Salmon, 1956, pp. 573–574, pl. 21  
fig. 4 and 5 (*Melanchra*). New synonymy.

Bold Peak OL, J.T. Salmon; HT ♀ designated by Salmon,  
NMNZ.

Note. The characteristic wing pattern and elongate wing  
shape of *dotata* are present in designated *boldensis*, and  
specimens in NZAC from localities between central North

Island and Bold Peak are uniform in appearance, hence the proposed synonymy.

***exquisita*** Philpott, 1903, pp. 246–247 (*Melanchra*)  
new combination

West Plains SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, pp. 63–64, pl. viii fig. 1, as *Melanchra exquisita*. Dugdale 1971, p. 119, in “*Erana*” group.

***grandiosa*** Philpott, 1903, p. 246 (*Melanchra*) new combination

West Plains SL, A. Philpott; HT ♀ designated by Philpott, NZAC.

Hudson 1928, p. 64, pl. viii fig. 9, as *Melanchra grandiosa*. Dugdale 1971, p. 119, in “*Erana*” group.

***inchoata*** Philpott, 1920, p. 43 (*Melanchra*) new combination

Stephens Island SD, H. Hamilton; HT ♂ designated by Philpott, NMNZ.

Hudson 1928, p. 69, pl. xlvi fig. 31 and 32, as *Melanchra inchoata*.

***levis*** Philpott, 1905, p. 329, pl. xx fig. 4 (*Melanchra*) new combination

West Plains SL, A. Philpott; HT ♂ designated by Philpott, NZAC.

Hudson 1928, p. 74, pl. viii fig. 28, pl. xlvi fig. 18, as *Melanchra levis*. Dugdale 1971, p. 119, in “*Erana*” group.

***merope*** Hudson, 1898, p. 19, pl. v fig. 2 (*Melanchra*) new combination

Wellington WN, G.V. Hudson; HT ♂ unique, labelled “268a”, NMNZ.

Hudson 1928, p. 72, pl. x fig. 24, as *Melanchra merope*. Dugdale 1971, p. 119, in “*Erana*” group.

***chlorographa*** Hampson, 1905, p. 452 (*Morrisonia*). Synonymised by Longstaff (1912, p. 111, footnote).

Orepuki SL, Dunlop; HT ♀ designated by Hampson, Agrotidae genitalia slide no. 343, BMNH.

Hudson 1928, p. 72, as synonym.

***meyricci*** Hampson, 1911, p. 421 (*Miselia*; as new name)

Macetown OL, H. Hamilton; HT ♂ labelled as “Type ♂” by Hampson, BMNH.

Hudson 1898, p. 19, pl. iv fig. 87; 1928, p. 63, pl. vii fig. 33; as *Melanchra pictula*.

***pictula*** not of White (in Taylor 1855), but in the sense of Butler (1877, p. 386, pl. xlvi fig. 1) (*Diantheocia*). Synonymised by Hampson (1911, p. 421).

***pictula*** White in Taylor, 1855, p. xiii (list), pl. 1 fig. 3 (*Diantheocia*)

[North Island], R. Taylor; HT not located.

Hudson 1898, pl. iv fig. 38; 1928, p. 63, pl. vii fig. 32; as *Melanchra rhodopleura*.

Note. White's illustration of *pictula* clearly represents the North Island population, lacking the prominent white reniform characteristic of South Island *meyricci*, as pointed out by Philpott (1928g, p. 483). Sunley (1911, p. 129) described the larva (as *rhodopleura*).

***rhodopleura*** Meyrick, 1887, p. 19 (*Mamestra*). New synonymy.

Wellington WN, R.W. Fereday; LT ♂ here designated, labelled “Wellington New Zealand R.W.F. /85”, “*Melanchra rhodopleura* Meyr. 2/1 E. Meyrick det. in Meyrick Coll.”, BMNH.

Hudson 1898, p. 19; 1928, p. 63; as species.

***ochthistis*** Meyrick, 1887, pp. 20–21 (*Mamestra*) new combination

Christchurch MC, R.W. Fereday; LT ♂ here designated, labelled “*Melanchra ochthistis* Meyr. 5/1 E. Meyrick det. in Meyrick Coll.”, “Christchurch New Zealand RWF /85”, BMNH.

Hudson 1898, p. 20, pl. iv fig. 42, as *Melanchra vitiosa* and as synonym; 1928, p. 73, pl. ix fig. 14, as *Melanchra ochthistis*. Dugdale 1971, p. 119, in “*Erana*” group.

***octans*** Hudson, 1898, p. 25, pl. v fig. 1 (*Melanchra*) new combination

Mount Linton SL, A. Philpott; HT ♀ unique, labelled “378a”, NMNZ.

Hudson 1928, p. 64, pl. viii fig. 2, as *Melanchra octans*. Dugdale 1971, p. 119, in “*Erana*” group.

***pansicolor*** Howes, 1912, p. 204 (*Morrisonia*) new combination

Vauxhall, Dunedin DN, W.G. Howes; LT ♂ here designated, labelled “Vauxhall 14.11.09”, “Fenwick Coll.” (the 2 PLTs are ♀♀), NMNZ.

Hudson 1928, p. 69, pl. ix fig. 8, as *Melanchra pansicolor*. Dugdale 1971, p. 119, in “*Erana*” group and wrongly ascribed to Meyrick.

***pascoi*** Howes, 1912, pp. 205–206 (*Morrisonia*) new combination

Queenstown OL, M.O. Pasco; 9♂ 4♀ STs each labelled “Cotype”, AMNZ.

Hudson 1928, p. 71, pl. ix fig. 1–3, as *Melanchra pascoi*. Dugdale 1971, p. 119, in “*Erana*” group.

***captiosa*** Philpott, 1927d, pp. 80–81 (*Melanchra*). New synonymy.

Mount Arthur Tableland NN, 4,000 ft, A. Philpott; HT ♀ designated by Philpott, NZAC.

Hudson 1939, p. 400, pl. lv fig. 3, as species.

***saeva*** Meyrick, 1929, pp. 484–485 (*Melanchra*). New synonymy.

Arthur's Pass [village] NC/WD, G.V. Hudson; LT ♀ by unknown designator, labelled “*Melanchra saeva* Meyr. E. Meyrick det. in Meyrick Coll.”, “Arthurs Pass New Zealand GVH bred 1.28”, “Lectotype”, BMNH.

Hudson 1939, pp. 397–398, pl. lv fig. 17 and 18, as species.

Note. Comparison of the type material and of material in NZAC from intermediate localities showed no differences, hence the proposed synonymies.

**pauca** Philpott, 1910, p. 544 (*Melanchra*) new combination

"Wairarapa" WA, ?collector; HT ♀ designated by Philpott, not located in NMNZ; PT ♂ labelled by Philpott, NZAC.

Hudson 1928, p. 64, pl. viii fig. 6, as *Melanchra pauca*. Dugdale 1971, p. 119, in "Erana" group.

**praesignis** Howes, 1911, p. 128, pl. 1 fig. 4 (*Morrisonia*) new combination

Orepuki SL, W.G. Howes; HT ♂ labelled as "Type" by Howes, not located.

Hudson 1928, p. 66, pl. ix fig. 9, pl. xviii fig. 6, as *Melanchra praesignis*. Dugdale 1971, p. 119, in "Erana" group.

**ludibunda** Philpott, 1929a, p. 300 (*Melanchra*).

New synonymy.

Mount Arthur NN, A. Philpott; HT ♀ designated by Philpott, NZAC.

Hudson 1939, p. 398, pl. iv fig. 9, as species.

**stipata** Walker, 1865b, pp. 753–754 (*Xylina*) new combination

[Nelson NN], T.R. Oxley; LT ♀ here designated, labelled "Xylina stipata", "N. Zealand Auckland Oxley 60-73", "Type" (circular, black-rimmed label), BMNH.

Hudson 1898, p. 25, pl. v fig. 17; 1928, pp. 71–72, pl. viii fig. 35 and 36; as *Melanchra stipata*. Dugdale 1971, p. 119, in "Erana" group.

**tartarea** Butler, 1877, p. 384 (*Graphiphora*) new combination

[MC], J.D. Enys; HT ♂ labelled by Butler as "Type", Agrotidae genitalia slide no. 326, BMNH.

Hudson 1898, p. 21, pl. v fig. 6; 1928, pp. 72–73, pl. viii fig. 32 and 33; as *Melanchra tartarea*. Dugdale 1971, p. 119, in "Erana" group.

**vitiosa** Butler, 1877, p. 384, pl. xlivi fig. 3 (*Apamea*) new combination

[MC], J.D. Enys; HT ♂ labelled by Butler as "Type", BMNH.

Hudson 1898, p. 20, pl. iv fig. 40, as *Melanchra proteastis*; 1928, p. 73, pl. ix fig. 13, as *Melanchra vitiosa*. Dugdale 1971, p. 119, in "Erana" group.

**proteastis** Meyrick, 1888a, p. 45 (*Mamestra*); replacement name for *M. vitiosa* not of Butler (1877) but in the sense of Meyrick (1887, p. 20). Synonymised by Meyrick (1912b, p. 101).

Christchurch MC, R.W. Fereday; LT ♂ by unknown designator, labelled "Melanchra vitiosa Butler, 3/3 E. Meyrick det. in Meyrick Coll.", "Mamestra proteastis Meyr., E. Meyrick det. in Meyrick Coll.", "Christchurch New Zealand RWF 6/69", LT label, BMNH.

Hudson 1898, p. 20, as species; 1928, p. 73, as synonym.

Also 1 undescribed species (NZAC, BMNH).

• **Mythimna** in the sense of Kloet & Hincks (1972, p. 77)

**loreymima** Rungs, 1953, pp. 139 and 141, fig. 2C (*Leucania*, subgenus *Acantholeucania*)

Vietnam, Saigon, J. de Joannis; HT ♂ designated by Rungs, MNHN.

Papuaasia.

New Zealand: variably established in ND, AK, WO, TK, WI, WN (Fox 1978), and present on the Kermadec Islands (Meyrick 1912, p. 97).

Hudson 1938, p. 393, pl. lxii fig. 19, as *Aletia loreyi* (not of Duponchel).

**separata** Walker, 1865, pp. 626–627 (*Leucania*)

China, Shanghai; HT ♂ unique, BMNH.

Old World: widely distributed.

New Zealand: resident, ND–WN, NN, MB.

Hudson 1898, pp. 13–14, pl. iv fig. 24, as *Leucania unipuncta* in the sense of Meyrick (1887, p. 12; as *Leucania extranea* [form A] Guenée, 1852, p. 78), not of Haworth; 1928, pp. 55–56, pl. vii fig. 14, as *Aletia unipuncta* in the sense of Meyrick (1912, p. 97).

Note. The larva is an agricultural pest known as the northern armyworm. Common (1965a, pp. 14–17) gives a useful guide to species in Australia, one of which has been intercepted here, in cargo.

• **Persectania** Hampson, 1905, pp. xii (key) and 386.

Type species *Noctua ewingii* Westwood, 1839, by original designation (as *Persectania ewingi*, an unjustified emendation – Nye 1965, p. 379); Australia.

**aversa** Walker, 1856c, p. 113 (*Leucania*)

"Country unknown"; HT ♀ unique, BMNH.

Hudson 1898, pp. 22–23, pl. v fig. 8 and 9, as *Mamestra composita*, after Meyrick (1887, p. 22), not of Guenée, 1852; 1928, p. 61, pl. vii fig. 27, as *Persectania aversa*, different from Australian *P. composita* (a synonym of *P. ewingii* Westwood).

**peracuta** Morrison, 1874a, p. 114 (*Morrisonia*).

Formally synonymised by Franclemont (1981, pp. 135–136, fig. 3 and 4).

?California (locality unknown); LT ♀ designated and illustrated by J.G. Franclemont, USNM.

Hudson 1928, p. 61, as synonym of *composita*, after Meyrick (1912, p. 99). Common 1954a, p. 88, as probable synonym of *aversa*.

Note. The ♂ labelled "M. peracuta Morr. Type" from the Grote Collection, no. 1738, Agrotidae genitalia slide no. 148, BMNH, was never indicated in a publication. Franclemont's action has precedence.

**maori** Felder & Rogenhofer, 1875, pl. cix fig. 24 (*Mamestra*). Synonymised by Meyrick (1887, p. 22).

[Nelson NN, T.R. Oxley]; HT ♀ unique, Agrotidae genitalia slide no. 261, BMNH.

Hudson 1898, p. 22; 1928, p. 61; as synonym.

**dentigera** Butler, 1880, p. 542 (*Leucania*). Synonymised by Meyrick (1887, p. 22).

Blenheim MB, W. Skellon; HT ♀ unique, Agrotidae genitalia slide no. 151, BMNH.  
Hudson 1898, p. 22; 1928, p. 61; as synonym.

● **Physetica** Meyrick, 1887, p. 5. Type species *Spaelotis caerulea* Guenée, by original monotypy.

**caerulea** Guenée, 1868, p. 38 (*Spaelotis*)  
[Rakaia MC], R.W. Fereday; HT ♂ unique, BMNH.  
Hudson 1898, p. 8, pl. iv fig. 7; 1928, p. 59, pl. vii fig. 20–22.

Note. Specimens from OL and FD in NZAC have the wings grey ventrally, not shining buff.

**hudsoni** Howes, 1906, p. 510, pl. xliv fig. 1 (*Physetica*). Synonymised by Hudson (1928, p. 59).  
Tuturau SL, R. Gibb; HT ♂ unique, not located.  
Hudson 1928, p. 59 as synonym.

● **Tmetolophota** Hampson, 1905, pp. xiv (key) and 470. Type species *Leucania propria* Walker, 1856, by original designation.

Note. Dugdale (1971b, p. 120) transferred to this genus all New Zealand species placed by Hudson (1928, 1939) in *Leucania* and *Persectania* except *P.aversa* and *P. disjungens*.

**acontistis** Meyrick, 1887, p. 9 (*Leucania*)  
Castle Hill MC, J.D. Enys; HT ♂ unique, CMNZ.  
Hudson 1898, p. 11, pl. iv fig. 14; 1928, p. 52, pl. vii fig. 5; as *Leucania acontistis*.

Note. *T. acontistis*, *T. stulta* (Philpott), and *T. paraxysta* (Meyrick) form an intergrading group on colour pattern.

**alopa** Meyrick, 1887, p. 10 (*Leucania*)  
Lake Guyon MB/BR, R.W. Fereday; LT ♂ here designated, labelled "Leucania alopa Meyr." in Meyrick's handwriting, "Feb-Mar /75 Lake Guyon", CMNZ.  
Hudson 1898, p. 12, pl. iv fig. 165; 1928, p. 53, pl. vii fig. 8; as *Leucania alopa*.

**atristriga** Walker, 1865b, p. 756 (*Xylina*)  
[Nelson NN], T.R. Oxley; LT ♂ here designated, labelled "Xylina atristriga", "New Zealand Auckland Oxley 60-73", "Type" (circular label, green margin). BMNH.  
Hudson 1898, p. 10, pl. iv fig. 12, as *Leucania astristriga*; 1928, pp. 61–62, pl. vii fig. 28, as *Persectania atristigma*.

**antipoda** Felder & Rogenhofer, 1875, pl. cix fig. 23 (*Mamestra*). Synonymised by Meyrick (1887, p. 8).  
[Nelson NN, T.R. Oxley]; HT ♀ unique, BMNH.  
Hudson 1898, p. 10; 1928, p. 61; as synonym.

**arotis** Meyrick, 1887, p. 11, line 20 (*Leucania*)  
Blenheim MB, W. Skellon; LT ♂ here designated, labelled "From Skellon", "Fereday Collection", "Lectotype ♂ Leucania arotis Meyr.", CMNZ.  
Hudson 1898, p. 12, pl. vi fig. 18, as *Leucania arotis*; 1928, p. 61, pl. vii fig. 26, as *Persectania arotis*.

**aulacias** Meyrick, 1887, p. 11, line 1 (*Leucania*).  
Synonymised by Hudson (1898, p. 12).

Dunedin DN, [R.W. Fereday]; HT ♂ unique. CMNZ.  
Hudson 1898, p. 12 and footnote; 1928, p. 61; as synonym.  
Note. Hudson's synonymy is the action of a "first reviser" as defined by Article 24(i) of the International Code of Zoological Nomenclature, 1964, p. 25, and is mandatorily here upheld, as Recommendation 24A (which would reverse the synonymy) is not a Rule, i.e., is not mandatory.

**obsoleta** Howes, 1906, p. 511 (*Leucania*), preoccupied. Synonymised by Longstaff (1912, p. 112).  
Dunedin DN, W.G. Howes; HT ♀ not located.  
Hudson 1928, p. 61, as synonym.

Note. *Leucania obsoleta* Howes is a junior homonym of *L. obsoleta* Hübner 1803 (Europe) in Hampson (1905, p. 599), now *Mythimna obsoleta* in Heath et al. (1979, p. 269). A search of BMNH, CMNZ, and NMNZ failed to reveal any ♀ collected in December 1904 or any earlier year. Howes's description was read to the Otago Institute on 12 September 1905.

**innotata** Howes, 1908, p. 534, (*Leucania*), replacement name for *obsoleta* Howes. Synonymised by Longstaff (1912, p. 112).

Anderson's Bay DN, W.G. Howes; LT ♀ here designated, labelled "Dunedin 2.10.06", "Museum Collection Coll. G. Howes", NMNZ.

Hudson 1928, p. 61, as synonym.  
Note. As well as altering the name *obsoleta* to *innotata* Howes gave a new description and cited particular specimens collected around Dunedin in October 1906. I have identified these as being in NMNZ, and have designated a LT from that series.

**blenheimensis** Fereday, 1883c, p. 196 (*Leucania*)  
Blenheim MB, W. Skellon; LT ♀ here designated, labelled "10", "Fereday Collection", CMNZ.

Hudson 1898, p. 13, pl. iv fig. 23; 1928, p. 53, pl. vii fig. 9; as *Leucania blenheimensis*.

**hartii** Howes, 1914, p. 95 (*Leucania*)  
Cape Egmont TK, S. Hart; HT ♀ unique, not located.  
Hudson 1928, p. 54, pl. ix fig. 16, as *Leucania harti*.

**lissoxyla** Meyrick, 1911b, p. 70 (*Leucania*)  
Mount Arthur Tableland NN, 4,000 ft, G.V. Hudson; HT ♂ unique, BMNH.

Hudson 1928, p. 53, pl. vii fig. 7, as *Leucania lissoxyla*.

**micrastra** Meyrick, 1897, p. 383 (*Aletia*)  
Wellington WN, G.V. Hudson; HT ♀ unique, BMNH.  
Hudson 1898, p. 12, pl. iv fig. 10, as *Leucania micrastra*; 1928, p. 54, pl. vii fig. 13, as *Aletia micrastra*, after Meyrick (1912b, p. 97).

**sapiens** Meyrick, 1929, p. 484 (*Aletia*). New synonymy.

Waipoua TO, G.V. Hudson; HT ♂ unique, BMNH.  
Hudson 1939, p. 393, pl. lv fig. 10, as *Aletia sapiens*.  
Note. No feature could be found distinguishing between the two HTs, and all North Island material in NZAC agrees outwardly with both, hence the proposed synonymy.

➤ Noctuidae, *Tmetolophota*

**paraxysta** Meyrick, 1929, p. 483 (*Leucania*)  
 Waiaouru TO, G.V. Hudson; LT ♂ selected by D.S. Fletcher and here published, labelled "Leucania paraxysta Meyr. 4/1 E. Meyrick det. in Meyrick Coll.", "Waiaouru New Zealand G.V.H. 12.28", BMNH.  
 Hudson 1939, p. 393, pl. iv fig. 25 and 26, as *Leucania paraxysta*.

Note. See note under *T. acontistis*, above.

**phaula** Meyrick, 1887, p. 10 (*Leucania*)  
 Christchurch MC, [R.W. Fereday]; LT ♂ here designated, labelled "Nov. 1872 Sumner, bred fr. larva found in tussock", "Fereday Collection", CMNZ.

Hudson 1898, p. 11, pl. iv fig. 15; 1928, p. 53, pl. vii fig. 6; as *Leucania phaula*.

**dunedinensis** Hampson, 1905, p. 591 (*Leucania*). Synonymised by Meyrick (1912b, p. 96). Dunedin DN, A. Purdie; HT ♂ designed by Hampson, BMNH.

Hudson 1928, p. 53, as synonym.

Note. Hampson's HT also bears the label "Spaelotis dunedinensis Butler, type".

**neurae** Philpott, 1905, p. 330, pl. xx fig. 5 (*Leucania*). Synonymised by Meyrick (1909a, p. 5). West Plains SL, A. Philpott; 2 ST ♂♂, both labelled "West Plains" [no date], NZAC.  
 Hudson 1928, p. 53, as synonym.

**propria** Walker, 1856c, pp. 111–112 (*Leucania*)  
 [Waikouaiti DN], P. Earl; HT ♂ unique, BMNH.

Hudson 1898, p. 11, pl. iv fig. 13, as *Leucania propria*; 1928, p. 62, pl. viii fig. 29, as *Persectania propria*, after Meyrick (1912b, p. 99).

**basifascia** Hampson, 1913b, p. 598 (*Persectania*). New synonymy.

[OL or CO], A. Hamilton; HT ♂ designated by Hampson, Agrotidae genitalia slide no. 171, BMNH.  
 Hudson 1928, p. 62, not figured, as species.

**purdii** Fereday, 1883c, pp. 195–196 (*Leucania*)  
 Fairfield, Dunedin DN, A. Purdie; HT ♂ unique, CMNZ.  
 Hudson 1898, p. 10, pl. iv fig. 11; 1928, p. 52, pl. vi fig. 13; as *Leucania purdii*.

**semivittata** Walker, 1865a, p. 628 (*Leucania*)  
 [Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
 Hudson 1898, p. 13, pl. iv fig. 21 and 22; 1928, p. 53, pl. vii fig. 10; as *Leucania semivittata*.

**similis** Philpott, 1924a, pp. 207–208 (*Persectania*)  
 Gouland Downs NN, A. Philpott; HT ♂ designated by Philpott, NZAC.  
 Hudson 1928, p. 61; 1939, p. 394, pl. iv fig. 22; as *Persectania similis*.

**steropastis** Meyrick, 1887, pp. 22–23 (*Mamestra*)  
 Christchurch MC, R.W. Fereday; LT ♂ by unknown designator, labelled "Mamestra steropastis Meyr. 2/1 E.

Meyrick det. in Meyrick Coll.", "New Zealand RWF .85", BMNH.

Hudson 1898, p. 23, pl. v fig. 10 and 11, as *Melanchra steropastis*; 1928, p. 60, pl. vii fig. 24, as *Persectania steropastis*.

Note. A Dunedin DN specimen from Purdie (accession no. 88-8) in BMNH bears a Butler MS. name (*aberrans*) which is not listed in the Museum's Noctuidae card catalogue.

**stulta** Philpott, 1905, p. 329, pl. xx fig. 1 (*Leucania*)  
 West Plains SL, A. Philpott; HT ♀ so labelled by Philpott, NZAC.

Hudson 1928, p. 54, pl. vii fig. 12, as *Leucania stulta*.

Note. See note under *T. acontistis*, above.

**sulcana** Fereday, 1883, pp. 267–268, pl. ix fig. 3  
 (?*Leucania*)

Rowe's Bush, Akaroa MC, R.W. Fereday; LT ♀ here designated, labelled "Type", "5.2.78 Akaroa Rowes Bush at night", CMNZ.

Hudson 1898, p. 13, pl. iv fig. 19 and 20; 1928, pp. 53–54, pl. vii fig. 11; as *Leucania sulcana*.

**toroneura** Meyrick, 1901, p. 565 (*Leucania*)

Mount Cook MK, G.V. Hudson; LT ♂ by unknown selector, labelled "Leucania toroneura Meyr. 6/4 lectotype E. Meyrick det. in Meyrick Coll.", "Mt Cook New Zealand G.V.H. 12/99", BMNH.

Hudson 1928, pp. 52–53, pl. vi fig. 19, as *Leucania toroneura*.

**unica** Walker, 1856c, p. 12 (*Leucania*)  
 [Waikouaiti DN], P. Earl; HT ♀ unique, BMNH.

Hudson 1898, p. 12, pl. iv fig. 17; 1928, p. 52, pl. vi fig. 2; as *Leucania unica*.

**juncicolor** Guenée, 1868, p. 2 (*Nonagria*). Synonymised by Meyrick (1887, p. 10).  
 [?Rakaia MC], H.G. Knaggs; HT ♂ unique, BMNH.

Hudson 1898, p. 12; 1928, p. 52; as synonym.

Also 3 undescribed species (NZAC and K.J. Fox Collection).



#### Subfamily HELIOTHINAE

(as in Hardwick 1970, Bretherton et al. 1983)

● **Heliothis** Ochsenheimer, 1816, p. 91. Type species *Phalaena dipsacea* Linnaeus (= *Phalaena viriplaca* Hufnagel), by subsequent designation (Samouelle, 1819, p. 252); Europe.

Note. The gender of *Heliothis* has now been ruled on (ICZN 1985b, p. 158). The feminine ending for specific names is retained here, following an ICRISAT recommendation (Nye 1982, pp. 7–8).

**Subgenus *Helicoverpa*** Hardwick, 1965, p. 9. Type species *Noctua armigera* Hübner, 1808, pl. 79 fig. 370, by original designation.

Note. Hardwick (1970, pp. 7–19) reviewed heliothine genera and reiterated two clear characters separating *Helicoverpa* from *Heliothis*: (a) presence of cornuti on the aedeagal vesica; and (b) presence of a group of stiff, modified scales on the fore femur, both lacking in *Heliothis* in the sense of Hardwick (1965) and *Schinia* in the sense of Hardwick (1970). Scott (1980, p. 1) synonymised *Helicoverpa* with *Heliothis* on the grounds that if the characters delimiting *Helicoverpa* from *Heliothis* and *Schinia* are given a different ‘weighting’, Hardwick’s premise does not hold. Nye (1982, p. 5) proposed subgeneric status for *Helicoverpa*, a course followed here. In redescribing *H. armigera*, Bretherton *et al.* (1983, p. 296) place it without comment in *Heliothis*, and do not list *Helicoverpa* (of which *armigera* is the type species) in the synonymy under *Heliothis*.

***armigera conferta*** Walker, 1857b, p. 690 (*Heliothis*; as species)

Auckland AK, A. Sinclair; ST series in BMNH.  
Hudson 1898, p. 32, pl. v fig. 40 and 41; 1928, pp. 46–47, pl. vi fig. 26; as *Heliothis armigera*.

***punctigera*** Wallengren, 1860, p. 171 (*Heliothis*)  
Australia, Sydney; HT ♂ in NRSS.  
New Zealand: sporadic immigrant, occasionally establishing for a summer, but overwintering doubtful (Fox 1978).

Not mentioned by Hudson.

—♀—

#### Subfamily HYPENINAE

(as in Nye 1975, but excluding *Schrankia*)

Richards (1933) and Owada (1987) both define Herminiinae on the tympanal hood being pre-spiracular (in front of abdominal spiracle 1). Although *Rhapsa* and New Zealand *Trigonistis* accord well with most characters described for Japanese Herminiinae by Owada, in our genera the tympanal hood is post-spiracular, i.e., the spiracle is on the inner surface of the hood, facing anteriorly (Figures 114 and 115). As South Pacific representatives of *Bocana* Walker, *Hydrillodes* Guenée, *Lophocoleus* Robinson, and *Simplicia* Guenée show a range of spiracle 1 positions – from just outside the hood rim to on the rim to under the rim – the stated apomorphy is unsatisfactory, and may describe only one end of a grade. Therefore, I am reluctant at present to assign the New Zealand representatives to Herminiinae.

● ***Lithilaria*** Rosenstock, 1885, p. 425. Type species *Lithilaria ossicolor* Rosenstock, by original monotypy; Australia.

***Hyperaucha*** Meyrick, 1897b, p. 383. Type species *Hyperaucha octias* Meyrick, by original monotypy; New Zealand and Australia. Synonymised by Nye (1975, p. 250).

Note. In 1981 the New Zealand material was under *Trigonistis* Meyrick, in BMNH. Meyrick (1901, p. 566), specifically excluded from his concept of *octias* all New Zealand material, and made it plain that the name referred only to his Australian specimens. The situation requires clearing up.

● ***Rhapsa*** Walker, 1866a, p. 1149. Type species *Rhapsa scotosialis* Walker, by original monotypy.

***scotosialis*** Walker, 1866a, p. 1150 (*Rhapsa*) [Nelson NN], T.R. Oxley; HT ♂ unique, BMNH. Hudson 1898, p. 36, pl. vi fig. 5 and 6; 1928, p. 82, pl. x fig. 6 and 7.

***lilacina*** Butler, 1877, p. 388, pl. xlii fig. 11 (*Herminia*). Synonymised by Meyrick (1887, p. 88). [MC], J.D. Enys; HT ♂ unique, BMNH. Hudson 1898, p. 36; 1928, p. 82; as synonym.

● ***Trigonistis*** Meyrick, 1902a, p. 39. Type species *Trigonistis demonias* Meyrick, by original monotypy.

***anticlina*** Meyrick, 1901, p. 566 (*Hypenodes*) new combination

Wellington WN, G.V. Hudson; 2 ST ♀ ♀, one labelled “Wellington New Zealand G.V.H. /93”, the other “New Zealand GVH /99”, BMNH.

Hudson, 1898, p. 37 pl. vi fig. 7, as *Rhapsa octias* (not of Meyrick, 1897, p. 383); 1928, p. 78, pl. x fig. 16, as *Hypenodes anticlina*.

Note. The ST series of *anticlina* is under *Trigonistis* in BMNH.

Also 1 undescribed species (NZAC).

—♀—

#### Subfamily HYPENODINAE

(as in Forbes 1954)

● ***Schrankia*** Hübner, [1825], p. 345. Type species *Pyralis taenalis* Hübner, by monotypy; Europe.

***costaestrigalis*** Stephens, 1834, p. 45 (*Hypenodes*) Cosmopolitan.  
New Zealand: ND–SL, throughout.  
Hudson 1898, p. 34, as *Hypenodes exsularis*; 1928, p. 77, pl. x fig. 9, as *Hypenodes costaestrigalis*.

➤ Noctuidae, *Schrankia costaestrigalis*

***exsularis*** Meyrick, 1888a, p. 46 (*Hypenodes*).  
Synonymised by Meyrick (1910b, p. 69).  
[New Plymouth] TK, E. Meyrick; HT ♂ unique, BMNH.  
Hudson 1898, p. 34, as species; 1928, p. 77, as synonym.

***triangulalis*** Hudson, 1923a, p. 64 (*Scoparia*).  
Synonymised by Hudson (1928, p. 77).  
Kao ND, C.E. Clarke; HT not located.  
Hudson 1928, p. 77, as synonym.

—⊗—

**Subfamily NOCTUINAE**  
(as in Nye 1975)

● ***Agrotis*** Ochsenheimer, 1816, p. 66. Type species *Noctua segetum* Denis & Schiffermüller, 1775, p. 81, by subsequent designation (Curtis 1827, p. 165); Austria.

***infusa*** Boisduval, 1838, p. 240 (*Noctua*)  
“Australie”, ?collector; HT not located (Common 1958a, p. 76).  
New Zealand: regular spring migrant (Fox 1978). Philpott (1920, p. 42) first recorded adults here.  
Hudson 1928, p. 48, pl. xlvi fig. 35, as *Agrotis spina* Guénée.  
Note. Common (1954b, pp. 223–263; 1958, pp. 69–88, pl. 1) gives comprehensive biological and taxonomic information.

***innominata*** Hudson, 1898, p. 31, pl. v fig. 39  
(*Agrotis*)  
Wellington WN, J.H. Lewis and W.R. Morris; 2 ST ♂♂, not located, NMNZ.  
Hudson 1928, p. 48, pl. vi fig. 20.  
Note. The entry in Hudson's Register at NMNZ reads “442a Newtown Wellington. J.H. Lewis Dec. 92”, but the specimen is gone.

***ipsilon aneitura*** Walker, 1865a, p. 701 (*Agrotis*; as species)  
Vanuatu, Aneityum; HT ♂ unique, Agrotidae genitalia slide no. 562, BMNH.  
Hudson 1898, p. 30, pl. v fig. 35 and 36; 1928, p. 48, pl. vi fig. 21; as *Agrotis ypsilon*.  
Note. Common (1958a, p. 73) lists differences in colour pattern between the nominate subspecies (*ipsilon*) from Europe and subspecies *aneitura* from Australia, New Zealand, and Papua New Guinea. He could discern no genital differences.

***munda*** Walker, 1857a, p. 348 (*Agrotis*)  
Australia, Adelaide, Wilson; LT ♂ labelled as “type”, Agrotidae genitalia slide no. 546, BMNH.  
New Zealand: sporadic, non-establishing (Fox 1978).

Hudson 1928, p. 47, pl. vi fig. 25, as *Euxoa radians*.  
Note. One of Walker's series was sent from New Zealand by Rev. J.F. Churton, of Auckland.

● ***Diarsia*** in the sense of Holloway (1977, p. 62)

***intermixta*** Guenée, 1852a, p. 337 (*Noctua*)  
“sans indication de patrie”; HT ♀ unique, MNHN.  
Hudson 1898, p. 7, pl. v fig. 29, as *Orthosia immunis*; 1928, pp. 48–49, pl. vii fig. 1 and 2, as *Graphiphora comptata*.

***compta*** Walker, 1857a, p. 404 (*Graphiphora*).  
Synonymised by implication (Holloway 1977, p. 62).  
Australia, New South Wales, Wood; HT ♂ unique, BMNH.  
Hudson 1928, pp. 48–49, pl. vii fig. 1 and 2.

***immunis*** Walker, 1857a, p. 430 (*Taeniocampa*).  
Synonymised with *compta* by Meyrick (1912b, p. 95).  
[?Hawkes Bay HB or Taupo TO], W. Colenso; HT ♀ unique, BMNH.  
Hudson 1898, p. 7, as species; 1928, p. 48, as synonym.

***communicata*** Walker, 1865b, pp. 716–717 (*Orthosia*).  
Synonymised with *compta* by Meyrick (1912b, p. 95).  
[Nelson NN], T.R. Oxley; HT ♂ unique, BMNH.  
Hudson 1928, p. 48 as synonym.

***acetina*** Felder & Rogenhofer, 1875, pl. cix fig. 6 (*Agrotis*).  
Synonymised with *immunis* by Meyrick (1887, p. 30).  
[Nelson NN, T.R. Oxley]; HT ♂ unique, BMNH.  
Hudson 1898, p. 7; 1928, p. 48; as synonym.  
Note. There are 8 other synonyms of *D. intermixta*, 6 by Walker, 2 by Warren; none refer to New Zealand.

● ***Ectopatria*** Hampson, 1903a, pp. xx (key) and 652.  
Type species *Agrotis subrufescens* Walker, 1865; Australia.

***aspera*** Walker, 1857b, p. 601 (*Hadena*)  
Australia, Argent [45–48]; HT ♂ unique, BMNH.  
Hudson 1939, p. 392, pl. iv fig. 30.  
Note. Coastal, from Three Kings Islands ND to NN.

***provida*** Walker, 1858c, pp. 1737–1738 (*Xylina*).  
Synonymised by Hampson (1903, p. 654).  
Auckland AK, D. Bolton; HT ♂ unique, BMNH.  
Hudson 1939, p. 392, as synonym.

***canescens*** Walker, 1865b, p. 757 (*Xylina*).  
Synonymised by Hampson (1903, p. 654).  
[Nelson NN], T.R. Oxley; LT ♂ labelled as “Type”, BMNH.  
Hudson 1939, p. 392, as synonym.  
Note. Philpott (1928g, p. 482) erroneously gave the locality of *canescens* as Auckland.

● ***Euxoa*** in the sense of Meyrick (1912b, p. 94)

**admirationis** Guenée, 1868, p. 38 (*Agrotis*) [?MC], H.G. Knaggs; HT ♂ unique, BMNH. Hudson 1898, p. 31, pl. v fig. 37, as *Agrotis admirationis*; 1928, p. 47, pl. vi fig. 7, as *Euxoa admirationis*, after Meyrick (1912b, p. 94).

**sericea** Butler, 1879a, p. 490 (*Chersotis*). Synonymised by Hampson (1903, p. 173).

Wairarapa WA, J. Lorimer; HT ♂ labelled as "type", BMNH.

Hudson 1898, p. 31, pl. v fig. 38, as species; 1928, p. 47, as synonym.

**inconspicua** Butler, 1880, p. 545 (*Chersotis*). Synonymised by Hudson (1898, p. 31, with *sericea*); and by Hampson (1903, p. 173, with *admirationis*).

Blenheim MB, W. Skellon; HT ♀ labelled as "Type", BMNH.

Hudson 1898, p. 31; 1928, p. 47; as synonym.

**veda** Howes, 1906, p. 511 (*Agrotis*). Synonymised by Meyrick (1912b, p. 94).

Motueka NN, W.G. Howes; HT ♂ unique, not located. Hudson 1928, p. 47, as synonym.

**obscura** Salmon, 1946, p. 3 (*Euxoa*; as var. of *admirationis*). **New synonymy.**

Arthur riverbed, Milford Sound FD, J.T. Salmon; HT ♂ designated by Salmon, NMNZ.

Note. Dark specimens from GB to SL are present in NZAC; the shape of the orbicular marking is variable in both pale and dark specimens.

**cerapachoides** Guenée, 1868, p. 39 (*Agrotis*) [MC], R.W. Fereday; HT ♂ unique, Noctuidae genitalia slide no. 9528 ♂, BMNH.

Hudson 1898, p. 32, pl. vi fig. 1, as *Agrotis cerapachoides*; 1928, p. 47, pl. vii fig. 4, as *Euxoa cerapachoides*, after Hampson (1903, p. 174).




## Subfamily NOLINAE

● **Celama** Walker, 1865, p. 500. Type species *Celama liparisalis* Walker, by original monotypy.

**parvitis** Howes, 1917, p. 274 (*Adeixis*)

Harbour Cone, Broad Bay DN, C.E. Clarke; HT ♂ unique, labelled as "Type" by Clarke, AMNZ.

Hudson 1928, p. 44, pl. xliv fig. 7, as *Celama parvitis*, after Philpott (1927a, p. 703).



---

Noctuidae: Nolinae  
(181) *Celama parvitis* Howe

---



## Subfamily PLUSIINAE (as in Kitching 1987)

● **Chrysodeixis** Hübner, [1821b], p. 252. Type species *Phalaena chalcites* Esper, by subsequent designation (Dyar 1902, p. 81; cited as *Crysodeixia* and *chalcitis*); Italy.

**argentifera** Guenée, 1852b, pp. 352–353 (*Plusia*) "Nouvelle Hollande". New Zealand: a regular, intermittently establishing immigrant (Fox 1978). Not mentioned by Hudson.

**eriosoma** Doubleday, 1843, p. 285 (*Plusia*). Auckland AK, A. Sinclair; HT ♂ unique, BMNH. Hudson 1898, p. 35, pl. vi fig. 3; 1928, p. 79, pl. x fig. 5; as *Plusia chalcites*.

Note. There are consistent differences in pheromone complement (Benn *et al.* 1982) and morphology between Mediterranean and South-east Asian populations. Kitching (1987, p. 144) notes that there is some evidence of morphological overlap in intermediate areas.

● **Ctenoplusia** Dufay, 1970, p. 91. Type species *Plusia limbirena* Guenée, by original designation.

**albostriata** Bremer & Grey, 1853, p. 18 (*Plusia*) China.

New Zealand: regular immigrant, probably established and overwintering in northern areas (Fox 1978). Hudson 1928, p. 80, pl. x fig. 22 and 23, as *Plusia oxygramma* not of Geyer, but in the sense of Meyrick (1912a, p. 105; wrongly ascribed to Hübner).

Note. Meyrick (1909a, p. 5) recorded this species as *Plusia transfixa* Walker. Kitching (1987, p. 145) discusses its generic position and host plants.

• ***Thysanoplusia*** Ichinose, 1973, p. 137. Type species *Phytometra intermixta* Warren, by original designation; China, Japan.

***orichalcea*** Fabricius, 1775, p. 607 (*Noctua*), as in Kostrowicki (1961, p. 395, fig. 44 and 113)

“India, Koenig”.

New Zealand: immigrant in large numbers, AK, Feb–May 1984; breeding AK-WO, from 1985 (NZAC).

Not mentioned by Hudson.

—♂—

#### SPECIES OF UNCERTAIN FAMILY POSITION

• ***Lysiphragma*** of authors, but not Meyrick (1888e)

***argentaria*** Salmon, 1948, p. 310, pl. 60 fig. 2  
(*Lysiphragma*)

Great Island, Three Kings, E.G. Turbott; HT ♂ unique, AMNZ.

Not mentioned by Hudson.

Note. This species is excluded from Tineidae and Tineoidea on characters of venation (R5 to termen, pterostigma on R2), head scaling (broad, proclinate scales), presence of ocelli, antennal scaling (dorsal only), haustellum (long, locked), labial palpi (lacking bristles), and maxillary palpi (reduced, porrect). The pleural area on abdominal sternite 8 is broad but not lobate. The habitus resembles that of the *Prays* group.

• ***Titanomis*** Meyrick, 1888e, p. 104. Type species *Titanomis sisyrota* Meyrick, by original monotypy.

***sisyrota*** Meyrick, 1888e, p. 104 (*Titanomis*)  
[Nelson NN]. G.V. Hudson; HT ♀ unique, BMNH.

Hudson 1928, p. 350, pl. xxv fig. 28, in Tineidae.

Note. Hudson (1928, p. 350) stated that the HT was caught in Nelson in 1882. He recorded no specimens from Wellington WN0 Meyrick's label reads “G.V. Hudson Wellington 10.5.85”, and there is a Hudson label “27a”. *T. sisyrota* has the following characters: a well developed, naked haustellum; 4-segmented, opposed, short maxillary palpi; no ocelli; a densely scaled metanotum; vein *M* in the discal cell forked, tubular in hindwing, vestigial in forewing; forewing discal cell with a chorda; and no tympanic organ on base of abdomen. This combination excludes it from Nocuoidea, but may not exclude it from Coccoidea. Dr P.W. Schoorl (pers. comm.) recently examined the only known male of this species, and could not include it in Coccoidea.

—♂—

## REFERENCES

- Ackery, P.R.; Vane-Wright, R.I. 1984: Milkweed butterflies; their cladistics and biology, being an account of the natural history of the Danainae, a subfamily of the Lepidoptera, Nymphalidae. London, Dept. of Entomology, British Museum (Natural History). 425 p.
- Agassiz, D.J.L. 1982: *Oegoconia caradjai* Popescu-Gorj & Capuse (Lep: Gelechiidae) recognised as British. *Proceedings and transactions of the British Entomological and Natural History Society* 15 : 1-5.
- Alfsken, J.D. 1899: *Pyrameis ida* sp.n., ein neuer Tagfalter nebst einer Varietät von den Chatham Inseln. *Zoologischer Anzeiger*, Leipzig, 22 : 5-8.
- 1903: Beitrag zur Insectenfauna der Hawaiischen und Neuseelandischen Inseln (Ergebnisse einer Reise nach dem Pacific), Schauinsland 1896-97. *Zoologische Jahrbücher*, Jena, 29 : 561-628.
- Andrews, J.R.H. 1986: The southern ark: zoological discovery in New Zealand 1769-1900. Auckland, Century Hutchinson. 237 p.
- Bates, H.W. 1867: New species of insects from the Province of Canterbury, New Zealand, collected by R.W. Fereday, Esq. *Entomologists' monthly magazine* 4 : 52-56.
- Benn, M.H.; Galbreath, R.A.; Young, H.; Down, G.; Preisner, E. 1982: The sex pheromone of the silver Y moth *Chrysodeixis eriosoma* (Doubleday) in New Zealand. *Zeitschrift für Naturforschung (C)* 37 (11-12) : 1130-1135.
- Berg, C. 1899: Substicucion de nombres genericos. III. *Communicaciones del Museo Nacional de Buenos Aires*, I : 77-79.
- Billberg, G.J. 1820: *Enumeratio Insectorum in Museo G.J. Billberg*. 138 p.
- Blanchard, E. 1852: Insectos: Ordern VI Lepidopteros. Pp. 1-112 in Gay, Historia física y política de Chile (zoología) 7.
- Bleszynski, S. 1955: Studies on the Crambidae (Lepidoptera). Part IX. *Polski pismo entomologiczne* 25 : 227-231.
- 1962: Studies on the Crambidae (Lepidoptera), Part XXXVII. Changes in the nomenclature of some Crambidae, with the descriptions of new genera and species. *Polski pismo entomologiczne* 32 : 5-48.
- Bleszynski, S.; Collins, R.J. 1962: A short catalogue of the world species of the family Crambidae (Lepidoptera). *Acta zoologica Cracoviensis* 7 (12) : 197-389.
- Boisduval, J.B.A.D. de 1832: Faune entomologique de l'Océan Pacifique, avec l'illustration des insectes nouveaux recueillis pendant le voyage. Pt 1. Lépidoptères. Pp. 5-267 in d'Urville, Voyage de découvertes de l'Astrolabe. Paris, Tastu. IV.
- Bradley, J.D. 1956a: A new genus for *Tortrix postvittana* (Walker) and certain other Australian and New Zealand species (Lepidoptera: Tortricidae). *Bulletin of entomological research*, London 47 : 101-105.
- 1956b: Records and descriptions of Microlepidoptera from Lord Howe Island and Norfolk Island collected by the British Museum (Natural History) Rennell Island Expedition, 1953. *Bulletin of the British Museum (Natural History), entomology* 4 (4) : 143-164.
- 1961: Microlepidoptera from the Solomon Islands. Additional records and descriptions of Microlepidoptera collected in the Solomon Islands by the Rennell Island Expedition 1953-54. *Bulletin of the British Museum (Natural History), entomology* 10 (4) : 113-168.
- 1966: Some changes in the nomenclature of British Lepidoptera. Part 4. Microlepidoptera. *Entomologists' gazette* 17 : 213-235.
- 1967: Some changes in the nomenclature of British Lepidoptera. Part 5. Microlepidoptera. *Entomologists' gazette* 18 : 45-47.
- Bradley, J.D.; Tremewan, W.G.; Tuck, K.; Hamilton, C.J. 1985: Comments on the proposed conservation of *Laspeyresia* Hübner [1825]. Z.N. (S.) 2421. *Bulletin of zoological nomenclature* vol. 42 : 9-10.
- Brants, A.A. 1913: (Reported in Verslag van de Acht-en-Zestigste Zomervergadering der Nederlandse Entomologische Vereeniging). *Tijdschrift voor Entomologie* 56 : i-xxii.
- Braun, A. 1948: Elachistidae of North America (Microlepidoptera). *Memoirs of the American Entomological Society*, no. 13. ii + 110 p.
- Bremer, O.; Grey, W. 1853: Beiträge zur Schmetterlings-Fauna des Nordlichen China's & St Petersburg. 23 p.
- Bretherton, R.F.; Goater, B.; Lorimer R.I. 1983: Noctuidae: Cuculliinae to Hypeninae. In Heath, J.; Emmett, A.M. eds, *The moths and butterflies of Great Britain and Ireland*. 459 p.
- Brock, J.P. 1971: A contribution towards an understanding of the morphology and phylogeny of the ditrysian Lepidoptera. *Journal of natural history* 5 : 29-102.
- Brown, R.L. 1979: The valid generic and tribal names for the Codling Moth, *Cydia pomonella* (Olethreutinae: Tortricidae). *Annals of the Entomological Society of America* 72 : 565-567.
- Bryk, F. 1937: Arctiidae: subfamilies Callimorphinae, Nyctemeralinae. In *Lepidopterorum Catalogus*, pars 82. 105 p.
- Buller, A.P. 1905: Notes on the occurrence of some rare species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 37 : 331-333.
- Buller, W.L. 1873: Notice of a new species of moth from New Zealand. *Transactions and proceedings of the New Zealand Institute* 5 : 279-280.
- Busck, A. 1908: A generic revision of American moths of the family Oecophoridae, with descriptions of new species. *Proceedings of the United States National Museum* 35 : 187-207.
- Busck, A. in Quaintance, A.L.; Wood, W.B. 1916: *Laspeyresia molesta*, an important new insect enemy of the peach. *Journal of agricultural research, Dept. of Agriculture, Washington D.C.*, vol. VII no. 8 : 373-377.

- Butler, A.G. 1868: Catalogue of diurnal Lepidoptera of the family Satyridae in the collection of the British Museum. London, printed by order of the Trustees. vi + 211 pp.
- 1874: The Lepidoptera of New Zealand. In *Insects*, by Adam White and Arthur Gardiner Butler. In Richardson, J.; Gray, J.E. eds. *The zoology of the voyage of H.M.S. Erebus & Terror*, under the command of Sir James Clark Ross R.N., F.R.S. during the years 1839 to 1843. London, E.W. Janson. 51 p., 10 plates.
- 1876: List of the butterflies now known to inhabit New Zealand, with descriptions of a new genus, and a new species, in the collection of John D. Enys, Esq. *Entomologists' monthly magazine* 13 : 152–154.
- 1877: On two collections of heterocerous Lepidoptera from New Zealand, with descriptions of new genera and species. *Proceedings of the Zoological Society of London for 1877* : 379–407.
- 1878: On the butterflies of New Zealand. *Transactions and proceedings of the New Zealand Institute* 10 : 263–276.
- 1879a: On a small collection of Heterocerous Lepidoptera, from New Zealand. *Cistula Entomologica* 2 : 487–511.
- 1879b: Descriptions of new species of Lepidoptera from Japan. *Annals and magazine of natural history (5th series)* vol. 4, no. 24 : 437–457.
- 1880: On a collection of Lepidoptera Heterocera from Marlborough Province, New Zealand. *Cistula Entomologica* 2 : 541–562.
- 1881a: On a second collection of Lepidoptera, made in Formosa, by H.E. Hobson. *Proceedings of the Zoological Society of London*, 1880 : 666–691.
- 1881b: On a collection of nocturnal lepidoptera from the Hawaiian Islands. *Annals and magazine of natural history (5th series)*, vol. 7: 392–408.
- 1882a: On a small collection of Lepidoptera from Melbourne. *Annals and magazine of natural history (5) IX* : 84–103.
- 1882b: Heterocerous Lepidoptera collected in Chili by Thomas Edmonds, Esq. *Transactions of the Entomological Society of London*, 1882, pt III (Sept.) : 339–427.
- 1884a: On a new genus of butterfly from New Zealand. *Annals and magazine of natural history (5) 13* : 171–173. [Reprinted in *New Zealand journal of science (Dunedin)* 2 : 159–160, July 1884.]
- 1884b: On the duplication of generic names employed by Walker. *Entomologists' monthly magazine* 21 : 133.
- 1886: Descriptions of 21 new genera and 103 new species of Lepidoptera - Heterocera from the Australian Region. *Transactions of the Entomological Society of London for 1886*, pt IV : 381–441.
- 1892: On the genus *Hypocala*, a group of Noctuid Moths. *Annals and magazine of natural history (6) 10* : 17–22.
- Chapman, T.A. 1910: On *Zizeeria* (Chapman), *Zizera* (Moore), a group of Lycaenid Butterflies. *Transactions of the Entomological Society of London*, 1910 : 479–497.
- 1981: On the British (and a few continental) species of *Scoparia*. *Transactions of the Entomological Society of London*, 1911 : 501–518.
- Chappell, A.V. 1934: Life histories of some New Zealand Lepidoptera, with notes on the male genitalia and larval chaetotaxy. *Stylops* 3 : 163–177.
- Clarke, C.E. 1920: New Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 52 : 35.
- 1926: New species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 56 : 417–421.
- 1934: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 64 : 11–16.
- Clarke, J.F.G. 1955: Catalogue of the type specimens of Microlepidoptera in the British Museum (Natural History) described by Edward Meyrick, vol. 1. London, Trustees of the British Museum (Natural History). 332 p.
- 1969: Gelechiidae (D–Z). In Catalogue of the type specimens of Microlepidoptera in the British Museum (Natural History) described by Edward Meyrick, vol. 7. London, Trustees of the British Museum (Natural History). 531 p.
- 1971: The Lepidoptera of Rapa Island. *Smithsonian contributions to zoology* 56. 282 p.
- Clemens, B. 1859: Contributions to American entomology. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1859 : 256–262.
- 1860: Contributions to American Lepidopterology. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1860 : 4–15.
- Clerck, C.A. 1759–1764: *Icones Insectorum rariorum cum nominibus eorum trivialibus, locisque e C. Linnaei ... Systema Naturae allegatis. 2nd pt. Holmiae. [viii, iii, 3:] 55 pl., coloured.*
- Cockerell, T.D.A. 1913: The noctuid genus *Alysia*. *The entomologist* 46 : 15.
- Common, I.F.B. 1954a: The Australian armyworms of the genus *Persectania* (Lepidoptera: Noctuidae). *Australian journal of zoology* 2 : 86–99.
- 1954b: A study of the ecology of the adult bogong moth, *Agrotis infusa* (Boisd.) (Lepidoptera: Noctuidae), with special reference to its behaviour during migration and aestivation. *Australian journal of zoology* 2 : 223–263.
- 1958a: The Australian cutworms of the genus *Agrotis* (Lepidoptera: Noctuidae). *Australian journal of science* 6 : 69–88.
- 1958b: The genera of the Australian Tortricidae (Lepidoptera). *Proceedings of the Tenth International Congress of Entomology*, 1956, vol. 1 : 289–295.
- 1961: The generic position of the Australian light-brown apple moth (Lepidoptera: Tortricidae). *Proceedings of the Linnean Society of New South Wales* 86 : 177–182.

- 1964: A new genus for the Australian lucerne leafroller (Lepidoptera: Tortricidae). *Proceedings of the Linnean Society of New South Wales* 88 : 298–300.
- 1965a: The identity and distribution of species of *Pseudaletia* (Lepidoptera: Noctuidae) in Australia. *Journal of the Entomological Society of Queensland* 4 : 14–17.
- 1965b: A revision of the Australian Tortricini, Schoenotenini and Chlidanotini (Lepidoptera: Tortricidae: Tortricinae). *Australian journal of zoology* 13 : 613–726.
- 1970: Lepidoptera (moths and butterflies). Pp. 765–866 in MacKerras, I.M. ed, *The insects of Australia*. Victoria, CSIRO / Melbourne University Press.
- 1975: Evolution and classification of the Lepidoptera. *Annual review of entomology* 20 : 183–203.
- Common, I.F.B.; Waterhouse, D.F. 1972: Butterflies of Australia. Sydney, Angus & Robertson. 498 p.
- 1981: Butterflies of Australia. Sydney, Angus & Robertson. Revised edition, xiv + 682 p.
- Corbet, A.S.; Tams, W.H.T. 1943: Keys for the identification of the Lepidoptera infesting stored food products. *Proceedings of the Zoological Society of London (B)* 118 : 55–148.
- Cramer, P. [1775–] 1779–1782: De uitlandsche Kapellen, &c – Papillons exotiques des trois parties du Monde – l’Asie, l’Afrique et l’Amerique, &c. 4 Deel.
- Craw, R.C. 1978: Revision of genus *Argyrophenga* (Lepidoptera: Satyridae). *New Zealand journal of zoology* 5 : 751–768.
- 1986: Review of the genus *Notoreas* (*sensu auctorum*) (Lepidoptera: Geometridae: Larentiinae). *New Zealand journal of zoology* 13 : 131–140.
- 1987: Revision of the genus *Helastia* sensu stricto with description of a new genus (Lepidoptera: Geometridae: Larentiinae). *New Zealand journal of zoology* 14 : 269–293.
- Crosby, T.K.; Dugdale, J.S.; Watt, J.C. 1976: Recording specimen localities in New Zealand: an arbitrary system of areas and codes defined. *New Zealand journal of zoology* 3 : 69 + map.
- Curtis, J. 1825: *British entomology* 2: 51–98. London.
- 1827: *British entomology* 4: (esp. pl. 165). London.
- 1828: *British entomology* 5: (esp. p. 5, pl. 12). London.
- 1832: *British entomology* 9: (esp. folio 391). London.
- 1833: *The entomological magazine* 1: 190.
- Dalla Torre, K.W. von; Strand, E.** 1929: Psychidae. *Lepidopterorum Catalogus* 34: 1–211.
- Davis, D.R. 1967: A revision of the moths of the subfamily Prodoxinae (Lepidoptera: Incurvariidae). *United States National Museum bulletin* 255. 170 p.
- 1978: The North American moths of the genera *Phaeoses*, *Opogona*, and *Oinophelia*, with a discussion of their suprageneric affinities (Lepidoptera: Tineidae). *Smithsonian contributions to zoology* 282. 39 p.
- 1986: A new family of monotypian moths from austral South America (Lepidoptera: Paleaphatidae), with a phylogenetic review of the Monotrysia. *Smithsonian contributions to zoology* 434. 202 p.
- Denis, J.N.C.M.; Schiffermüller, I. 1775: Ankündigung eines systematischen Werkes von den Schmetterlingen der Wienergegend ... Wien. 322 p.
- Desmarest, E. 1857: Papillons nocturnes de Paris. In Chenu, J.C. ed, *Encyclopédie d’Histoire Naturelle*. [iii] + 312 p., 39 pl.
- Diakonoff, A. 1941: Notes and descriptions of microlepidoptera (I) (7th paper on Indo-Malayan and Papuan microlepidoptera). *Treubia* 18 (2) : 395–438, pl. 17–22.
- 1956: Records and descriptions of microlepidoptera (8). *Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie te Leiden*, 29. 60 p.
- 1959: Revision of *Cryptaspasma* Walsingham, 1900 (Lepidoptera, Tortricidae). *Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie te Leiden*, 43. 60 p., 13 pl.
- 1964: Further records and descriptions of the species of *Bactra* Stephens (Lepidoptera, Tortricidae). *Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie te Leiden*, 70. 81 p.
- 1974: The south asiatic Polyorthini, with notes on species of *Polyortha* Dognin (Lepidoptera, Tortricidae). *Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie te Leiden*, 131. 86 p.
- 1986: Glyptapterigidae auctorum sensu latu (Glyptapterigidae sensu Meyrick, 1913); containing Tortricidae: Hilarographini, Choreutidae, Brachodidae (partim), Immidae, and Glyptapterigidae. *Microlepidoptera Palaearctica* 7. 436 p.
- Dollimore, E.S. (compiler) 1962: The New Zealand Guide, superseding Wise’s Index to every place in New Zealand. 3rd edition. Dunedin, H. Wise & Co.
- Donovan, E. 1805: An epitome of the natural history of the insects of New Holland, New Zealand, New Guinea, Otaheite and other islands in the Indian, Southern and Pacific Oceans &c. London. 41 pl., coloured, with descriptive letterpress.
- Doubleday, E. 1843: Lepidoptera. Pp. 283–289 in List of the annulose animals hitherto recorded as found in New Zealand, with the descriptions of some new species. In Dieffenbach, E. ed., *Travels in New Zealand: with contributions to the geography, geology, botany and natural history of the country*, vol. 2. London, John Murray.
- 1845: Description of new or imperfectly described diurnal Lepidoptera. *Annals and magazine of natural history* (1) 16 : 304–308.
- Doubleday, E.; Westwood, J.O. 1845–1852: The genera of diurnal Lepidoptera: comprising their generic

- characters, a notice of their habits and transformations, and a catalogue of the species of each genus. Illustrated with eighty-six plates by William C. Hewitson. [1851]: vol. 2, pp. 251–234: Nymphalidae continued, Morphidae, Brassolidae, Satyridae, Libytheidae, Erycinidae, Lycaenidae and Hesperiidae; with a supplement by John O. Westwood. [1852]: vol. 3, plates. London, Longman, Brown, Green, and Longmans.
- [1849] 1850: A synonymic list of British Lepidoptera ... excluding the family Tineidae, &c. 27 p.
- Druce, H. 1884: Zygaenidae. Pp. 33–112, pl. VI–XI in Godman, F.D. & Salvin, O., Biologia Centrali-America: Lepidoptera Heterocera.
- Drury, D. 1770–1782: Illustrations of natural history, wherein are exhibited figures of exotic insects, &c. 3 vols.
- Dufay, C. 1970: Insectes lépidoptères: Noctuidae, Plutinae. *Faune de Madagascar* 31 : 1–198.
- Dugdale, J.S. 1964: Insects of Campbell Island. Appendix. Lepidoptera; Geometridae. *Pacific insects monograph* 7 : 607–623.
- 1966a: A new genus for the New Zealand 'elusive Tortrix' (Lepidoptera: Tortricidae: Tortricinae). *New Zealand journal of science* 9 : 391–398.
- 1966b: A revision of New Zealand Schoenotenini and Cnephasini (Lepidoptera: Tortricidae: Tortricinae). *New Zealand journal of science* 9 : 731–775.
- 1971a: A new species of *Proterodesma* Meyrick from the Chatham Is. (Lepidoptera: Tineidae). *New Zealand journal of science* 14 : 62–65.
- 1971b: Entomology of the Aucklands and other islands south of New Zealand: Lepidoptera, excluding non-crambine Pyralidae. *Pacific insects monograph* 27 : 55–172.
- 1971c: The Muehlenbeckia leafminer, *Zapryasta* [sic] *calliphana* (Lepidoptera: Cosmopterigidae). *New Zealand entomologist* 5 (1) : 65–69.
- 1973a: A note on Kermadec Islands Lepidoptera. *New Zealand entomologist* 5(2) : 197–203.
- 1973b: The genus *Plutella* (Hyponomeutidae) in New Zealand and the family position of *Circoxena* (Lepidoptera). *New Zealand journal of science* 16 : 1009–1023.
- 1974: Female genital configuration in the classification of Lepidoptera. *New Zealand journal of zoology* 1 (2) : 127–146.
- 1978: A new name for *Tortrix encausta* Philpott, 1930 (Lepidoptera: Tortricidae). *New Zealand entomologist* 6 (4) : 443.
- 1979: A new generic name for the New Zealand species previously assigned to *Simaethis auctorum* (Lepidoptera: Choreutidae), with description of a new species. *New Zealand journal of zoology* 6 : 461–466.
- 1986: *Trioxycanus* Dumbleton, 1966 (Lepidoptera), a genus based on a misidentified type species, with proposal of new names for the taxonomic genus and species involved. Z.N.(S.)2462.
- Bulletin of zoological nomenclature 43 (1) : 46–49.
- 1987a: *Proditrix*, a new genus for *Titanomis tetragona* Hudson and *Plutella megalynta* Meyrick and allies (Lepidoptera: Yponomeutidae), shoot destroyers of Monocotyledones. *New Zealand journal of zoology* 14 : 99–112.
- 1987b: A new genus for G.V. Hudson's "curved case moth" (Lepidoptera: Psychidae). *The New Zealand entomologist* 9 : 107–115.
- Dumbleton, L.J. 1966: Genitalia, classification and zoogeography of the New Zealand Hepialidae (Lepidoptera). *New Zealand journal of science* 9 (4) : 920–981.
- Duponchel, P.A.J. 1838: Histoire naturelle des Lépidoptères ou Papillons du France, vol. 11. Paris. 720 p., pl. 287–314.
- Dyar, H.G. 1920: A generic subdivision of the genus *Plutella*. *Journal of the New York Entomological Society* 10 : 79–82.
- Edwards, E.D. 1977: *Junonia erigone* (Cramer) (Lepidoptera: Nymphalidae) recorded from Australia. *Australian entomological magazine* 4 (3) : 41–43.
- Enderlein, G. 1905: Beitrag zur Kenntnis der antarktischen Landarthropoden. *Pringleophaga*, eine neue Schmetterlings Gattung aus dem Antarktischen Gebiet. *Zoologische Anzeiger* 29 : 119–125.
- Evans, W.P. 1931: Traces of a lepidopterous insect from the Middle Waikato coal measures. *Transactions and proceedings of the New Zealand Institute* 62 : 99–100, pl. 12.
- Fabricius, J.C. 1775: *Systema entomologiae, sistens insectorum classes, ordines, genera, species adiectis synonymis, locis, descriptionibus, observationibus. Flensburgi et Lipsiae, Korte.* 832 p.
- 1781: Species insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosis adiectis observationibus, descriptionibus. *Hamburgi et Kilonii, Bohn.* 2 vols.
- 1787: *Mantissa insectorum sistens eorum species nuper detectus adiectis characteribus, genericis differentiis specificis, amendmentibus, observationibus. Hafniae, Proft.* 2 vols.
- 1793: *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species, adiectis synonymis, locis, observationibus, descriptionibus. Hafniae, Proft. Tome III, pars I.* 347 p.
- 1794: *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species, adiectis synonymis, locis, observationibus, descriptionibus. Hafniae, Proft. Tome III, pars II.* 349 p.
- 1807: *Magazin für Insectenkunde* 6 : 281. Herausgegeben von K. Illiger.
- Falkovitsh, N. 1965: New eastern-asiatic species of leaf-rollers (Lepidoptera, Tortricidae). *Entomologiske Obozreniye* 44 : 414–437.
- Felder, C. 1861: Lepidoptera nova Columbiae. *Wiener entomologische Monatschrift* 5: 97–111.

- 1862: Lepidoptera nova Columbiae. *Wiener entomologische Monatschrift* 6 : 405-427.
- Felder, C.; Felder, R. 1865: Reise der österreichischen Fregatte Novara um die Erde (zoologischer Theil), Band 2 (Abtheilung 2), Hefte 2 (plates 22-47).
- Felder, C.; Rogenhofer, A.F. 1874: Reise der österreichischen Fregatte Novara um die Erde (zoologischer Theil), Band 2 (Abtheilung 2), Hefte 4 (Plates 75-107).
- 1875: Reise der österreichischen Fregatte Novara um die Erde (zoologischer Theil), Band 2 (Abtheilung 2), Hefte 5 (plates 121-140).
- Fereday, R.W. 1869: *Sesia tipuliformis* in New Zealand. *Entomologists' monthly magazine* 6 : 146.
- 1872: Observations on a paper read by Mr A. Bathgate before the Otago Institute; 11th January, 1870. "On the Lepidoptera of Otago". *Transactions and proceedings of the New Zealand Institute* 4 : 214-218.
- 1874: Observations on the occurrence of a butterfly new to New Zealand, of the genus *Danais*. *Transactions and proceedings of the New Zealand Institute* 6 : 183-186.
- 1876: Description of a species of butterfly belonging to the family Satyridae Westwood. *Transactions and proceedings of the New Zealand Institute* 8 : 302-304.
- 1877a: Description of a species of *Catocala*, new to science. *Transactions and proceedings of the New Zealand Institute* 9 : 457-459.
- 1877b: Brief observations on the genus *Chrysophanus*, as represented in New Zealand. *Transactions and proceedings of the New Zealand Institute* 9 : 460-463.
- 1878a: Supplementary description of species or varieties of *Chrysophani* (Lepidoptera Rhopalocera) inhabiting New Zealand. *Transactions and proceedings of the New Zealand Institute* 10 : 252-259.
- 1878b: Description of new genera and species of Psychidae. *Transactions and proceedings of the New Zealand Institute* 10 : 260-263.
- 1879: Description of a new genus? and species of butterfly of the subfamily Satyrinae. *Entomologists' monthly magazine* 16 : 129-130.
- 1880: Description of a ?new species of the family Leucanidae, and a ?new species of the genus ?*Chlenias*. *Transactions and proceedings of the New Zealand Institute* 12 : 267-270.
- 1883a: Occurrence of a species of *Ophideres*, Boisd., new to New Zealand. *Transactions and proceedings of the New Zealand Institute* 15 : 192-193.
- 1883b: Description of a species of butterfly new to New Zealand and probably to science. *Transactions and proceedings of the New Zealand Institute* 15 : 193-195.
- 1883c: Description of two new species of Heteropterous [sic] Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 15 : 195-196.
- 1883d: Description of a new species of *Cidaria* (Lepidoptera). *New Zealand journal of science (Dunedin)* 1 : 531. (See Fereday 1884.)
- 1884: Description of a new species of *Cidaria* (Lepidoptera). *Transactions and proceedings of the New Zealand Institute* 16 : 119-120.
- Fernald, C.H. 1908: The genera of the Tortricidae and their types. Amherst (Mass.), Carpenter & Morehouse. 67 p.
- Field, W.D. 1971: Butterflies of the genus *Vanessa* and of the resurrected genera *Bassaris* and *Cynthia* (Lepidoptera: Nymphalidae). *Smithsonian contributions to zoology* 84 : 1-105.
- 1975: Ctenuchid moths of *Ceramidia* Butler, *Ceramidiodes* Hampson, and the *caca* species group of *Antichloris* Huebner. *Smithsonian contributions to zoology* 198 : 1-45.
- Fletcher, D.S. 1966: Some changes in the nomenclature of British Lepidoptera. Part 1. *Entomologists' gazette* 17 : 9-18.
- 1967: A revision of the Ethiopian species and a checklist of the world species of *Cleora* (Lepidoptera: Geometridae). *Bulletin of the British Museum (Natural History), entomology, supplement* 8 : 1-119.
- 1979: In Nye, I.W.B., The generic names of moths of the world, volume 3. London, Trustees of the British Museum (Natural History). 243 p.
- Fletcher, D.S.; Nye, I.W.B. 1984: The generic names of moths of the world, volume 5. London, Trustees of the British Museum (Natural History). 185 p.
- Fletcher, T.B. 1929: A list of the generic names used for Microlepidoptera. *Memoirs of the Department of Agriculture in India, entomological series* 11 : 1-246.
- Forbes, W.T.M. 1954: Lepidoptera of New York and neighbouring states: part III, Noctuidae. *Cornell University Agricultural Experiment Station memoir* 329. 433 p.
- Foster, S.P.; Clearwater, J.R.; Muggleston, S.J.; Dugdale, J.S.; Roelofs, W.L. 1986: Probable sibling species complexes within two described New Zealand leaf-roller moths. *Naturwissenschaften* 73 : 156-158.
- Foster, S.P.; Roelofs, W.L. 1987: Sex pheromone differences in populations of the brown-headed leaf-roller, *Ctenopseustis obliquana*. *Journal of chemical ecology* 13 : 623-639.
- Fox, K.J. 1970: A new species of *Melanchia* (Lepidoptera: Noctuidae) from New Zealand. *Records of the Dominion Museum, Wellington*, 7 : 21-24.
- 1975: Migrant Lepidoptera in New Zealand 1973-1974. *New Zealand entomologist* 6 (1) : 66-69.
- 1978: The transoceanic migration of Lepidoptera to New Zealand - a history and a hypothesis on colonisation. *New Zealand entomologist* 6 (4) : 368-380.
- Franclemont, J.G. 1981: The identity of *Mamestra passa* and *Morrisonia peracuta* of Morrison (Lepidoptera: Noctuidae: Hadenniae). *Proceedings of the Entomological Society of Washington* 83 : 133-136.

- Gaede, M. 1939: Oecophoridae II. *Lepidopterorum Catalogus* pars 92 [continuation of pars 88], pp. 209–476.
- Gaedike, R. 1978: Versuch der phylogenetischen Gliederung der Epermeniidae der Welt (Lepidoptera). *Beiträge zur Entomologie* 28 : 201–209.
- Gaskin, D.E. 1971: A revision of the New Zealand Diptychophorini (Lepidoptera: Pyralidae: Crambinae). *New Zealand journal of science* 14: 759–809.
- 1973: Revision of New Zealand Chilonini (Lepidoptera: Pyralidae) and redescription of some Australian species. *New Zealand journal of science* 16 : 435–463.
- 1974: The species of *Pareromene* Osthelder (Pyralidae: Crambinae: Diptychophorini) from the western South Pacific, with further notes on the New Zealand species. *Journal of entomology (B)* 43 (2) : 159–184.
- 1975: Revision of the New Zealand Crambini (Lepidoptera: Pyralidae: Crambinae). *New Zealand journal of zoology* 2 (3) : 265–363.
- 1986: Morphology and reclassification of the Australian, Melanesian and Polynesian *Glaucocaris* Meyrick (Lepidoptera: Crambinae: Diptychophorini). *Australian journal of zoology, supplementary series* 115. 75 p.
- 1987: Supplement to New Zealand Crambinae (Lepidoptera: Pyralidae) – corrections, descriptions of females of two species, and notes on structure, biology, and distribution. *New Zealand journal of zoology* 14 : 113–121.
- Gibbs, G.W. 1962: The New Zealand genus *Metacrias* Meyrick (Lepidoptera: Arctiidae). Systematics and distribution. *Transactions of the Royal Society of New Zealand (zoology)* 2 : 153–167.
- 1980a: Reinstatement of a New Zealand copper butterfly, *Lycaena rauparaha* (Fereday, 1877). *New Zealand journal of zoology* 7 (1) : 105–114.
- 1980b: New Zealand butterflies, identification and natural history. Auckland, William Collins Ltd. 207 p.
- 1983: Evolution of Micropterigidae (Lepidoptera) in the SW Pacific. *Geojournal* 76 : 505–510.
- Godart, J.B. 1819: Histoire naturelle. Entomologie, ou histoire naturelle des Crustacés, des Arachnides et des Insectes. Encyclopédie Méthodique, vol. 9, pp. 1–328. Paris & Liège. [Note: Catalogue of the books ... in the British Museum (Natural History), vol. 2, E-K, 1904, p. 527, notes that "pp. 1–328, 706, 707 and all extra-European Hesperides to p. 793 were by Latreille; the remaining portion was by J.B. Godart".]
- 1824: Histoire naturelle des Lépidoptères ou papillons diurnes des environs de Paris, décrits par M. Godart, peints par M. C. Vautier. Paris, Crevot.
- Goeze, J.A.E. 1783: Entomologische Beiträge zu des Ritter Linné Zwölften Ausgabe des Natursystems. Pp. i–xx, 1–178. Leipzig.
- Gozmany, L. 1978: Lecithoceridae. Microlepidoptera Palaeoarctica, vol. 5 (text), pp. 1–306. Vienna, Verlag Georg Fromme.
- Gozmany, L.A.; Vari, L. 1973: The Tineidae of the Ethiopian Region. *Transvaal Museum memoir* 18. 238 p.
- Green, C.J.; Dugdale, J.S. 1982: Review of the genus *Ctenopseustis* Meyrick (Lepidoptera: Tortricidae) with reinstatement of two species. *New Zealand journal of zoology* 9(4) : 427–435.
- Grote, A.R. 1873a: Description of a butterfly new to the Lower Lake Region. *Bulletin of the Buffalo Society of Natural Sciences* 1 : 178–179.
- 1873b: Description of three genera of Noctuidae. *Bulletin of the Buffalo Society of Natural Sciences* 1 : 180–182.
- 1874: On the Noctuidae of North America. *Report of the Peabody Academy of Science* 6 : 21–38.
- Guenée, A. 1845: Essai sur une nouvelle classification des Microlépidoptères (Part 1). *Annales de la Société Entomologique de France* (2) 3 : 297–344.
- 1852a: Noctuélites, Tome 1. In *Histoire naturelle des Insectes. Species général des Lépidoptères*, par Mm. Boisduval et Guenée, vol. 5. 407 p.
- 1852b: Noctuélites, Tome 2. In *Histoire naturelle des Insectes. Species général des Lépidoptères*, par Mm. Boisduval et Guenée, vol. 6. 444 p.
- 1852c: Noctuélites, Tome 3. In *Histoire naturelle des Insectes. Species général des Lépidoptères*, par Mm. Boisduval et Guenée, vol. 7. 442 p.
- 1854: Deltoides et Pyralites. In *Histoire Naturelle des Insectes. Species général des Lépidoptères*, par Mm. Boisduval et Guenée, vol. 8. 446 p.
- 1857a: Uranides et Phalénites, Tome 1. In *Histoire naturelle des Insectes. Species général des Lépidoptères*, par Mm. Boisduval et Guenée, vol. 9. 514 p.
- 1857b: Uranides et Phalénites, Tome 2. In *Histoire naturelle des Insectes. Species général des Lépidoptères*, par Mm. Boisduval et Guenée, vol. 10. 584 p.
- 1868: New species &c, of heterocerous Lepidoptera from Canterbury, New Zealand collected by Mr R.W. Fereday. *Entomologists' monthly magazine* 5 : 1–6, 38–43, 61–65, 92–95.
- Hamilton, A. 1909: Notes on Lepidoptera collected by H. Hamilton in various localities in the Queenstown district, Otago, between November 1907 and March 1908. *Transactions and proceedings of the New Zealand Institute* 41: 44–48.
- Hampson, G.F. 1894: Moths. In Blanford, W.T. (ed.), *The Fauna of British India, including Ceylon and Burma*. Vol. ii. xxii + 609 p.
- 1895: On the classification of the Schoenobiinae and Crambinae, two subfamilies of moths of the family Pyralidae. *Proceedings of the Zoological Society of London*, 1895 : 897–974.

- 1896: On the classification of three subfamilies of moths of the family Pyralidae: the Epipachinae, Endotrichinae and Pyralinae. *Transactions of the Entomological Society of London*, 1896: 451–550.
- 1897: On the classification of two subfamilies of moths of the family Pyralidae: the Hydrocampinae and Scoparianae. *Transactions of the Entomological Society of London*, 1897: 127–240.
- 1903a: Catalogue of the Noctuidae in the collection of the British Museum, London. Vol. 4, Catalogue of the Lepidoptera Phalaenae in the British Museum. xx + 689 p.
- 1903b: The moths of India. Supplementary paper to the volumes in the 'Fauna of British India', series ii, part 8. *Journal of the Bombay Natural History Society* 14: 639–659.
- 1905: Catalogue of the Noctuidae in the collection of the British Museum; Vol. 5, Hadeninae. Catalogue of the Lepidoptera Phalaenae in the British Museum. xvi + 634 p.
- 1906: Catalogue of the Lepidoptera Phalaenae in the British Museum, Vol. 6. 532 p.
- 1909: Catalogue of the Lepidoptera Phalaenae in the British Museum, Vol. 8. xiv + 583 p.
- 1911: Descriptions of new genera and species of Syntomidae, Arctiidae, Agaristidae and Noctuidae. *Annals and magazine of natural history, London*, (8) 8: 393–445.
- 1913a: Catalogue of the Noctuidae in the collection of the British Museum, Vol. 12. Catalogue of the Lepidoptera Phalaenae in the British Museum. xiii + 626 p.
- 1913b: Descriptions of new genera and species of Noctuidae. *Annals and magazine of natural history, London*, (8) 12: 580–601.
- 1914: New genera and species of Noctuidae. *Annals and magazine of natural history, London*, 13: 197–223.
- 1918a: Descriptions of new genera and species of Amatidae, Lithosiidae and Noctuidae. *Novitates Zoologicae, Tring*, 25: 93–217.
- 1918b: Descriptions of the subfamily Pyraustinae (continuation). *Annals and magazine of natural history, London*, (9) 2: 181–196.
- Hardwick, D.F. 1965: The corn earworm complex. *Memoirs of the Entomological Society of Canada* 40. 246 p.
- 1970: A generic revision of the North American Heliothidinae (Lepidoptera: Noctuidae). *Memoirs of the Entomological Society of Canada* 73. 59 p.
- Harris, A.C. 1982: Notes on introduced Lepidoptera associated with bananas in New Zealand. *New Zealand entomologist* 7(3): 329–332.
- Haworth, A.H. 1811: Lepidoptera Britannica, ... sistens digestionem novam Insectorum Lepidopterorum quae in magna Britannia reperiuntur, [etc.], Pt 3, pp. 377–512.
- [1812] 1828: Lepidoptera Britannica, ... [etc.], Pt 4, pp. 513–609.
- Hawthorne, E.F. 1897: Descriptions of two new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 29: 282–283.
- Heinemann, H. von 1870: Kleinschmetterlinge, 2. *In Die Schmetterlinge Deutschlands und der Schweiz, systematisch bearbeitet von H.V. Heinemann [(and afterwards)] M.F. Wocke, nebst analytischen Tabellen, etc.* Braunschweig. Vol. 2, Heft 1.
- Helson, G.A.H. 1967: Adult periodicity of *Wiseana* spp. (family Hepialidae) in New Zealand as revealed by light traps. *Transactions of the Royal Society of New Zealand, zoology* 9, 8: 79–91.
- Hemming, F. 1933: Additional notes on the types of certain butterfly genera. *The Entomologist, London*, 66: 222–225.
- 1937: Hübner. A bibliographical and systematic account of the entomological works of Jacob Hübner and of the supplements thereto by Carl Geyer, Gottfried von Frölich, and Gottlieb August Wilhelm Herrich Schaeffer. Vol. 1. London, Royal Entomological Society of London. 605 p.
- 1967: The generic names of the butterflies and their type-species (Lepidoptera: Rhopalocera). *Bulletin of the British Museum (Natural History), entomology, supplement* 9. 509 p.
- Heppner, J.B. 1982: Synopsis of the Glyphipterigidae (Lepidoptera: Copromorphoidea) of the World. *Proceedings of the Entomological Society of Washington* 84 (1): 38–66.
- (ed.) 1984: Atlas of Neotropical Lepidoptera. Checklist: Part 1, Micropterigoidea – Immoidea. The Hague, Boston, and Lancaster, Dr W. Junk Publishers. 112 p.
- 1985: The sedge moths (Lepidoptera: Glyphipterigidae) of North America. *Flora & fauna handbook no. 1*. Gainesville, Florida. 254 p.
- Heppner, J.; Duckworth, D.W. 1981: Classification of the superfamily Sesioidea (Lepidoptera: Ditrysia). *Smithsonian contributions to zoology* 314. 144 p.
- Herrich-Schäffer, G.A.W. 1853: Die Schaben und Feder-motten. *In Systematische Bearbeitung der Schmetterlinge von Europa, zugleich als Text Revision und Supplement zu Jacob Hübner's Sammlung europäischer Schmetterlinge*. Vol. 5. 394 p.
- 1855: Synopsis familiarum Lepidopterorum. Pp. 82–132 in *Systematische Bearbeitung der Schmetterlinge von Europa*, etc. Vol. 6.
- Hewitson, W.C. 1875: Description of three new butterflies. *Entomologist's monthly magazine* 12: 9–10.
- Heyden, C. von 1843: Amtlicher Bericht der Versamm-lung der Naturforscher zu Mainz. 208 p.
- Higgins, L.G. 1975: The classification of European butterflies. London, Glasgow, etc., William Collins Sons & Co. Ltd. 320 p.
- Hodges, R.W. 1966: Review of New World species of *Batrachedra*, with description of three new genera (Lepidoptera: Gelechioidae). *Transactions of the American Entomological Society* 92: 585–651.
- 1974: Gelechioidae Oecophoridae (in part). Fascicle 6.2 of Dominick, R.B. et al. (eds), *The moths of America north of Mexico*. London, E.W. Classey & R.B.D. Publications Inc. 142 p.

- 1978: Gelechioidea Cosmopterigidae. Fascicle 6.1 of Dominick, R.B. et al. (eds), *The moths of America north of Mexico, including Greenland*. London, E.W. Classey Ltd and The Wedge Entomological Foundation. 166 p.
- (ed.) 1983: Checklist of the Lepidoptera of America north of Mexico. London, E.W. Classey and The Wedge Entomological Research Foundation. 284 p.
- 1985: Comments on the proposed conservation of *Laspeyresia* Hübner, [1825]. Z.N.(S)2421. *Bulletin of zoological nomenclature* 42: 8.
- Holloway, B.A. 1962: *Melanitis leda* (Linn.) and other migrant butterflies in New Zealand during 1962. *Records of the Dominion Museum [Wellington N.Z.]* 4 (8): 79–82.
- Holloway, J.D. 1977: The Lepidoptera of Norfolk Island: their biogeography and ecology. *Series Entomologica* 13. 291 p.
- 1979: A survey of the Lepidoptera, biogeography and ecology of New Caledonia. *Series Entomologica* 15. 588 p.
- Holloway, J.D.; Bradley, J.D.; Carter, D.J. (in press): CIE *Guides to insects of importance to man. 1. Lepidoptera*. Wallingford, CAB International.
- Horak, M. 1984: Assessment of taxonomically significant structures in Tortricinae (Lep., Tortricidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft (Bull. Soc. Ent. Suisse)* 57: 3–64.
- Horning, D.S.; Greenwood, D.J. 1977: A portable cabinet for rearing immature insects. *The New Zealand Entomologist* 6 (30): 293–295.
- Howes, W.G. 1906: Some new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 38: 510–511, pl. 44 fig. 1, 2, 3, and 5.
- 1908: Further notes on Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 40: 533–534.
- 1911: New species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 43: 127–128, pl. I.
- 1912: New species of Lepidoptera, with notes on the larvae and pupae of some New Zealand butterflies. *Transactions and proceedings of the New Zealand Institute* 44: 203–208.
- 1914: New Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 46: 95–96.
- 1917: New Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 49: 274.
- 1942: New Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 71: 277–278.
- 1943: Descriptions of two new species of Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 72: 371–372.
- 1945: New Lepidoptera. *Transactions of the Royal Society of New Zealand* 75: 65–67, pl. 7.
- 1946: Lepidoptera collecting at the Homer, with descriptions of new species. *Transactions of the Royal Society of New Zealand* 76: 139–147, pl. 8 and 9.
- Hübner, J. 1796: Sammlung Europäischer Schmetterlinge, Horde 8. Tineae-Schaben. Augsburg. 78 p., pl. 1–34.
- 1800–1809: Sammlung Europäischer Schmetterlinge, Horde 6. Pyralides-Zunsler. Augsburg. Pl. 1–32.
- 1810–1813: Sammlung Europäischer Schmetterlinge, Horde 8. Augsburg. Pl. 45–63.
- 1818: Zuträge zur Sammlung exotischer Schmetterlinge, Vol. 1. 40 p.
- 1819: Verzeichniss bekannter Schmettlinge [sic]. Augsburg. Signatures 2–11, pp. 17–176.
- 1820: Verzeichniss bekannter Schmettlinge [sic]. Augsburg. Signatures 12 and 13, pp. 177–208.
- 1821a: Sammlung Exotischer Schmetterlinge. Vol. 2 [1819–1827], pl. 1–225.
- 1821b: Verzeichniss bekannter Schmettlinge [sic]. Augsburg. Signature 16, pp. 241–256.
- 1822: Systematisch-Alphabetisches verzeichniss aller bisher bei den Fürbildungen zur Sammlung europäischer Schmetterlinge angegebenen Gattungsbennungen. Augsburg. 81 p.
- 1823: Verzeichniss bekannter Schmettlinge [sic]. Augsburg. Signatures 17–19, pp. 257–304.
- 1825: Verzeichniss bekannter Schmettlinge [sic]. Augsburg. Signatures 20–27, pp. 305–431.
- Hudson, G.V. 1889: Description of a new species of Arcitiidae from New Zealand. *The entomologist* 22: 53.
- 1898: New Zealand moths and butterflies (Macro-lepidoptera). London, West, Newman & Co. 144 p., 13 pl.
- 1903: On some new species of Macro-lepidoptera. *Transactions and proceedings of the New Zealand Institute* 35: 243–245.
- 1905a: On some new species of macro-lepidoptera in New Zealand. *Transactions and proceedings of the New Zealand Institute* 37: 355–358.
- 1905b: On macro-lepidoptera observed during the summer of 1903–4, including a note on the occurrence of a hawk-moth new to New Zealand. *Transactions and proceedings of the New Zealand Institute* 37: 358–360.
- 1908: Recent observations on New Zealand Macro-lepidoptera, including descriptions of new species. *Transactions and proceedings of the New Zealand Institute* 40: 104–107.
- 1909: Descriptions of four new species of Macro-lepidoptera from the southern islands. Pp. 67–69 in Chilton, C. (ed.), *Subantarctic islands of New Zealand*, vol. 1.
- 1913: A new *Scoparia* from New Zealand. *Entomologists' monthly magazine* 49: 250–251.
- 1918: Descriptions of new species of Lepidoptera from New Zealand. *Entomologists' monthly magazine (new series, vol. 4)* 64: 61–63.
- 1920: Description of two new species of Hepialidae from New Zealand. *Entomologists' monthly magazine* 56: 277.

- 1921: Description of a new species of the lepidopterous genus *Melanchra* Hübn. from New Zealand. *Entomologists' monthly magazine* 57 : 255.
- 1922: Descriptions of two new species of Lepidoptera from New Zealand. *Entomologists' monthly magazine* 58 : 196-197.
- 1923a: Descriptions of two new species of Lepidoptera from New Zealand. *Entomologists' monthly magazine* 59 : 64-65.
- 1923b: Descriptions of two new species of Lepidoptera from New Zealand. *Entomologists' monthly magazine* 59 : 129-130.
- 1923c: Descriptions of three new species of Lepidoptera from New Zealand. *Entomologists' monthly magazine* 59 : 179-181.
- 1923d: Descriptions of a large new tineid from New Zealand. *Entomologists' monthly magazine* 59 : 218.
- 1924: Description of a new noctuid from New Zealand. *Entomologists' monthly magazine* 60: 7-8.
- 1925: Descriptions of three new species of Lepidoptera from New Zealand. *Entomologists' monthly magazine* 61 : 220-221.
- 1928: The butterflies and moths of New Zealand. Wellington, Ferguson & Osborn Ltd. 386 p., 52 pl.
- 1939: A supplement to the butterflies and moths of New Zealand. Wellington, Ferguson & Osborn Ltd. Pp. 387-481, pl. 53-62.
- 1950: Fragments of New Zealand entomology. Wellington, Ferguson & Osborn Ltd. 188 p., ii + 17 pl.
- Hummel, A.D. 1823: *Essais entomologiques*, Vol. 1, part 3: Observations sur les insectes de 1823. *Monographia Pelophilarum. Novae species*. 48 p., 1 pl.
- Hutton, F.W. (ed.) 1904: *Index faunae Novae Zealandiae*. London, Dulau & Co. 372 p.
- Ichinose, T. 1973: A revision of some genera of the Japanese Plusiniae, with descriptions of a new genus and two new sub-genera (Lepidoptera, Noctuidae). *Kontyu* 41 (2) : 135-140.
- International Commission on Zoological Nomenclature 1961: International code of zoological nomenclature adopted by the XVth International Congress of Zoology. London, International Trust for Zoological Nomenclature. 176 p.
- 1985a: International code of zoological nomenclature, third edition, adopted by the XXth general assembly of the International Union of Biological Sciences. London, International Trust for Zoological Nomenclature. 338 p.
- 1985b: Opinion 1288. *Sphinx tipuliformis* Clerck, 1759 (Insecta, Lepidoptera): conserved. *Bulletin of zoological nomenclature* 42 : 17-18.
- 1985c: Opinion 1312. *Heliothis* Ochsenheimer, 1816 (Insecta, Lepidoptera): gender and stem designated. *Bulletin of zoological nomenclature* 42 : 158-159.
- Janse, A.T. 1951: The moths of South Africa, vol. 5, pt 3 (Gelechioidea), pp. 175-300. Pretoria, Transvaal Museum.
- 1954: The moths of South Africa, vol. 5, pt 4 (Glechiidae), pp. 301-464. Pretoria, Transvaal Museum.
- Jordan, K. 1939: On the constancy and variability of the differences between the Old World species of *Utetheisa* (Lepid., Arctiidae). *Novitates Zoologicae* 41 : 251-291.
- Karsholt, O.; Nielsen, E.S. 1976a: Systematisk forlegnelse over Danmarks sommerfugle. Klampenborg, Scandinavian Science Press. 128 p.
- 1976b: Notes on some Lepidoptera described by Linnaeus, Fabricius, and Ström. *Entomologica Scandinavica* 7 : 241-251.
- Kasy, F. 1973: Beitrag zur Kenntnis der Familie Stathmopodidae Meyrick, 1913 (Lepidoptera: Gelechioidea). *Tijdschrift voor Entomologie* 116 : 227-299.
- Kay, M. 1980: *Nyctemera amica* × *N. annulata* colony at Woodhill (Lepidoptera: Arctiidae). *New Zealand entomologist* 7 (2) : 154-158.
- Kennel, J. von; Eggers, F. 1933: Die Tympanalorgane der Lepidopteren. *Zoologische Jahrbücher, Abteilung für Anatomie und Ontogenie der Tiere* 57 : 1-104.
- Kirby, W.F. 1892: A synonymic catalogue of Lepidoptera Heterocera (moths). Vol. 1: Sphinges and Bombyces. London. 951 p.
- Kirkaldy, G.W. 1910: On some preoccupied generic names in insects. *Canadian entomologist* 42 : 8.
- Kitching, I.D. 1984: An historical review of the higher classification of the Noctuidae (Lepidoptera). *Bulletin of the British Museum (Natural History), entomology series* 49 (3) : 153-234.
- 1987: Spectacles and silver Ys: a synthesis of the systematics, cladistics and biology of the Plutinae (Lepidoptera: Noctuidae). *Bulletin of the British Museum (Natural History), entomology series* 54 (2) : 75-261.
- Klima, A. 1939: Pyralidae: subfamily Pyraustinae II. *Lepidopterorum Catalogus, pars* 94 : 225-384.
- Kloet, G.S.; Hincks, W.D. 1972: Handbooks for the identification of British insects. Vol. 11, A checklist of British insects. Part 2: Lepidoptera. London, Royal Entomological Society. 2nd edition, revised. 153 p.
- Kluk, K. 1802: Zwierząt domowych i dzikich osobliwie Kraiowych. Hist. nat. poczatki i gospodarstwo. 4 vols. [See Ackery, P.; Vane-Wright, R.I., 1984, p. 268; publication date may be in doubt].
- Knaggs, H.G. 1867: New species of *Scoparia* (Lepidoptera) from New Zealand, collected by R.W. Fereday. *Entomologists' monthly magazine* 4 : 80-81.
- Kollar, V. 1832: Systematisches Verzeichniss der Schmetterlinge im Erzherzogthume Oesterreich. *Beiträge Landesk. Oesterreich.* 2 (1).
- Kozhanchikov, I.V. 1956: Fauna of the USSR. Lepidoptera, Vol. III (2), Psychidae. 516 p.
- Kristensen, N.P. 1980: *Sphinx tipuliformis* Clerck, 1759 (Insecta, Lepidoptera): proposed conservation.

- Z.N.(S.)2139. *Bulletin of zoological nomenclature* 37 (3) : 154-156.
- 1984: Studies on the morphology and systematics of primitive Lepidoptera (Insecta). *Steenstrupia* 10 : 141-191.
- Kristensen, N.P.; Nielsen, E.S. 1979: A new subfamily of micropterigid moths from South America. A contribution to the morphology and phylogeny of the Micropterigidae, with a generic catalogue of the family (Lepidoptera: Zeugloptera). *Steenstrupia* 5 (7) : 69-147.
- Kudrna, O. 1974: *Artogeia* Verity, 1947, gen. rev. for *Papilio napi* Linnaeus (Lep., Pieridae). *Entomologist's gazette* 25 (1) : 9-12.
- Kumata, T. 1977: On the Japanese species of the genera *Macarostola*, *Aristaea*, and *Systoloneura*, with descriptions of three new species (Lepidoptera: Gracillariidae). *Insecta Matsumurana (new series)* 9 : 1-51.
- Kuznetsov, V.I.; Kerzhner, I.M. 1984: *Laspeyresia* Hübner, [1825] (Insecta, Lepidoptera): proposed conservation by the suppression of *Cydia* Hübner, [1825]. Z.N.(S.) 2421. *Bulletin of zoological nomenclature* 41 : 110-113.
- Kyrki, J. 1983: *Roeslerstamnia* Zeller assigned to Amphitheridae, with notes on the nomenclature and systematics of the family (Lepidoptera). *Entomologica Scandinavica* 14 : 321-329.
- 1984: The Yponomeutoidea - a reassessment of the superfamily and its constituent suprageneric taxa (Lepidoptera). *Entomologia Scandinavica* 15 : 71-84.
- Lange, W.H., Jr 1950: Biology and systematics of plume moths of the genus *Platyptilia* in California. *Hilgardia* 19 (19) : 561-668.
- 1956: A generic revision of the aquatic moths of North America: (Lepidoptera: Pyralidae, Nymphulinae). *Wasmann journal of biology (San Francisco)* 14 : 59-144.
- Latreille, P.A. 1796: Précis de caractères génériques des Insectes, disposés dans un ordre naturel par le Citoyen Latreille. Bordeaux, Brive. 210 p.
- Leach, W.E. 1815: Entomology. Pp. 57-172 in Brewster (ed.), Edinburgh Encyclopedia, vol. 9.
- Lederer, J. 1863: Beitrag zur kentniss der Pyralidinen. *Wiener entomologische Monatsschrift* 7 : 427-504.
- Lesse, H. de 1952: Nôte sur les genres *Precis* Hb. et *Junonia* Hb. (Lep. Nymphalidae). *Bulletin de la Société entomologique de France* 57 : 74-77.
- Lewin, J.W. 1805: Prodromus entomology. Natural history of lepidopterous insects of New South Wales. Collected, engraved and faithfully painted from Nature by John William Lewin, A.L.S. of Parramatta, New South Wales. London. 18 plates. [Names and descriptions by A. Macleay - Fletcher 1929, p. vii, as "W. Macleay".]
- Lienig, F.; Zeller, P.C. 1846: Lepidopterologische Fauna von Livland und Kurland mit Anmerkungen von Zeller. *Isis* 1846 (III) : 175-302.
- Linnaeus, C. 1758: *Systema Naturae*, 10. Holmiae, Laurentii Salvii. 824 p.
- 1761: Fauna Suecica. Holmiae, Laurentii Salvii. 560 p.
- 1767: *Systema Naturae*, 12, 1 (2). Holmiae, Laurentii Salvii. Pp. 533-1327.
- Longstaff, G.B. 1912: On the nomenclature of the Lepidoptera of New Zealand. *Transactions and proceedings of the New Zealand Institute* 44 : 108-115.
- Lower, O.B. 1897: Descriptions of new Australian Lepidoptera. *Transactions of the Royal Society of South Australia* 21 : 50-60.
- Lucas, T.P. 1892: On twenty new species of Australian Lepidoptera. *Proceedings of the Linnaean Society of New South Wales* (2) 7: 249-266.
- Macleay, W.S. 1826: Annulosa. Catalogue of insects collected by Captain King, R.N. Appendix B, pp. 438-469, tab. B in King, Capt. Phillip P., Narrative of a survey of the intertropical and western coasts of Australia performed between the years 1818 and 1822, vol. 2.
- McCoy, F. 1868: (Letter in) *Annals and Magazine of Natural History* (4) 1 : 76.
- McGuffin, W. 1967: Guide to the Geometridae of Canada (Lepidoptera). I. Subfamily Sterrhinae. *Memoirs of the Entomological Society of Canada* 50. 67 p.
- 1972: Guide to the Geometridae of Canada (Lepidoptera). II. Subfamily Ennominae. 1. *Memoirs of the Entomological Society of Canada* 86. 159 p.
- McQuillan, P.B.; Ireson, J.E. 1987: The identity and immature stages of *Patagoniodes farinaria* (Turner) comb. nov. (Lepidoptera: Pyralidae: Phycitinae) in Australia and New Zealand. *Journal of the Australian Entomological Society* 26 : 239-247.
- Maes, K. 1985: A comparative study of the abdominal tympanal organs in Pyralidae (Lepidoptera). I. Description, terminology, preparation technique. *Nota lepidopterologica* 8(4) : 341-350.
- Mann, J. 1857: Verzeichniss der in Jahre 1853 in der Gegend von Fiume gesammelten Schmetterlinge. Schluss. *Wiener entomologischer Monatschrift* 1 : 161-189.
- Marion, H. 1954: Contribution à l'étude des Pyralidae de Madagascar. *Mémoires de l'Institut Scientifique du Madagascar (E)* 5 : 39-62.
- Meyrick, E. 1879a: Descriptions of Australian Microlepidoptera. I. Crambites. *Proceedings of the Linnaean Society of New South Wales* 3 : 175-216.
- 1879b: Descriptions of Australian Micro-Lepidoptera. II. Crambites. *Proceedings of the Linnaean Society of New South Wales* 4 : 205-242.
- 1880a (22 November): Descriptions of Australian Microlepidoptera. III. Tineina. *Proceedings of the Linnaean Society of New South Wales* 5 : 132-182.
- 1880b: Descriptions of Australian Microlepidoptera. IV. Tineina (cont.). *Proceedings of the Linnaean Society of New South Wales* 5 : 204-271.
- 1881a: Descriptions of Australian Microlepidoptera. V. Tortricina. *Proceedings of the Linnaean Society of New South Wales* 6 : 410-536.

- 1881b: Descriptions of Australian Microlepidoptera. VI. Tortricina (continued). *Proceedings of the Linnaean Society of New South Wales* 6 : 629-706.
- 1882a: Descriptions of New Zealand Microlepidoptera. I. Abstract. *New Zealand journal of science (Dunedin)* 1 : 186-187.
- 1882b: Descriptions of Australian Microlepidoptera. VII. Revisional. *Proceedings of the Linnaean Society of New South Wales* 7 : 148-202.
- 1882c: Descriptions of New Zealand Microlepidoptera. II. Abstract. *New Zealand journal of science (Dunedin)* 1 : 277-278.
- 1883a: Descriptions of Australian Microlepidoptera. VIII. Oecophoridae. *Proceedings of the Linnaean Society of New South Wales* 7 : 415-547.
- 1883b: Descriptions of New Zealand Microlepidoptera. I and II. Crambidae and Tortricina. *Transactions and proceedings of the New Zealand Institute* 15 : 3-33, 33-68.
- 1883c: Descriptions of New Zealand Microlepidoptera. III. Oecophoridae [abstract]. *New Zealand journal of science* 1 : 522-525.
- 1883d: Monograph of New Zealand Geometrina [abstract]. *New Zealand journal of science (Dunedin)* 1 : 526-531.
- 1883e: Descriptions of Australian Microlepidoptera. IX. Oecophoridae (continued). *Proceedings of the Linnaean Society of New South Wales* 8 : 320-383.
- 1884a: Descriptions of New Zealand Microlepidoptera. III. Oecophoridae. *Transactions and proceedings of the New Zealand Institute* 16 : 3-49.
- 1884b: A monograph of the New Zealand Geometrina. *Transactions and proceedings of the New Zealand Institute* 16 : 49-113.
- 1884c: Supplement to a monograph of the New Zealand Geometridae. *New Zealand journal of science* 2 : 234-235.
- 1884d: Descriptions of New Zealand Microlepidoptera. IV. Scopariidae. *New Zealand journal of science* 2 : 235-237.
- 1884e: On the classification of the Australian Pyralidina. *Transactions of the entomological society of London*, 1884 : 277-350.
- 1885a: Description of New Zealand Microlepidoptera. V [VI]. Pyralidina [abstract]. *New Zealand journal of science (Dunedin)* 2 : 346-348.
- 1885b: Tortricina (supplementary). Additions to former paper. Abstract. *New Zealand journal of science (Dunedin)* 2 : 348-349.
- 1885c: Descriptions of Australian Microlepidoptera. XII. Oecophoridae (continued). *Proceedings of the Linnaean Society of New South Wales* 9 : 1045-1082.
- 1885d: Supplement to a monograph of the New Zealand Geometrina. *Transactions and proceedings of the New Zealand Institute* 17 : 62-68.
- 1885e: Descriptions of New Zealand Microlepidoptera. IV. Scopariidae. *Transactions and proceedings of the New Zealand Institute* 17 : 68-120.
- 1885f: Descriptions of New Zealand Microlepidoptera. VI. Pyralidina. *Transactions and proceedings of the New Zealand Institute* 17 : 121-140.
- 1885g: Descriptions of New Zealand Microlepidoptera. VII. Tortricina (supplementary). *Transactions and proceedings of the New Zealand Institute* 17 : 141-149.
- 1885h: Notes on the nomenclature of the New Zealand Geometrina. *New Zealand journal of science* 2 : 589.
- 1885i: Descriptions of New Zealand Microlepidoptera. Gelechiidae. VIII. Tineina (part). *New Zealand journal of science* 2 : 589-592.
- 1885j: On the classification of the Australian Pyralidina. *Transactions of the Entomological Society of London*, 1885 : 421-456.
- 1886a: Descriptions of Australian Microlepidoptera. XII. Oecophoridae (continued). *Proceedings of the Linnaean Society of New South Wales* 10 : 765-832.
- 1886b: Description of New Zealand Microlepidoptera. VIII. Tineina (Part). *Transactions and proceedings of the New Zealand Institute* 18 : 163-183.
- 1886c: Notes on nomenclature of New Zealand Geometrina. *Transactions and proceedings of the New Zealand Institute* 18 : 184.
- 1886d: On the synonymy of some species of Nyctemera. *Entomologists' monthly magazine* 23 : 15-16.
- 1886e: Revision of Australian Lepidoptera. *Proceedings of the Linnaean Society of New South Wales, new series* 1 : 687-802.
- 1886f: Notes on synonymy of Australian Mircolepidoptera [sic]. *Proceedings of the Linnaean Society of New South Wales, new series* 1 : 803-806.
- 1886g: Descriptions of Lepidoptera from the South Pacific. *Transactions of the Entomological Society of London*, 1886 : 189-296.
- 1887: Monograph of the New Zealand Noctuina. *Transactions and proceedings of the New Zealand Institute* 19 : 3-40.
- 1888a: Supplement to a monograph of New Zealand Noctuina. *Transactions and proceedings of the New Zealand Institute* 20 : 44-47.
- 1888b: Notes on New Zealand Geometrina. *Transactions and proceedings of the New Zealand Institute* 20 : 47-62.
- 1888c: Notes on New Zealand Pyralidina. *Transactions and proceedings of the New Zealand Institute* 20 : 62-73.
- 1888d: Notes on New Zealand Tortricina. *Transactions and proceedings of the New Zealand Institute* 20 : 73-76.
- 1888e: Descriptions of New Zealand Tineina. *Transactions and proceedings of the New Zealand Institute* 20 : 77-106.
- 1889a: Descriptions of Australian Microlepidoptera. XV. Oecophoridae. *Proceedings of the*

- Linnaean Society of New South Wales* (2) 3 : 1565-1703.
- 1889b: Descriptions of New Zealand Microlepidoptera. *Transactions and proceedings of the New Zealand Institute* 21 : 154-188.
- 1890: Descriptions of New Zealand Lepidoptera. *Transactions of the New Zealand Institute* 22 : 204-220.
- 1891: New species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 23 : 97-101.
- 1892: On new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 24 : 216-220.
- 1893: Descriptions of Australian Microlepidoptera. XVI. Tineidae. *Proceedings of the Linnaean Society of New South Wales* (series 2) 7 : 477-612.
- 1897a: Descriptions of Australian Microlepidoptera. XVII. Elachistidae. *Proceedings of the Linnaean Society of New South Wales* 22 : 297-435.
- 1897b: Descriptions of new Lepidoptera from Australia and New Zealand. *Transactions of the entomological society of London*, 1897 : 367-390.
- 1901: Descriptions of new Lepidoptera from New Zealand. *Transactions of the entomological society of London*, 1901 : 565-579.
- 1902a: Descriptions of new Australian Lepidoptera. *Transactions of the Entomological Society of London*, 1902 : 25-48.
- 1902b: A new genus of Gelechiidae. *Entomologists' monthly magazine* 38 : 103-104.
- 1902c: Lepidoptera from the Chatham Islands. *Transactions of the Entomological Society of London*, 1902 : 273-279.
- 1902d: Descriptions of new species of Lepidoptera (Oecophoridae). *Transactions of the Royal Society of South Australia* 26 (2) : 133-174.
- 1904: Descriptions of Australian Microlepidoptera. XVIII. Gelechiidae. *Transactions and proceedings of the Linnaean Society of New South Wales* 29 : 255-441.
- 1905: Notes on New Zealand Lepidoptera. *Transactions of the Entomological Society of London*, 1905 : 219-244.
- 1906: Descriptions of Australian Tineina. *Transactions and proceedings of the Royal Society of South Australia* 30 : 33-66.
- 1907a: Descriptions of Indian Microlepidoptera. *Journal of the Bombay Natural History Society* 17 : 730-754.
- 1907b: Descriptions of Australasian Microlepidoptera. XIX. Plutellidae. *Proceedings of the Linnaean Society of New South Wales* 32 : 47-150.
- 1907c: Notes and descriptions of Lepidoptera. *Transactions and proceeding of the New Zealand Institute* 39 : 106-121.
- 1909a: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 41 : 5-16.
- 1909b: Lepidoptera from Auckland Island. Pp. 70-74 in Chilton, C. (ed.), *Subantarctic islands of New Zealand*, vol. 1. Wellington.
- 1910a: Notes on New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 42 : 63-66.
- 1910b: Lepidoptera from the Kermadec Islands. *Transactions and proceedings of the New Zealand Institute* 42 : 67-73.
- 1910c: Revision of Australian Tortricina. *Proceedings of the Linnaean Society of New South Wales* 35 : 139-294.
- 1910d: Lepidoptera Heterocera (Pyrales). Fam. Pterophoridae. *Genera insectorum fasc.* 100. 22 p.
- 1911a: A new *Crambus* from New Zealand. *Entomologists' monthly magazine* 47 : 82-83.
- 1911b: Notes and descriptions of New Zealand Lepidoptera. Parts I and II. *Transactions and proceedings of the New Zealand Institute* 43 : 57-78.
- 1911c: A revision of the classification of the New Zealand Tortricina. *Transactions and proceedings of the New Zealand Institute* 43 : 78-91.
- 1911d: Revision of Australian Tortricina. *Proceedings of the Linnaean Society of New South Wales* 36 : 224-303.
- 1911e: Descriptions of Indian Microlepidoptera. *Journal of the Bombay Natural History Society* 20 : 706-736.
- 1912a: Adelidae, Micropterygidae, Gracilaridae. In Wagner, H. (ed.), *Lepidopterorum catalogus*, pars 6. 68 p.
- 1912b: A revision of the classification of the New Zealand Caradrinina. *Transactions and proceedings of the New Zealand Institute* 44 : 88-107.
- 1912c: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 44 : 117-126.
- 1912d: *Exotic Microlepidoptera* 1 (2) : 33-64. London, Taylor & Francis.
- 1912e: Tortricidae. In Wagner, H. (ed.), *Lepidopterorum catalogus*, pars 10. 86 p.
- 1912f (precise date unknown): Lepidoptera Heterocera: family Micropterygidae. *Genera insectorum, fasc.* 132. 9 p.
- 1913a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 45 : 22-29.
- 1913b: A revision of New Zealand Pyralidina. *Transactions and proceedings of the New Zealand Institute* 45 : 30-51.
- 1913c: Carposinidae, Heliodinidae, Glyphipterygidae. *Lepidopterorum catalogus*, pars 13. 53 p.
- 1913d: Pterophoridae, Orneodidae. *Lepidopterorum catalogus*, pars 17. 44 p.
- 1913e: Lepidoptera Heterocera. Fam. Tortricidae. *Genera insectorum, fasc.* 149. 81 p.
- 1914a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 46 : 101-118.

- 1914b: *Exotic Microlepidoptera* 1 : 241–256.
- 1915a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 47 : 201–204.
- 1915b: Revision of New Zealand Tineina. *Transactions and proceedings of the New Zealand Institute* 47 : 205–244.
- 1915c: *Exotic Microlepidoptera* 1 : 289–320.
- 1916a: Occurrence of a New Zealand Tineid in Britain. *The entomologists' monthly magazine* 52 : 17–18.
- 1916b: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 48 : 414–419.
- 1917a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 49 : 245–247.
- 1917b: Revision of New Zealand Notodontina. *Transactions and proceedings of the New Zealand Institute* 49 : 248–273.
- 1918a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 50 : 132–134.
- 1918b: *Exotic Microlepidoptera* 2 : 177–208.
- 1919: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 51 : 349–354.
- 1920a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 52 : 30–32.
- 1920b: *Exotic Microlepidoptera* 2 : 321–352.
- 1921: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 53 : 334–336.
- 1922a: *Exotic Microlepidoptera* 2 : 513–544.
- 1922b: A new genus and species of Microlepidoptera from New Zealand. *The Entomologist* 55 : 270–271.
- 1922c: Lepidoptera Heterocera (Tineae). Fam. Heliodinidae. *Genera insectorum*, fasc. 165. 29 p.
- 1922d: Lepidoptera Heterocera. Fam. Oecophoridae. *Genera insectorum*, fasc. 180. 224 p.
- 1923: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 54 : 162–169.
- 1924a: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 55 : 202–206.
- 1924b: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 55 : 661–662.
- 1924c: *Exotic Microlepidoptera* 3 : 65–96.
- 1925a: Lepidoptera of the Chatham Islands. *Records of the Canterbury Museum* 2 : 269–275.
- 1925b: Lepidoptera Heterocera, fam. Gelechiidae. *Genera insectorum*, fasc. 184. 290 p.
- 1926a: *Exotic Microlepidoptera* 3 : 257–288.
- 1926b: Descriptions of New Zealand Lepido-  
ptera. *Transactions and proceedings of the New Zealand Institute* 56 : 415–416.
- 1927a: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 57 : 697–702.
- 1927b: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 58 : 313–316.
- 1929: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 60 : 483–490.
- 1931a: New species of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 62 : 92–97.
- 1931b: Notes on New Zealand Lepidoptera. *Records of the Canterbury Museum* 3 : 367–369.
- 1932: Descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 63 : 24.
- 1934: Notes on New Zealand Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 64 : 151–153.
- 1935: Descriptions of two new species of New Zealand Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 65 : 304.
- 1936: Descriptions and notes on New Zealand Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 66 : 281–283.
- 1937: *Exotic Microlepidoptera* 5 (4) : 97–128.
- 1938: New species of New Zealand Lepidoptera. *Transactions and proceedings of the Royal Society of New Zealand* 67 : 426–429.
- Miller, L.D. 1968: The higher classification, phylogeny and zoogeography of the Satyridae (Lepidoptera). *Memoirs of the American Entomological Society* no. 24. 174 p.
- Miller, L.D.; Brown, F.M. 1979: Studies in the Lycaeninae (Lycaenidae). 4. The higher classification of the American coppers. *Bulletin of the Allyn Museum* 51. 30 p.
- Miller, W.E. 1985: (Comments on the proposed conservation of *Laspeyresia* Hübner, [1825]. Z.N.(S.)2421.) *Bulletin of zoological nomenclature* 42 : 8–9.
- Minet, J. 1981: Les Pyraloidea et leurs principales divisions systématiques. *Bulletin de la Société entomologique de France* 86 : 262–280.
- 1983: Etude morphologique et phylogénétique des organes tympaniques des Pyraloidea. 1. Généralités et homologies (Lep. Glossata). *Annales de la Société Entomologique de France (new series)* 19 : 175–207.
- 1984: Contribution à l'analyse phylogénétique des Néolépidoptères (Lepidoptera: Glossata). *Nouvelle revue d'entomologie* 1 (2) : 139–149.
- 1985: Etude morphologique et phylogénétique des organes tympaniques des Pyraloidea. 2 - Pyralidae; Crambidae, première partie (Lepidoptera Glossata). *Annales de la Société Entomologique de France* 21 : 69–86.

- 1986: Ebauche d'une classification moderne de l'ordre des Lépidoptères. *Alexanor* 14(7) : 291–313.
- Moore, F. 1880: Lepidoptera of Ceylon 1 (1). London. 40 p.
- 1881: Descriptions of new genera and species of Asiatic nocturnal Lepidoptera. *Proceedings of the Zoological Society of London*, 1881 : 326–380.
- [1885]: Lepidoptera of Ceylon 3. London. Pp. 1–304.
- 1886: The Lepidoptera of Ceylon 3. London. Pp. 305–392, pl. clxxi–cxcv.
- Moriuti, S. 1977: Fauna Japonica: Yponomeutidae s. lat. (Insecta: Lepidoptera). Tokyo, Keigaku Publishing Co. 326 p., xcvi pl.
- Morrison, B. 1968: A further record of *Dryadaula pacotlia* Meyrick (Lep., Tineidae) in Britain with notes on its life history. *Entomologist's gazette* 19 : 181–183.
- Morrison, H.K. 1874a: V. Descriptions of new Noctuidae. *Bulletin of the Buffalo Society for Natural Science* 2 : 109–117.
- 1874b: Description of new Noctuidae. *Proceedings of the Boston Society for Natural History* 17 : 131–166.
- Müller-Rutz, J. 1920: Aus der Welt der Kleinschmetterlinge. *Mitteilungen der Entomologie, Zurich und Umgebung*??, 5 : 334–349.
- Munroe, E. 1964: Insects of Campbell Island. Lepidoptera: Pyralidae. *Pacific Insects Monograph* 7 : 260–271.
- 1972: The moths of America north of Mexico. Part 13 1A. Pyraloidea, Pyralidae, comprising subfamilies Scopariinae, Nymphulinae. London, E.W. Classey Ltd & R.B.D. Publications Inc. 134 p.
- Murray, R.P. 1873: Descriptions of new species of exotic Rhopalocera. *Entomologists' monthly magazine* 10 (1873–74) : 107–108.
- Newman, E. 1856: Characters of a few Australian Lepidoptera, collected by Mr Thomas R. Oxley. *Transactions of the Entomological Society of London (n.s.) iii* 8 : 281–300.
- Nielsen, E.S. 1982: Review of the higher classification of the Lepidoptera, with special reference to the lower heteroneurans. *Tyō to Ga* 33 (1,2) : 98–101.
- 1985a: Primitive (non-ditrysian) Lepidoptera of the Andes; diversity, distribution, biology, and phylogenetic relationships. *Journal of research on the Lepidoptera, supplement 1* : 1–16.
- 1985b: The monotrysian heteroneuran phylogeny puzzle: a possible solution (Lepidoptera). *Proceedings of the 3rd Congress of European Lepidopterology, Cambridge, 1982* : 138–143.
- Nielsen, E.S.; Common, I.F.B. (in press): Lepidoptera (moths and butterflies). In *The insects of Australia* (2nd edn). CSIRO/Melbourne University Press.
- Nielsen, E.S.; Davis, D.R. 1985: The first southern hemisphere prodoxid and the phylogeny of the Incurvarioidea (Lepidoptera). *Systematic entomology* 10 : 307–322.
- Nowicki, M.S. 1864: Microlepidopterorum species novae, auctore ... Accedit una tabella. Cracoviae, sumptibus V. Dzieduszycki. 31 p.
- Nye, I.W.B. 1975: The generic names of moths of the world. Volume 1: Noctuoidea (part): Noctuidae, Agaristidae and Nolidae. *Publications of the British Museum (Natural History)* 770. 568 p.
- 1982: Nomenclature of *Heliothis* and associated taxa (Lepidoptera: Noctuidae): past and present. In ICRISAT (International Crop Research Institute for the Semi-arid Tropics) 1982, Proceedings of the International Workshop on *Heliothis* Management, 15–20 November 1981, ICRISAT Center, Patancheru A.P., India.
- Obraztsov, N.S. 1942: *Djakonovia* gen. nov., eine neue Tortricinen-Gattung (Lep. Tortricidae). *Iris (Deutsche entomologische Zeitung, Berlin)* 56 : 157–158. [Note: Obraztsov refers to this as 1942; *Zoological Record* gives "(1942) 1943".]
- 1949: Materialien zu einer Revision der palaearktischen Tortricinae-Gattungen (Lep.: Tortricidae). 1. Allgemeine Aufteilung der unterfamilie Tortricinae und die Tribe Ceracini. *Entomon (Munich)* 1 : 200–206.
- 1955: Die gattungen der Palaearktischen Tortricidae. 1. Allgemeine Aufteilung der Familie und die unterfamilien Tortricinae und Sparganothinae. *Tijdschrift voor Entomologie* 98 (3) : 147–228.
- Ochsenheimer, F. 1816: Die Schmetterlinge von Europa. 4. Nachträge zu Bande 1–3. 223 p.
- Olivier, A.G. 1789: Encyclopédie méthodique. Dictionnaire des Insectes, vol. 4. Paris, Pankouke. 331 p.
- Owada, M. 1987: A taxonomic study on the subfamily Herminiinae of Japan (Lepidoptera: Noctuidae). Tokyo, National Science Museum. 208 p.
- Pactl, J. 1953: Genera of the Hepialidae (Insecta, Lepidoptera). *Journal of the Asiatic Society, Calcutta, (science)* 19 : 141–148.
- Patzak, H. 1974: Beiträge zur Insectenfauna der DDR: Lepidoptera – Coleophoridae. *Beiträge zur Entomologie* 24 : 153–278.
- Petersen, G. 1957: Die Genitalien der paläarktischen Tineiden (Lepidoptera: Tineidae). *Beiträge zur Entomologie* 7 : 55–176, 338–379, 557–595.
- 1969: Beiträge zur Insekten-Fauna der DDR: Lepidoptera: Tineidae. *Beiträge zur Entomologie* 19 : 311–388.
- Philpott, A. 1903: On some new species of Lepidoptera (moths) from Southland. *Transactions and proceedings of the New Zealand Institute* 35 : 246–249.
- 1905: On some new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 37 : 328–331.
- 1910: Description of a new species of moth. *Transactions and proceedings of the New Zealand Institute* 42 : 544.
- 1912: Descriptions of three new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 44 : 115–116.

- 1913: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 45 : 76–78.
- 1914: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 46 : 118–121.
- 1915: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 47 : 192–201.
- 1916: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 48 : 420–423.
- 1917a: A list of the Lepidoptera of Otago. *Transactions and proceedings of the New Zealand Institute* 49 : 195–238.
- 1917b: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 49 : 239–245.
- 1918: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 50 : 125–132.
- 1919: Descriptions of new species of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 51 : 224–225.
- 1920: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 52 : 42–44.
- 1921: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 53 : 337–342.
- 1922: The venation of the genus *Mnesarchaea* (Lepidoptera) with a description of a new species. *New Zealand journal of science and technology* 5 : 80–82.
- 1923: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 54 : 148–154.
- 1924a: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 55 : 207–214.
- 1924b: The tibial strigil of the Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 55 : 215–224.
- 1924c: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 55 : 663–669.
- 1925: Australian Lepidoptera in New Zealand. Occurrence in Nelson during 1924–25. *New Zealand journal of science and technology* 7 : 364–366.
- 1926a: New Zealand Lepidoptera: notes and descriptions. *Transactions and proceedings of the New Zealand Institute* 56 : 387–399.
- 1926b: List of New Zealand species of *Borkhausenia* (Oecophoridae: Lepidoptera), including new species. *Transactions and proceedings of the New Zealand Institute* 56 : 399–413.
- 1926c: A new endemic genus of Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 56 : 414.
- 1927a: N.Z. Lepidoptera: notes and descriptions. *Transactions and proceedings of the New Zealand Institute* 57 : 703–709.
- 1927b: The genitalia of the genus *Gymnobathra* (Oecophoridae: Lepidoptera). *Transactions and proceedings of the New Zealand Institute* 57 : 716–721.
- 1927c: The male genitalia of the Hepialidae. *Transactions of the Entomological Society of London* 75 : 35–41.
- 1927d: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 58 : 80–92.
- 1927e: The male genitalia of the New Zealand Tineidae. *Transactions and proceedings of the New Zealand Institute* 58 : 93–101.
- 1927f: The male genitalia of the New Zealand Oecophoridae. *Transactions and proceedings of the New Zealand Institute* 58 : 102–113.
- 1927g: The male genitalia of the New Zealand Plutellidae. *Transactions and Proceedings of the New Zealand Institute* 58 : 317–326.
- 1927h: The male genitalia of the New Zealand Lyonetiidae. *Transactions and proceedings of the New Zealand Institute* 58 : 327–336.
- 1927i: The male genitalia of the New Zealand Glyphipterygidae. *Transactions and proceedings of the New Zealand Institute* 58 : 337–347.
- 1927j: The male genitalia of the New Zealand Gelechiidae. *Transactions and proceedings of the New Zealand Institute* 58 : 348–356.
- 1928a: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 58 : 359–370.
- 1928b: Notes on *Isonomeutis amauropa* Meyr. (Lepidoptera). *Transactions and proceedings of the New Zealand Institute* 58 : 371–374.
- 1928c: Some new species of Lepidoptera. *Records of the Canterbury Museum [N.Z.]* 3 : 181–183.
- 1928d: The male genitalia of the New Zealand Tortricidae. *Transactions and proceedings of the New Zealand Institute* 59 : 443–468.
- 1928e: The male genitalia of the New Zealand Eucosmidae. *Transactions and proceedings of the New Zealand Institute* 59 : 469–475.
- 1928f: The male genitalia of the New Zealand Carposinidae. *Transactions and proceedings of the New Zealand Institute* 59 : 476–480.
- 1928g: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 59 : 481–490.
- 1928h: The male genitalia of the New Zealand Pterophoridae. *Transactions and proceedings of the New Zealand Institute* 59 : 645–649.
- 1929a: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 60 : 300–304.
- 1929b: The male genitalia of the New Zealand Crambidae. *Transactions and proceedings of the New Zealand Institute* 60 : 491–514.

- 1930a (March): Descriptions of Lepidoptera in the Canterbury Museum. *Records of the Canterbury Museum [N.Z.]* 3 : 247–250.
- 1930b (June): New species of Lepidoptera in the collection of the Auckland Museum. *Records of the Auckland Institute and Museum* 1 : 1–16.
- 1930c (August): The Lepidoptera of Mount Cook District, with descriptions of new species. *Transactions and proceedings of the New Zealand Institute* 61 : 419–439.
- 1931: Notes and descriptions of New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 62 : 26–36.
- Pierce, F.N.; Metcalfe, J.W. 1938: The genitalia of the Pyrales with the Deltoids and Plumes. An account of the morphology of the male clasping organs and the corresponding organs of the female. Warmington, F.N. Pierce. 69 p., 30 pl.
- Poole, R.W. 1970: Transfer of four New Zealand geometrids from the genus *Azelina* (Lepidoptera: Geometridae). *Proceedings of the Entomological Society of Washington* 72 : 135.
- Popescu-Gorj, A.; Căpușe, I. 1965: Revision d'*Oegoconia quadripuncta* (Hw.) (Lepidoptera Gelechioidea) des collections de Roumanie. *Revue Roumaine de biologie zoologie (Academia Republicii Socialiste România, Bucharest)* 6 : 389.
- Povolny, D. 1967: Genitalia of some Nearctic and Neotropical members of the tribe Gnorimoschemini (Lepidoptera: Gelechiidae). *Acta entomologica Musei nationalis Pragae* 37 : 51–127.
- 1974: Revision of the genus *Empista* Povolny (*Zeempista* subgen. n.) from New Zealand (Lepidoptera: Gelechiidae). *Acta entomologica Bohemoslovaca* 71 : 414–428.
- 1977: Notes on Gnorimoschemini of Australia and New Zealand (Lepidoptera: Gelechiidae). *Acta entomologica Musei nationalis Pragae* 39 : 403–443.
- Prout, L.B. 1910: Lepidoptera Heterocera fam. Geometridae subfam. Oenochrominae. *Genera insectorum, fasc. 104.* 120 p.
- 1912: Notes on the nomenclature of the New Zealand Geometridae, with descriptions of a new species. *Proceedings of the New Zealand Institute* 44 : 52–54. [Note: although usually bound together, the *Transactions* and the *Proceedings* pages are separately numbered.]
- 1914: New Lepidoptera (Geometridae). *Transactions and proceedings of the New Zealand Institute* 46 : 122–123.
- 1920: Geometrae II: Fauna Indo-Australica. In Seitz, A., *The Macrolepidoptera of the World (Stuttgart)* 12 (English edition) : 1–116.
- 1927: Notes on New Zealand Geometridae. *Transactions and proceedings of the New Zealand Institute* 58 : 75–79.
- 1934: Geometridae: Fauna Indo-Australica. In Seitz, A., *The Macrolepidoptera of the World (Stuttgart)* 12 (English edition) : 133–140.
- 1939: Geometridae: Fauna Indo-Australica. In Seitz, A., *The Macrolepidoptera of the World (Stuttgart)* 12 : 237–292.
- 1958: New species of Indo-Australian Geometridae. *Bulletin of the British Museum (Natural History)* 6 (12) : 367–463.
- Purdie, A.A. 1884: *Orocrambus* sp. *New Zealand journal of science (Dunedin)* 2 : 167–168.
- 1887: Description of a new species of moth (*Pasiphila lichenodes*). *Transactions and proceedings of the New Zealand Institute* 19 : 69–72.
- Quail, A. 1901: On *Lysiphragma howesii*, sp. nov. *Transactions and proceedings of the New Zealand Institute* 33 : 154–158.
- 1903: On *Charagia virescens* Dbld. *Transactions and proceedings of the New Zealand Institute* 35 : 249–255.
- Ragonot, E.L. 1888: Nouveaux genres et espèces de Phycitidae et Galleridae. Paris. 52 p.
- 1894: Description d'une nouvelle espèce de Tineite (*Trichophaga coprobiella*) provenant d'Obrock (mer Rouge). *Annales de la Société Entomologique de France* 62 : 120–124.
- Richards, A.G. Jr 1933: Comparative skeletal morphology of the noctuid tympanum. *Entomologica Americana* 13 : 1–43.
- Riedl, T. 1969: Matériaux pour la connaissance des Momphidae paléarctiques (Lepidoptera). Partie 9. Revue des Momphidae européennes, y compris quelques espèces d'Afrique du Nord et du Proche-Orient. *Polskie pismo entomologiczne* 39 : 635–919.
- Riley, C.V. 1869: First annual report on the noxious, beneficial and other insects of the State of Missouri, p. 180, pl. II (fig. 13 and 14), fig. 98.
- Robinson, G.S. 1976: The preparation of slides of Lepidoptera genitalia with special reference to the Microlepidoptera. *Entomologists' gazette* 27 : 127–132.
- 1979: Clothes moths of the *Tinea pellionella* complex: a revision of the world's species (Lepidoptera: Tineidae). *Bulletin of the British Museum (Natural History) (entomology)* 38 (3) : 57–128.
- 1983: Darwin's moth from St Paul's Rocks: a new species of *Erechthias* (Tineidae). *Systematic entomology* 8 : 303–311.
- Robinson, G.S.; Nielsen, E.S. 1983: The Microlepidoptera described by Linnaeus and Clerck. *Systematic entomology* 8 : 191–242.
- Roesler, U. 1966: Die deutschen Arten des *Homoeosoma - Ephestia*-Komplexes (Lepidoptera, Phycitidae). *Mitteilungen der München Entomologischen Gesellschaft* 56 : 104–160.
- 1969: Phycitinen-Studien VII (Lepidoptera: Pyralidae). *Entomologische Zeitung, Frankfurt-am-Main*, 79 : 245–260.
- Rosenstock, R. 1885: Notes on Australian Lepidoptera, with descriptions of new species. *Annals and magazine of natural history (5)* 16 : 377–384.
- Rothschild, L.W. 1896: Some undescribed Lepidoptera. *Novitates Zoologicae* 3 : 231–232.
- 1914: Indo-Australian Arctiidae. In Seitz, A., *The Macrolepidoptera of the World: Fauna Indo-*

- Australica* 10: 236–263.
- Rungs C. 1953: Le complex de *Leucania loreyi* auct. nec Dup. (Lep. Phalaenidae). *Bulletin de la Société Entomologique de France* (1953) 9: 138–141.
- Salmon, J.T. 1946: New Lepidoptera from the Homer – Milford District. *Dominion Museum (New Zealand) records in entomology* 1: 1–11.
- 1948: New species and records of Lepidoptera from the Three Kings Islands, New Zealand. *Records of the Auckland Institute and Museum* 3: 309–311.
- 1956: New species of New Zealand Lepidoptera. *Transactions of the Royal Society of New Zealand* 83: 573–576.
- Salmon, J.T.; Bradley, J.D. 1956: Lepidoptera from the Cape Expedition and Antipodes Islands. *Records of the Dominion Museum, Wellington*, 3: 61–81.
- Samouelle, G. 1819: Entomologists' useful compendium; or an introduction to the knowledge of British insects etc. London. 496 p., 12 pl.
- Sattler, K. 1973: A catalogue of the family-group and genus-group names of the Gelechiidae, Holcogonidae, Lecithoceridae and Symmocidae (Lepidoptera). *Bulletin of the British Museum (Natural History), entomology*, 28 (4): 155–282.
- 1978: The identity of the genus *Athrips* Billberg, 1820 (Lep., Gelechiidae). *Deutsche entomologische Zeitschrift* 25: 57–61.
- 1988: The systematic status of the genera *Ilseopsis* Povolny, 1965 and *Empista* Povolny, 1968 (Lepidoptera: Gelechiidae: Gnornimoschemini). *Nota lepidopterologica* 10: 224–235.
- Schrank, F. von P. 1801: Fauna Boica 2 (1). Nürnberg, Ingolstadt, Landshut. 374 p.
- 1802: Fauna Boica 2 (2). Nürnberg, Ingolstadt, Landshut. 412 p.
- Scoble, M. 1986: The structure and affinities of the Hedyoidea: a new concept of the butterflies. *Bulletin of the British Museum (Natural History), entomology* 53(5): 251–286.
- Scopoli, J.A. 1763: *Entomologia Carniolica exhibens insecta Carnioliae indigena et distributa in ordines, genera, species, varietates, methodo Linneana. Vindobonae, Trattner.* 36 + 420 p.
- Scott, A.W. 1859: On a new lepidopterous insect from Australia. *Proceedings of the Zoological Society of London*, XXVII, pt 2, 396: 207–209.
- 1864: Australian Lepidoptera and their transformations, drawn from the life, by Harriet and Helena Scott; with descriptions, general and systematical by A.W. Scott, M.A. London. 30 p., ix pl.
- Scott, J.A. 1986: On the monophyly of the Macrolepidoptera, including a reassessment of their relationship to the Coccoidea and Castnoidea, and a reassignment of Mimallonidae to Pyraloidea. *Journal of research on the Lepidoptera* 25: 30–38.
- Scudder, S.H. 1875: Historical sketch of the generic names proposed for butterflies. A contribution to systematic nomenclature. *Proceedings of the American Academy of Arts and Science* 10: 91–293.
- Sibatini, A. 1974: A new genus for two new species of Lycaeninae (s. str.) (Lepidoptera: Lycaenidae) from Papua New Guinea. *Journal of the Australian Entomological Society* 13 (2): 95–110.
- Snellen, P.C.T. 1880: Nieuwe Pyraliden op het eiland Celebes gevonden door Mr M.C. Piepers. *Tijdschrift voor Entomologie* 23: 198–250.
- Stainton, H.T. 1849: An attempt at a systematic catalogue of the British Tineidae and Pterophoridae. London, van Voorst. 32 p.
- 1851: A supplementary catalogue of the British Tineidae and Pterophoridae. London, van Voorst. 28 p.
- 1854: *Insecta Britannica, Tineina*. London, Lovell Reeve. 313 p.
- 1859: A new *Tinea*. *The entomologist's weekly intelligencer* 6: 183.
- Standinger, O. 1879: Ueber Lepidopteren des südöstlichen europäischen Russlands. *Stettiner entomologischer Zeitung* 40: 315–328.
- Stempffer, H. 1967: The genera of African Lycaenidae (Lepidoptera: Rhopalocera). *Bulletin of the British Museum (Natural History) (entomology)*, suppl. 10. 322 p.
- Stephens, J.F. 1834: Illustrations of British entomology ... Haustellata, 4. London, Baldwin & Craddock. 436 p.
- Stephens, J.F. 1850: List of the specimens of British animals in the collection of the British Museum. Part 5, Lepidoptera. 353 p.
- Stoll, C. 1790: Supplementband zu Cramer's 'Papillons exotiques'. Amersterdam, Graevius. Lief. 4.
- Sunley, R.M. 1911: Notes on the larvae of some New Zealand Lepidoptera. *Transactions and proceedings of the New Zealand Institute* 43: 129–130.
- Swinhoe, C. 1902: New species of eastern and Australian Heterocera. *Annals and magazine of natural history* (7) 9: 415–424.
- 1916: New Indo-Malayan Lepidoptera. *Annals and magazine of natural history* (8) 18: 209–221.
- Taylor, R. 1843: Te Ika a Maui, or New Zealand and its inhabitants ... London, Wertheim & MacIntosh.
- Thierry-Meig, P. 1915: Descriptions de Lépidoptères nouveaux et nôtes lépidoptérologiques. *Miscellanea Entomologica, Narbonne*, 22: 61–66.
- Tilden, J.W. 1971: Comments on the nearctic members of the genus *Precis* Huebner. *Journal of research on the Lepidoptera* 9: 101–108.
- Tillyard, R.J. 1931: Mr Alfred Philpott [obituary]. *Transactions and proceedings of the New Zealand Institute* 62: 172–174.
- Todd, E.L. 1978: A checklist of species of *Heliothis* Ochsenheimer (Lepidoptera: Noctuidae). *Proceedings of the Entomological Society of Washington* 80: 1–14.
- Traugott-Olsen, E.; Nielsen, E.S. 1977: The Elachistidae (Lepidoptera) of Fennoscandia and Denmark. *Fauna entomologica Scandinavica* 6. 299 p.

- Treitschke, F. 1833: Die Schmetterlinge von Europa, 9 (2). Leipzig. Fleischer. 294 p.
- Turner, A.J. 1896: Descriptions of microlepidoptera from Queensland. *Transactions of the Royal Society of South Australia* 20 : 1-34.
- 1904: A preliminary revision of the Australian Thyrididae and Pyralidae. Part 1. *Proceedings of the Royal Society of Queensland* 18 : 109-199.
- 1913: Studies in Australian Microlepidoptera (Plutellidae). *Proceedings of the Linnaean Society of New South Wales* 38 : 174-228.
- 1917: Studies in Australian Lepidoptera. *Transactions of the Royal Society of South Australia* 41 : 57-120.
- 1919: The Australian Gelechianae (Lepidoptera). *Proceedings of the Royal Society of Queensland* 31 : 108-172.
- 1926: Studies in Australian Lepidoptera. *Transactions of the Royal Society of South Australia* 50 : 120-155.
- 1927: New and little-known Tasmanian Lepidoptera. Part II. *Papers and proceedings of the Royal Society of Tasmania*, 1926 : 119-164.
- Tutt, J.W. 1899: A natural history of the British Lepidoptera. A text book for students and collectors, 1. London. 560 p.
- Vari, L. 1961: South African Lepidoptera. I: Lithocolletidae. *Transvaal Museum memoir* 12. 238 p.
- 1971: Lepidoptera (Heterocera: Tineidae, Hyponomeutidae). Pp. 349-354 in van Zinderen Bakker, E.M.; Winterbottom, J.M.; Dyer, R.A. (eds), Marion and Prince Edward Islands. Report on the South African biological and geological expedition, 1965-66. Capetown, A.A. Balkema.
- Verity, R. 1947: Le farfalle diurne d'Italia. Volume terzo. Divisione Papilionida. Sezione Papilionina (Famiglie Papilionidae e Pieridae). Florence, Marzocco. 318 p.
- Viette, P.E.L. 1950: Contribution à l'étude des Hepialidae (22ème note). Hepialidae du Musée de Leiden. *Zoologische Mededelingen* 31 : 67-77.
- 1951: Sur quelques noctuelles décrites par Guénée (1852-1854). *Bulletin mensuel de la Société Linnaéenne de Lyon* 20 : 159-162.
- 1954: Une nouvelle espèce de Lépidoptère brachyptère de l'île Campbell. *Entomologiske Meddelelser, Copenhagen*, 27 : 19-22.
- 1961: Notes on some synonyms or preoccupied names in the Lepidoptera. *The entomologist, London*, 94 : 38-39.
- Wagner, H.; Pfitzner, R. 1911: Hepialidae. Pp. 1-26 in Aurivillius, C.; Wagner, H. (eds), Lepidopterorum catalogus, pars 4.
- Walker, F. 1854: List of the specimens of lepidopterous insects in the collection of the British Museum. II. Pp. 279-581.
- 1855: List of the specimens of lepidopterous insects in the collection of the British Museum. IV. Pp. 777-976.
- 1856a: List of the specimens of lepidopterous insects in the collection of the British Museum. VII. Pp. 1509-1808.
- 1856b: List of the specimens of lepidopterous insects in the collection of the British Museum. VIII: Sphingidae. Pp. 1-271.
- 1856c: List of the specimens of lepidopterous insects in the collection of the British Museum. IX: Noctuidae. Pp. 1-252.
- 1857a: List of the specimens of lepidopterous insects in the collection of the British Museum. X: Noctuidae. Pp. 253-491.
- 1857b: List of the specimens of lepidopterous insects in the collection of the British Museum. XI: Noctuidae. Pp. 493-764.
- 1858a: List of the specimens of lepidopterous insects in the collection of the British Museum. XII: Noctuidae. Pp. 765-982.
- 1858b: List of the specimens of lepidopterous insects in the collection of the British Museum. XIII: Noctuidae. Pp. 983-1236.
- 1858c: List of the specimens of lepidopterous insects in the collection of the British Museum. XV: Noctuidae. Pp. 1521-1888.
- 1859a: List of the specimens of lepidopterous insects in the collection of the British Museum. XVII: Pyralides. Pp. 255-508.
- 1859b: List of the specimens of lepidopterous insects in the collection of the British Museum. XVIII: Pyralides. Pp. 509-798.
- 1859c: List of the specimens of lepidopterous insects in the collection of the British Museum. XIX: Pyralides. Pp. 799-1036.
- 1860a: List of the specimens of lepidopterous insects in the collection of the British Museum. XX: Geometrites. Pp. 1-276.
- 1860b: List of the specimens of lepidopterous insects in the collection of the British Museum. XXI: Geometrites (continued). Pp. 277-498.
- 1861: List of the specimens of lepidopterous insects in the collection of the British Museum. XXIII: Geometrites (continued). Pp. 757-1020.
- 1862a: List of the specimens of lepidopterous insects in the collection of the British Museum. XXIV: Geometrites (continued). Pp. 1021-1280.
- 1862b: List of the specimens of lepidopterous insects in the collection of the British Museum. XV: Geometrites (continued). Pp. 1281-1477.
- 1863a: List of the specimens of lepidopterous insects in the collection of the British Museum. XXVI: Geometrites (continued). Pp. 1479-1796.
- 1863b: List of the specimens of lepidopterous insects in the collection of the British Museum. XXVII: Crambites and Tortricites. Pp. 1-286.
- 1863c: List of the specimens of lepidopterous insects in the collection of the British Museum. XXVIII: Tortricites and Tineites. Pp. 287-561.
- 1864a: List of the specimens of lepidopterous insects in the collection of the British Museum. XXIX: Tineites. Pp. 563-835.

- 1864b: List of the specimens of lepidopterous insects in the collection of the British Museum. XXX: *Tineites*. Pp. 837–1096.
- 1865a: List of the specimens of lepidopterous insects in the collection of the British Museum. XXXII: supplement, part 2. Pp. 323–706.
- 1865b: List of the specimens of lepidopterous insects in the collection of the British Museum. XXXIII: supplement, part 3. Pp. 707–1120.
- 1866a: List of the specimens of lepidopterous insects in the collection of the British Museum. XXXIV: supplement, part 4. Pp. 1121–1533.
- 1866b: List of the specimens of lepidopterous insects in the collection of the British Museum. XXXV: supplement, part 5. Pp. 1535–2040.
- 1869: Characters of undescribed Lepidoptera Heterocera. London. 112 p.
- Wallengren, H.D.J. 1860: Lepidopterologische Mittheilungen. *Wiener entomologische Monatschritte* 4 : 161–176.
- 1880: Skandinaviens Arter at Tineidgruppen Plutellidae (Staint.). *Entomologisk Tidskrift* 1 : 53–63.
- Walsingham, F. (Rt Hon. Lord) 1897: Western equatorial African Micro-Lepidoptera. *Transactions of the Entomological Society of London*, 1897 : 33–67.
- 1900: Asiatic Tortricidae. *Annals and magazine of natural history* 5 : 451–469.
- 1907: Microlepidoptera of Tenerife. *Proceedings of the Zoological Society of London*, 1907 : 910–1026.
- Warren, B.C.S. 1967: Anatomical notes on the Satyrid genera *Percnodaimon* Butler and *Erebiola* Fereday (= *Dubierebia* Muschamp). In Wise, K.A.J., Taxonomy of two New Zealand butterfly species (Nymphalidae: Satyrinae). *Transactions of the Royal Society of New Zealand* 9 (4) : 39–41.
- Warren, W. 1892: Descriptions of new genera and species of Pyralidae, contained in the British Museum Collection. *Annals and magazine of natural history* (6) ix : 172–179.
- 1893: On new genera and species of moths of the family Geometridae from India, in the collection of H.J. Elwes, with notes by H.J. Elwes. *Proceedings of the Zoological Society of London*, 1893 : 341–434.
- 1894: New species and genera of Geometridae. *Novitates zoologicae* 1 : 366–466.
- 1895: New species and genera of Geometridae in the Tring Museum. *Novitates zoologicae* 2 : 82–159.
- 1896a: New genera and species of Pyralidae, Thyrididae and Epiplemidiae. *Annals and magazine of natural history* (6) xvii : 202–216.
- 1896b: New species of Drepanulidae, Thyrididae, Uraniidae, Epiplemidiae, and Geometridae in the Tring Museum. *Novitates zoologicae* 3 : 335–419.
- 1897a: New genera and species of moths from the old-world regions in the Tring Museum. *Novitates zoologicae* 4 : 12–130.
- 1897b: New genera and species of Drepanulidae, Thyrididae, Epiplemidiae, Uraniidae and Geometridae in the Tring Museum. *Novitates zoologicae* 4 : 195–262.
- 1903: New Drepanulidae, Thyrididae, Uraniidae and Geometridae from the Oriental Region. *Novitates zoologicae* 10 : 255–270.
- 1912 (25 September): Noctuidae. Pp. 65–96 in Seitz, A., Gross-Schmetterlinge der Erde. Fauna Indo-australica. 11. Stuttgart.
- Waterhouse, G.A.; Lyell, G. 1914: The butterflies of Australia. Sydney. 239 p.
- Watson, A.; Fletcher, D.S.; Nye, I.W.B. 1980: Noctuoidea (part). In Nye, I.W.B. (ed.), *The generic names of moths of the world*. 2. xiv + 228 p.
- Watt, J.C. 1979: Abbreviations for entomological collections. *New Zealand journal of zoology* 6 : 519–520.
- Watt, M.N. 1916: Description of a new species of *Melanochra* from Mount Egmont. *Transactions and proceedings of the New Zealand Institute* 48 : 413, with footnote.
- 1920: The leafmining insects of New Zealand. *Transactions and proceedings of the New Zealand Institute* 52 : 439–466.
- 1921a (August): The leafmining insects of New Zealand: part II. *Transactions and proceedings of the New Zealand Institute* 53 : 197–219.
- 1921b (November): A new tortricid moth. *New Zealand journal of science and technology* 4 : 257–258.
- 1924: The leafmining insects of New Zealand: part 5 – the genus *Nepticula* (Lepidoptera), and the Agromyzidae (Diptera), continued and *Gracilaria selenitis* Meyr. (Lepidoptera). *Transactions and proceedings of the New Zealand Institute* 55 : 674–687.
- Westwood, J.O. 1840: Synopsis of the British insects. In An introduction to the modern classification of insects, founded on the natural habitats and corresponding organisation of the different families. Vol. 2. London, Longman. 158 p.
- Whalley, P.E.S. 1971: The Thyrididae (Lepidoptera) of Africa and its islands. A taxonomic and zoogeographic study. *Bulletin of the British Museum (Natural History), entomology, supplement* 17. 198 p.
- 1973: The genus *Etiella* Zeller (Lepidoptera: Pyralidae): a zoogeographic and taxonomic study. *Bulletin of the British Museum (Natural History), entomology*, 28 (1) : 3–21.
- White, A. 1841: Notes on some insects from King George's Sound, collected and presented to the British Museum by Captain George Grey. Pp. 450–482 in Grey, G., *Journals of two expeditions of discovery in north-west and western Australia during the years 1837–39* [etc.], vol. 2, app. F.
- 1855: Pp. 418–425 in Taylor, R., *Te Ika a Maui, or New Zealand and its inhabitants* ... London, Wertheim & MacIntosh.
- 1862: [*Lycaena boldenarum* n.sp.]. *Proceedings of the Entomological Society of London*, 1861 : 26.

- Wilkinson, C. 1978: On the *Stigmella* - *Nepticula* controversy (Lepidoptera). *Tijdschrift voor Entomologie* 121 : 13-22.
- Wise, K.A.J. 1957: A new species of *Lithocolletis* (Lepidoptera: Gracillariidae) from New Zealand. *Proceedings of the Royal Entomological Society of London, B* 26 : 26-28.
- 1962: *Parectopa leucocyma* (Meyrick) (Lepidoptera: Gracillariidae) rediscovered as a leafminer of kauri (*Agathis australis* Salisb.). *Transactions of the Royal Society of New Zealand (zoology)* 1 : 373-375.
- 1967: Taxonomy of two New Zealand butterfly species (Nymphalidae: Satyrinae). *Transactions of the Royal Society of New Zealand (zoology)* 9 (4) : 39, 41-44.
- Wootton, R.J. 1979: Function, homology, and terminology in insect wings. *Systematic entomology* 4 : 81-93.
- Zeller, P.C.** 1839: Versuch einer naturgemässen Eintheilung der Schaben, *Tinea. Isis (Leipzig)*, 1839 : 167-220.
- 1846: Die Arten der Blattminirergattung *Lithocolletis* beschrieben. *Linnaea entomologica* 1 : 166-261.
- 1847: Bemerkungen über die auf einer Reise nach Italien und Sicilien gesammelten Schmetterlingsarten. *Isis (Leipzig)*, 1847 : viii + 561-549; xii + 881-914.
- 1848: Exotische Phyciden. *Isis (Leipzig)* : 857-890.
- 1849: Beitrag zur Kenntniss der Coleophoren. *Linnaea entomologica* 4 : 191-416.
- 1852: Lepidoptera Microptera quae J.A. Wahlberg in Caffrorum terra collegit. *Kongliga Svenska Vetenskaps-Akademiens Handligar*. 120 p.
- 1853: Drei javanische Nachfalter beschrieben. *Bulletin de la Société des Naturalistes de Moscou* 24 (3) : 502-516. [Note: (1) Horn & Schenkling (1928, p. 1361) give volume no. as 26. (2) Also known as *Byulleten' Moskovskogo obshchestva ispytatelei prirody*.]
- 1863: Chilonidarum et Crambidarum genera et species. Meseritz, also Berlin, Wiegandt & Hempe. 56 p.
- 1873: Beiträge zur Kenntniss der nordamerikanischen Nachfalter, besonders der Microlepidopteren, 2. *Verhandlung der Zoologisch-botanischen Gesellschaft in Wien* 23 : 201-334.
- 1875: Beiträge zur Kenntniss der nordamerikanischen Nachfalter, besonders der Microlepidopteren, 3. *Verhandlungen der Zoologisch-botanischen Gesellschaft in Wien* 25 : 207-360.
- 1877: Exotische Microlepidoptera. *Horae Societatis entomologicae rossicae* 13 : 3-493.
- 1879: Lepidopterologische Bemerkungen. *Stettiner entomologische Zeitung* 40 : 462-473.
- Zimmerman, E.C. 1958: Macrolepidoptera. *Insects of Hawaii* 7. 542 p.
- 1971: The New Zealand moth genus *Irenicodes* (Lepidoptera: Cycnodidae). *The New Zealand entomologist* 5 : 53-55.
- 1978: Microlepidoptera, part 1: Monotrysia, Tineoidea, Tortricoidea, Gracillarioidea, Yponomeutoidea and Alucitoidea. *Insects of Hawaii* 9. 881 p.

— 8 —

## APPENDIX

### Lepidoptera recorded from the Kermadec Islands (29°S, 177°W)

\*<sup>†</sup>, Raoul Island type locality  
†, recorded once only

#### Arctiidae

† *Nyctemera a. annulata* (Boisduval). Meyrick 1910  
*Uetetheisa pulchelloides vaga* Jordan. Meyrick 1910.  
NZAC, NMNZ

#### Blastobasiidae

*Blastobasis* sp. NZAC

#### Cosmopterigidae

*Cosmopterix attenuatella* (Walker). NZAC  
*Pyroderces aellotricha* (Meyrick). Meyrick 1910 (in  
*Sathrobrota*). NZAC  
*Pyroderces anarithma* Meyrick. NZAC

#### Geometridae

*Chloroclystis testulata* (Guenée). Meyrick 1910 (as  
*Phriessogonus denotatus*, *Chloroclystis indicata*). NZAC, NMNZ  
† *Epicyme rubropunctaria* (Doubleday). Meyrick 1910

\**Epyaxa lucidata* (Walker). Meyrick 1910, p. 69 (as  
*Hydriomena officiosa* Meyrick, new species).  
NZAC

*Phriessogonus laticostatus* (Walker). Meyrick 1910.  
NZAC  
*Scopula rubraria* (Doubleday). Meyrick 1910.  
NZAC

#### Glyptapterigidae

† \**Glyptapterix scolias* Meyrick, 1910, p. 73. HT  
unique, ?lost

#### Gracillariidae

† *Caloptilia octopunctata* (Turner). Meyrick 1910

#### Lycaenidae

*Zizina labradus labradus* (Godart). Meyrick 1910

#### Noctuidae

##### Acronyctinae

*Callopistria mailliardi* Guenée. Meyrick 1910 (in  
*Eriopus*). NZAC

*Leucocosmia nonagraria* (Walker). Meyrick 1910  
(as *Caradrina reclusa* Walker). NZAC

*Platysenta illecta* Walker. Meyrick 1910 (as *Per-*  
*igea capensis* Guenée). NZAC

*Spodoptera litura* (Fabricius). Meyrick 1910 (as  
*Spodoptera littoralis* Boisduval). NZAC

##### Hadeninae

† *Graphania insignis* (Walker). NMNZ (repre-  
sented by a single specimen)

*Mythimna loreyimima* (Rungs). Meyrick 1910 (as  
*Leucania loreyi* Duponchel). NZAC

*Mythimna separata* (Walker). Meyrick 1910 (as  
*Leucania unipuncta* Haworth). NZAC

*Tiracola plagiata* Walker. Meyrick 1910. NZAC

#### Heliothinae

*Heliothis armigera conferta* (Walker). Meyrick  
1910. NZAC

#### Hypeninae

\**Hydrillodes surata* Meyrick, 1910, p. 68. NZAC

#### Hypenodinae

† *Hypenodes gonospilalis* Walker. Meyrick 1910  
*Schrankia costaestrigalis* Stephens. Meyrick 1910.  
NZAC

#### Noctuinae

*Agrotis epsilon aneitura* (Walker). Meyrick 1910.  
NZAC

*Diasria intermixta* Guenée. NZAC

#### Catocalinae s.l.

† *Achaea janata* (Linnaeus). Meyrick 1910 (as  
*Achaea melicerete* Drury)

*Anomis flava* (Fabricius). NZAC

† *Anticarsia irrorata* Walker. Meyrick 1910 (as  
*Thermesia rubricans* Boisduval)

† *Hypocala deflorata australasiae* Butler. NZAC  
*Mocis frugalis* Walker. NZAC

† *Mocis trifasciata* Walker. NZAC

#### Plusiinae

*Chrysodeixis eriosoma* (Doubleday). Meyrick 1910  
(as *Plusia chalcites* Esper). NZAC

#### Nymphalidae

*Bassaris itea* Fabricius. Meyrick 1910. NZAC

*Danaus plexippus* Linnaeus. NZAC

*Hypolimnas bolina nerina* Felder. Meyrick 1910  
(as *Anosia bolina* Linnaeus). NZAC

† *Melanitis leda* Linnaeus. Meyrick 1910

† *Tiramula hamata* Macleay. NZAC

#### Oecophoridae

*Hofmannophila pseudospretella* (Stainton). NZAC

\**Pachyrhabda antinoma* Meyrick, 1910, pp. 72-73.  
NZAC

#### Pterophoridae

† *Sphenarches caffer* Zeller. Meyrick 1910

#### Pyralidae (in the broad sense)

† "Botys sp.". NZAC

*Cryptoblabes gnidiella* Millière. NZAC

† *Diasemia ramburialis* Duponchel. Meyrick 1910  
*Diplopseustis perieralis* (Walker). NZAC

† \**Dracaenura aegialitis* Meyrick, 1910, p. 71  
*Endotricha mesenterialis* in the sense of Meyrick.  
Meyrick 1910. NZAC

\**Eranistis pandora* Meyrick, 1910, p. 70. NZAC

*Herpetogramma licarsialis* (Walker). NZAC

*Homoeosoma anaspila* Meyrick. Meyrick 1910.  
NZAC

*Hymenia recurvalis* Fabricius. Meyrick 1910 (as

*Hymenia fascialis* Cramer). NZAC

\**Scoparia fragosa* Meyrick, 1910, p. 71. NZAC

#### Sphingidae

*Agrius convolvuli* (Linnaeus). NZAC

#### Tineidae

*Chloropleca terpsichorella* (Busck). NZAC

*Erechthias flavostriata* Walsingham. Meyrick 1910.

NZAC

*Monopis* sp. nr *ptilophaga* Enderlein. Meyrick

1910 (as *Monopis etherella* Newman). NZAC

*Opogona aurisquamosa* Butler. Meyrick 1910.

NZAC

*Opogona omoscopa* (Meyrick). NZAC

#### Tortricidae

*Bactra noteraula* Walsingham. Meyrick 1910.

NZAC

*Capua semiferana* complex. Meyrick 1910. NZAC

*Crocidosema plebejana* Zeller. NZAC

\**Polychrosis meliscia* Meyrick, 1910, p. 72

*Sperchia intractana* Walker. NZAC

\**Strepsicrates melanotreta* (Meyrick). Meyrick 1910,

p. 72 (as *Spilonota*). NZAC

#### Yponomeutidae

*Rhigognostis sera* Meyrick. NZAC

*Plutella xylostella* Linnaeus. Meyrick 1910. NZAC

*Prays* sp., *citri* group. NZAC

*Prays* sp., "brown speckled". NZAC

**Note.** The status of type material described from Raoul Island by Meyrick (1910) has not been elucidated. No specimens of *Glyptipterix scolias* were found, either in BMNH or NMNZ. Specimens of some species in NMNZ have red circular labels, but these could have been put on by someone other than Meyrick. Some specimens have labels seemingly in Meyrick's writing, but most are apparently in Hudson's hand. There is no doubt concerning the identity of any of the Kermadec species described by Meyrick (1910) in *Trans. N.Z. Inst.* 42, pp. 67-73.

#### Statistical summary

Total no. of species	69
No. of 'resident' species	50-55
No. of species recorded once only	16
No. not seen since 1908	10
No. recorded by Meyrick (1910)	45
No. not recorded in 1908	24
Immigration/extinction rate (%)	23

(suggesting a typical immigrant fauna; cf. 2-3% for 'mainland' New Zealand, i.e., an order of magnitude less)

## TAXONOMIC INDEX

Every name in the Taxonomic Catalogue is indexed. Junior synonyms are in a lighter type than currently valid names. Where a species-group name is listed with more than one genus, its status in each combination may be inferred from the type weight used for the generic name. In the case of synonyms, the combinations of generic and specific name listed are those originally published by authors, and may differ from combinations implicit in current usage.

- A**
- abdita*, *Epichorista* (auct. 2) 122
  - abditus*, *Orocrambus* 146
  - abjectana*, ?*Teras* 121
  - abnegatana*, *Grapholitha* 119
  - abrogata*, *Asaphodes* 171
  - absconditaria*, *Acidalia* 186
  - abstittella*, *Simaethis* 113
  - abyssina*, *Mallobathra* 67
  - accensa*, *Teras* 129
  - acceptella*, *Oxythecta* 97
  - aceptrix*, *Mamestra* 204
  - accurata*, *Aletia* 201
  - accusatrix*, *Tinea* sensu Meyrick 65
  - acerella*, *Gadira* 144
  - acetina*, *Agrotis* 212
  - Achaea* 196
  - acharis*, *Scoparia* 159
  - achlyoessa*, *Glyphipteryx* 76
  - Achroia* 141
  - achroiaria*, *Amilapis?* 192
  - achrosta*, *Harmologa* auct. 122
  - achyrota*, *Anisoplaca* 80
  - acidaliaria*, *Fidonia?* 193
  - Acipitilia* 132
  - Acleris* 129
  - acmonias*, *Izatha* 96
  - acmotypa*, *Thectophila* 86
  - acompa*, *Antiscopa* 154
  - acompsa*, *Pasiphila* 186
  - acontistis*, *Tmetolophota* 209
  - acridomima*, *Thylacosceles* 109
  - acrocausta*, *Cacoecia* 126
  - Acrocercops* 70
  - Acroclita* 114
  - acrodactyla*, *Anisoplaca* 80
  - acrodina*, *Erechthias* 61
  - acronoma*, *Glyphipteryx* 76
  - ACRONICTINAE** 195
  - acronyctoides*, *Spodoptera mauritia* 196
  - acrothecta*, *Glyphipteryx* 76
  - acroxantha*, *Parocystola* 98
  - Actias* 140
  - actinias*, *Tingena* 99
  - acuta*, *Mnesarchaea* 55
  - acutata*, *Samana* 192
  - adapertella*, *Gelechia* 95
  - Adeixis** 191
  - adelina*, *Graciliaria* 71
  - ademptella*, *Oecophora* 100
  - Adena* 152
  - admirationis*, *Euxoa* 213
  - admotella*, *Tinea* 119
  - adonata*, *Cidaria* 181
  - adonis*, *Asaphodes* 172
  - adreptella*, *Heterocrossa* 130, 131
  - aduncalis*, *Musotima* 151
  - adusta*, *Selidosema* 167
  - adversa*, *Mnesictena* 153
  - aegalis*, *Gauna* 153
  - aegrota*, *Asaphodes* 172
  - aegusalis*, *Gauna* 143
  - aellomacha*, *Acrocercops* 70
  - aellotricha*, *Pyroderces* auct. 86
  - aelodes*, *Platyptilia* 133
  - aemula*, *Sabatinca* 52
  - aenea*, *Gelophaula* 123  
    *Glyphipteryx* 56  
    *Sabatinca* 52
  - Aenetus* 56
  - aerata*, *Tinea* 65
  - aerifera*, *Glyphipteryx* 76
  - aerobatis*, *Kiwaia* 81
  - aerodana*, *Ericodesma* 117
  - aethaliana*, *Hendecasticha* 115
  - aethalota*, *Acrocercops* 70
  - aetherea*, *Tinea* sensu Meyrick 65
  - aethonellus*, *Orocrambus* 146
  - affinis*, *Tingena* 99
  - affinalis*, *Loxostege* 153
  - afflcta*, *Tingena* 99
  - agana*, *Scoparia* 155
  - AGARISTINAE** 196
  - agaura*, *Batrachedra* 84
  - aggregata*, *Cidaria* 183
  - agilitata*, *Sorensonata* 128
  - Aglossa* 143
  - agorastis*, *Graphania* 202
  - agrionata*, *Tatosoma* 190
  - agriopa*, *Crypsitricha* 60
  - Agriophara* 109
  - Agrius* 140
  - Agrotis* 212
  - alba*, *Helastia* 180
  - albafasciata*, *Pseudocoremia* 167
  - albafascicula*, *Scoparia* auct. 160
  - albalineata*, *Asaphodes* 172
  - albifasciata*, *Asterivora* 113
  - albilineata*, *Asaphodes* 172
  - albimaculata*, *Stathmopoda* 108
  - albiplaga*, *Chloroclystis* 176, 177

- albostriata*, *Ctenoplusia* 213  
*alcione*, *Meterana* 206  
*alcyonipenella*, *Ornix* 85  
*alectoraria*, *Xyridacma* 192  
*Aletia* 199  
*aletis*, *Philobota* 99  
    *Tingena* 99  
*allogama*, *Epichorista* (auct. 1) 122  
*alopa*, *Tmetolophota* 209  
*alopecana*, *Catamacta* 119  
*alopecias*, *Eudonia* 154  
*alta*, *Tatosoma* 190  
*alterna*, *Mocis* 198  
*Alysia* 201  
*alysidota*, *Acrocercops* 70  
*Alysina* 201  
*Alytopistis* n. gen. nr 69  
*amalodes*, *Carposina* 130  
*Amastris* 169  
*amauropa*, *Isonomeutis* 129  
*ambigua*, *Gymnobathra* 92  
*amblyterma*, *Homodotis* 182  
*amenana*, *Leptocroca* 96  
*amica*, *Nyctemera* 194  
*amiculata*, *Tingena* 99  
*amnopis*, *Borkhausenia* 100  
*amorbas*, *Izatha* 94  
*amotella*, *Oecophora* 98  
*amphileuca*, *Trachypepla* 106  
*Amphitape* 164  
*Amphixystis* 59  
*ampla*, *Pseudocoremia rufisata* 169  
*amplexana*, *Harmologa* 124  
*Anachloris* 170  
*anaema*, *Tingena* 99  
*analogia*, *Asterivora* 113  
*anarithma*, *Pyroderces* auct. 86  
*Anarsia* sensu Meyrick 80  
*anartoides*, *Teia* 195  
*anaspila*, *Homoeosoma* 142  
*anastrella*, *Trachypepla* 106  
*Anatossa* 164  
*anceps*, *Dasyuris anceps* 177  
*ancogramma*, *Tingena* 99  
*Andesia* 198  
*aneitura*, *Agrotis epsilon* 212  
*angularis*, *Trachypepla* 106  
*anguligera*, *Austrocidaria* 175  
*angusta*, *Helastia* 180  
    *Mallobathra* 68  
    *Mamestra* 203  
*Angustalius* 143  
*angustipennis*, *Orocrambus* 146  
*animosa*, *Scoparia* auct. 160  
*Anisoplaca* 80  
*annulata*, *Nyctemera* 194  
    *Porina* 57  
*annulatella*, *Caunaca* 76  
*anomalella*, *Stigmella* 53  
*Anomis* 197  
*antarctica*, *Chloroclystis* 187  
*Antherea* 140  
*anthracias*, *Aponotoreas* 171  
*Antichloris* 195  
*antichroa*, *Tortrix* auct. 128  
*anticlina*, *Trigonistis* 211  
*antigrapha*, *Asterivora* 113  
*antimorus*, *Crambus* 146  
*antiphona*, *Plutella* 74  
*antipoda*, *Epialus* 58  
    *Mamestra* 209  
*antipodaria*, *Epirranthis* 163  
*antipodea*, *Mnesictena* 153  
    *Nyctemera* 194  
*Antipodesma* 64  
*antipodium*, *Argyrophenga* 138  
*antiquana*, *Prothelymna* 128  
*Antiscopa* 154  
*antitypa*, *Harmologa* 125  
*Aochleta* 94  
*Aoria* 56  
*aoristana*, *Capua* 119  
*apanthes*, *Tingena* 99  
*Apatetris* 81  
*apertella*, *Tingena* 99  
*apeholes*, *Scoparia* 159  
*aphelias*, *Asaphodes* 172  
*aphrias*, *Epiblemma* 116  
*aphrontis*, *Tingena* 99  
*aphrosticha*, *Mallobathra* 68  
*apicata*, *Larentia* 174  
*apicellus*, *Orocrambus* 146  
*apicipallida*, *Tatosoma* 190  
*apocrypta*, *Phaeosaces* 98  
*apodoxa*, *Izatha* 94  
*Aponotoreas* 170  
*aposema*, *Stathmopoda* 108  
*apparitella*, *Pyroderces* auct. 86  
*apselias*, *Crambus* 149  
*aqilonaris*, *Leptocroca* 96  
*aquosata*, *Cidaria* 188  
*arachnias*, *Mamestra* 203  
*araneosa*, *Reductoderces* 69  
*archaeonoma*, *Cosmiotes* 111  
*Archipini* 117, 118  
*Archyala* 59  
*Arkte* 197  
*Arctesthes* 171  
**ARCTIIDAE** 194  
*arcuata*, *Notoreas* (Meyrick) 184  
    *Pyrgotis* 127  
*ardularia*, *Coremia* 180  
*arenacea*, *Hyperythra* 164  
*arenosa*, *Austrocidaria* 175

- arenosella*, *Batrachedra* 84  
*argentaria*, *Aletia* 199  
*Lysiphragma* auct. 214  
*Selidosema* 167  
*argentata*, *Pyrameis* 137  
*argentifera*, *Chrysodeixis* 213  
*argentosa*, *Ericodesma* 117  
*argillacea*, *Monopis* 63  
*argodelta*, *Tinea* sensu Meyrick 65  
*Argua* 164  
*Argyrophenga* 138  
*Argyroploce* 114  
*arida*, *Hydriomena* auct. 182  
*aridella*, *Gelophaula* 123  
*aristarcha*, *Chalastra* 163  
*aristias*, *Pasiphila* 187  
*aristodoxa*, *Stathmopoda* 108  
*Aristotelia* 81  
*armigera conferta*, *Heliothis* (*Helicoverpa*) 211  
*armigerella*, *Tingena* 100  
*arotis*, *Tmetolophota* 209  
*Arsinoe* 170  
*Artigisa* 197  
*asaleuta*, *Scoparia* auct. 160  
*Asaphodes* 171  
*ascendens*, *Porina* 57  
*Ascerodes* 118  
*ascomorpha*, *Tortrix* 126  
*ascotata*, *Cidaria* 180  
*aspera*, *Ectopatria* 212  
*aspersana*, *Acleris* 129  
*aspaltis*, *Leptocroca* 96  
*aspidephora*, *Trachypepla* 106  
*aspidotata*, *Eudonia* 155  
*aspistana*, *Epichorista* 122  
*assata*, *Cidaria* 182  
*asterisca*, *Eudonia* 155  
*Asterivora* 113  
*asteronota*, *Glyphipterix* 78  
*asterope*, *Meterana* 206  
*Asterophaga* 113  
*Astheniodes* 189  
*astraea*, *Tinea* sensu Meyrick 65  
*astragalota*, *Scoparia* auct. 160  
*astrapaea*, *Glyphipterix* 77  
*astrapia*, *Boarmia* 169  
*astricta*, *Batrachedra* 84  
*Astrogenes* 60  
*astrologana*, *Cacoecia* 125  
*astrosema*, *Diptychophora* 145  
*ataracta*, *Glyphipterix* 77  
*Athetis* 195  
*Atrips* auct. 80  
*atmogramma*, *Eudonia* 155  
*Notoreas* (Meyrick) 184  
*Tinea* sensu Meyrick 65  
*Atomotricha* 90  
*Atossa* 164  
*atra*, *Heliothela* 152  
*atristriga*, *Tmetolophota* 209  
*atronivea*, *Declana* 164  
*attactella*, *Izatha* 94  
*attenuata*, *Campbellana* 130  
*attracta*, *Boarmia* 165  
*attributa*, *Acidalia* 193  
*atychioides*, *Hierodoris* 89  
*auklandiae*, *Tinearupa sorenseni* 99  
*auklandica*, *Reductoderces* 69  
*augastis*, *Scoparia* 159  
*aulacias*, *Leucania* 209  
*aulistes*, *Crambus* 146  
*aulogramma*, *Glyphipterix* 77  
*Aulopola* 179  
*aurantiaca*, *Sabatinca* 52  
*aurata*, *Tingena* 100  
*aurella*, *Micropardalis* 52  
*Nepticula* 53  
*aurimaculata*, *Aoraia* 56  
*auriscriptella*, *Glaucoccharis* 144  
*austera*, *Izatha* 94  
*australiae*, *Hypocala deflorata* 197  
*Austramathes* 198  
*austrina*, *Dasyuris* 178  
*Oxythecta* 97  
*Austrocidiaria* 174  
*Austrotortrix* 117  
*autocharis*, *Epiphryne* 179  
*autochroa*, *Scoparia* 159  
*autolitha*, *Nesolocha* 162  
*autumna*, *Scoparia* auct. 160  
*autumnata*, *Porina* 57  
*averilla*, *Graphania* 202  
*aversa*, *Persectania* 208  
*axena*, *Eudonia* 155  
*axenana*, *Epalxiphora* 122  
*azaleella*, *Caloptilia* 71  
*Azelina* 166
- B**
- Bactra* 115  
*bactrias*, *Glyphipterix* 77  
*badia*, *Meterana* 206  
*balanophora*, *Izatha* 94  
*bankia*, *Melanitis leda* 138  
*barbarica*, *Helioscibes* auct. 89  
*Sabatinca* 52  
*barbata*, *Glyphipterix* 77  
*barbigera*, *Asterivora* 113  
*Barea* 90  
*Barsine* 163  
*Bascantis* 60  
*basella*, *Tingena* 100  
*basifascia*, *Persectania* 210  
*Bassaris* 136

*batis, Thyatira* 193  
*Batrachedra* 84  
**BATRACHEDRIDAE** 84  
*beata, Asaphodes* 172  
*Graphania* 202  
*Bedellia* 79  
*behrii, Etiella* 142  
*bellatula, Borkhausenia* 99  
*belonota, Tinea* sensu Meyrick 65  
*benedicta, Xanthorhoe* 172  
*berenice, Tingena* 100  
*berylia, Pseudocoremia* 167  
*bicomma, Nitocris* 196  
*bifaciella, Compsistis* 91  
*Tingena* 99, 100  
*bifasciella, Elachista* 111  
*biguttana, Teras* 125  
*bilineata, Hierodoris* 89  
*bilineolata, Pasiphila* 187  
*Biloba* 80  
*bipartita, Austrocidaria* 175  
*bipunctella, Glaucocharis* 144  
*bisectellus, Crambus* 150  
*bisignata, Cidaria* 183  
*bisignella, Tinea* 61  
*bisinalis, Eudonia* 155  
*bisselliella, Tineola* 66  
*bistonaria, Selidosema* 168  
*Bityla* 195  
*bjerkandrella, Tebenna* 114  
**BLASTOBASIIDAE** 85  
**BLASTOBASIINAE** 85  
*Blastobasis* 85  
*blax, Notoreas* (Meyrick) 184  
*blenheimensis, Tmetolophota* 209  
*boeticus, Lampides* 135  
*boisduvali, Xyleutes* 112  
*boldenarum, Lycaena boldenarum* 135  
*boldensis, Melanchra* 206  
*bolina nerina, Hypolimnas* 137  
**BOMBYCIDAE** 140  
**BOMBYCOIDEA** 140  
*Bombyx* 140  
*Boocara* 106  
*boreophilaria, Hybernia* 170  
*Borkhausenia* auct. 91, 99  
*Brachmia* sensu Meyrick 81  
*brachyacma, Tingena* 100  
*brachydelta, Glyphipterix* 77  
*Brachysara* 90  
*bradleyi, Tebenna* 114  
**BRASSOLINAE** 136  
*bephos, Fidonia* 186  
*bephosata, Paranotoreas* 186  
*brevicula, Harmologa* 124  
*bromias, Graphania* 202  
*brontophora, Kiwaia* 81

*brunneosa, Graphania* 202  
*bryaula, Gymnobathra* 92  
*bryopis, Larentia* 173  
*bulbulata, Xanthorhoe* auct. 190  
*butleri, Erebola* 138  
*byrsopola, Proterodesma* 64  
  
**C**  
*Cacopsodes* 191  
*Cadmogenes* 73  
*caerulea, Phyesetica* 209  
*caerulea, Chrysophanus* 136  
*Kiwaia* 81  
*Lycaena boldenarum* 135  
*caesia, Scoparia* auct. 160  
*calamogonus, Megacraspedus* 83  
*calaspidea, Kiwaia* 81  
*Calicotis* 109  
*calida, Helastia* 187  
*caliginosa, Gymnobathra* 92  
*Scoparia* auct. 160  
*calliactis, Glyptapterix* 77  
*calliarcha, Sabatinca* 52  
*callichlora, Austrocidaria* 175  
*callicrena, Dasyuris* 178  
*calligypsa, Pyrgotis* 127  
*calliphana, Zaprystra* 87  
*calliploca, Gymnobathra* 92  
*callirhous, Crambus* 146  
*callirrhous, Orocrambus* 146  
*callispura, Hierodoris* 89  
*callista, Declana* 164  
*callixyla, Gymnobathra* 92  
*Caloptilia* 71  
*calybe, Junonia villida* 137  
*camelias, Asaphodes* 172  
*camelina, Adoxophyes* 120  
*caminora, Stathmopoda* 108  
*Campbellana* 130  
*campbellensis, Asaphodes* 172  
*Eudonia* 158  
*campbelli, Pseudocoremia* 167  
*campbellica, Mallobathra* 68  
*campsiptera, Platyptilia* 133  
*campylocha, Stathmopoda* 108  
*cana, Ichneutica* 205  
*Mallobathra* 68  
*canalis, Pterophorus* 135  
*candida, Epichorista* (auct. 1) 122  
*canescens, Heterocrossa* 130  
*Hydriomena* auct. 103  
*Xylina* 212  
*caprealis, Aglossa* 143  
*caprimulgata, Hemerophila* 165

- captiosa*, *Melanchra* 207  
*Capua* 119  
*caradjai*, *Symmoca* 85  
*carcharodes*, *Epichorista* (auct. 2) 122  
*carduidactyla*, *Platyptilia* 133  
*carnifex*, *Proteodes* 88  
*Carposina* 130  
**CARPOSINIDAE** 130  
*caryochroa*, *Eutorna* 87  
*cassina*, *Opsiphanes* 136  
*casta*, *Coremia* 184  
*castanea*, *Dryadaula* 60  
*catacaustus*, *Orocrambus* 146  
*cataclysma*, *Mallobathra* 67  
*catadees*, *Dasyuris* 178  
*Catamacta* 119  
*cataphracta*, *Asaphodes*  
*catapyrrha*, *Arctesthes* 171  
*cataxesta*, *Eudonia* 155  
*Cateristis* 79  
*catocalaria*, *Larentia* 186  
**CATOCALINAE** 196  
*Catopsilia* 139  
*Caunaca* 76  
*caustica*, *Sabatinca* 52  
*caustopa*, *Izatha* 94  
*cauta*, *Phylacodes* 74  
*cautella*, *Ephestia* 142  
*cawthonella*, *Reductoderces* 69  
*Cebysa* 67  
*cecropia*, *Hyalophora* 140  
*cedrinodes*, *Astrocidaria* 175  
*Celama* 213  
*celerio*, *Hippotion* 140  
*celidota*, *Stenoptilia* 134  
*cenchrrias*, *Gymnobathra* 92  
*Cephalissa* 176  
*ceramodes*, *Xanthia* 198  
*cerapachoides*, *Euxoa* 213  
*ceraunias*, *Ichneutica* 205  
*cerealella*, *Sitotroga* 83  
*certella*, *Thamnosara* 66  
*cervinata*, *Wiseana* 58  
*chalara*, *Eudonia* 155  
*Chalastra* 163  
*chalcodelta*, *Caloptilia* 71  
*chalcophanes*, *Sabatinca* 52  
*chalicodes*, *Scoparia* 159  
*chaophilta*, *Strepsicrates* 116  
*chaotica*, *Cidaria* 183  
*characta*, *Eudonia* 155  
    *Microcolona* 86  
*charactana*, *Cacoecia* 120, 125  
*characterifer*, *Dumbletonius* 57  
*charadriias*, *Stenoptilia* 134  
*charadrota*, *Erechthias* 61  
*Charagia* 56  
*charaxias*, *Heterocrossa* 130  
*charidema*, *Epiphryne* 179  
*Charixena* 73  
*charopa*, *Strepsicrates* 116  
*chartularia*, *Orthenches* 73  
*charybdis*, *Pasiphila* 187  
*chasmatis*, *Erechthias* 61  
*chathamica*, *Proterodesma* 64  
*chatuidea*, *Asterivora* 113  
*cheradias*, *Kiwaia* 81  
*Chersadaula* 91  
*chimeria*, *Scoparia* 157  
*chioneres*, *Notoreas* (Meyrick) 184  
*chionochloae*, *Proditrix* 75  
*chionodira*, *Erechthias* 61  
*chionogramma*, *Asaphodes* 172  
*chirista*, *Thamnosara* 98  
*chlamydota*, *Asaphodes* 172  
    *Eudonia* 155  
**CHLIDANOTINAE** 114  
**CHLOEOPHORINAE** 198  
*chloradelpha*, *Tingena* 100  
*chloratma*, *Euthictis* 92  
*chlorias*, *Orthoclydon* 185  
*chloritis*, *Tingena* 100  
*chlorobela*, *Hierodoris* 89  
*chlorocapna*, *Asaphodes* 172  
*Chloroclystis* auct. 176  
*chlorocoma*, *Orthenches* 73  
*chlorodontia*, *Graphania* 202  
*chlorograpta*, *Meterana* 207  
*chloroleuca*, *Dolichernis* 70  
*chloronota*, *Atomotricha* 90  
*chlorosaris*, *Argyroploce* 114  
**CHOREUTIDAE** 113  
*chorica*, *Larentia* 174  
*choristis*, *Eudonia* 155  
*christiani*, *Pseudocoremia* 168  
*christinae*, *Helastia* 180  
*chrysargyra*, *Sabatinca* 52  
*chryserythra*, *Graphania* 202  
*chrysippus petilia*, *Danaus* 136  
*chrysitis*, *Caloptilia* 71  
*chrysochyta*, *Glaucocharis* 144  
*Chrysodeixis* 213  
*Chrysoesthia* 81  
*chrysogramma*, *Tingena* 100  
*chrysograpta*, *Astrogenes* 60  
*chrysomela*, *Pyrgotis* 127  
*chrysopeda*, *Notoreas* (Meyrick) 185  
*churtoni*, *Izatha* 94  
*cidiaria*, *Eupithecia* 188  
*cimmeria*, *Tephrosaria* 65  
*cinefacta*, *Scoparia* auct. 160  
*cineracia*, *Pseudocoremia* 167  
*cinerascens*, *Itama?* 163  
*cinerea*, *Euchlaena* 166

- cineraria*, *Helastia* 180  
*cinerosa*, *Agriophara* 109  
*cinigerella*, *Crocydopora* 141  
*cinnabari*, *Asaphodes* 172  
*cionophora*, *Glyphipteryx* 76, 77  
*Circica* 76  
*Circostola* 73  
*Circoxena* 86  
*ciserodes*, *Scoparia* 159  
*citharoda*, *Parectopa* 70  
*citrinata*, *Epiphryne* 179  
*citroena*, *Asaphodes* 173  
*Cladoxycanus* 57  
*clandestina*, *Helastia* 181  
*claranota*, *Scoparia* auct. 160  
*clarata*, *Asaphodes* 173  
*clarkei*, *Hydriomena* auct. 183  
*Orocrambus clarkei* 147  
*Planotortrix* 125  
*Proteodes* 88  
*Tingena* 100  
*clavata*, *Scoparia* auct. 160  
*claviferella*, *Aquita* 151  
*Cleora* 163  
*Clepsicosma* 152  
*Cnephasia* auct. 120  
*coarctatella*, *Gymnobathra* 93  
*Phaeosaces* 98  
*coctilis*, *Meterana* 206  
*codonias*, *Glyphipteryx*  
*coeleno*, *Meterana* 206  
*coerula*, *Arcte* 197  
*Coleophora* 85  
**COLEOPHORIDAE** 85  
**COLEOPHORINAE** 85  
*collectaria*, *Cidaria* 190  
*colligatella*, *Agriophara* 109  
*collitella*, *Tingena* 100  
*colpogramma*, *Pseudocoremia* 167  
*colpota*, *Asterivora* 113  
*Eudonia* 155  
*columbina*, *Stomopteryx* 80  
*columella*, *Harmologa* 124  
*comariana*, *Acleris* 129  
*comastis*, *Loxostege* 153  
*combinatana*, *Asterivora* 113  
*comma*, *Rictonis* 196  
*communicata*, *Orthosia* 212  
*comosaris*, *Borkhausenia* 105  
*compseuta*, *Bascantis* 62  
*Compsistis* 91  
*compsogramma*, *Tingena* 100  
*compta*, *Graphiphora* 212  
*comptella*, *Opogona* 63  
*compsotypa*, *Phaeosaces* 98  
*conditana*, *Adoxophyes* 119  
*Planotortrix* 125  
*conferta*, *Heliothis (Helicoverpa) armigera* 211  
*Tinea* sensu Meyrick 65  
*confusella*, *Barea* 90  
*congestana*, *Teras* 121  
*congregata*, *Cidaria* 183  
*congressata*, *Cidaria* 183  
*conifera*, *Scoparia* 160  
*conisalia*, *Scoriodyta* 69  
*conopias*, *Crambus* 150  
**Conopomorpha** 72  
*consentiens*, *Pyrgotis* 127  
*consignatella*, *Barea* 91  
*conspecta*, *Tinea* sensu Meyrick 65  
*conspicuella*, *Trachypepla* 106  
*constrictana*, *Capua* 119  
*contactella*, *Heterocrossa* 130  
*contexta*, *Scoparia* auct. 161  
*contextella*, *Tingena* 101  
*contractana*, *Teras* 121  
*contraria*, *Kiwaia* 81  
*contritella*, *Trachypepla* 106  
*conversata*, *Cidaria* 183  
*convolvuli*, *Agrius* 140  
*convulsella*, *Izatha* 94  
*cookaria*, *Zanclopteryx?* 169  
*copidota*, *Kessleria* 73  
*copiosella*, *Izatha* 95, 96  
*coprobiella*, *Trichophaga* 66  
**COPROMORPHIDAE** 129  
**COPROMORPHOIDEA** 129  
*coprosmae*, *Planotortrix* 125  
*copularis*, *Porina* 58  
*coracodes*, *Stathmopoda* 108  
*corcularia*, *Helastia* 181  
*cordalis*, *Sceliodes* 153  
*coricopa*, *Hypeuryntis* 109  
*Coridomorpha* 88  
*Corocosma* 91  
*correlata*, *Campptogramma* 193  
*corruptus*, *Orocrambus* 147  
*corylana*, *Crambus* 147  
*cosmia*, *Anisoplaca* 80  
*Cosmiotes* 110  
*Cosmodes* 195  
*cosmodora*, *Asaphodes* 173  
**COSMOPTERIGIDAE** 86  
*Cosmopterix*  
**COSSIDAE** 112  
**COSMOIDEA** 112  
*costaestrigalis*, *Schrankia* 211  
*cotinaea*, *Pasiphila* 187  
**CRAMBIDAE** 143  
**CRAMBINAE** 143  
*crassa*, *Xyleutes* 112  
*crassitibia*, *Amphitape* 165  
*crataea*, *Mallobathra* 68  
*Cremonogenes* 99

- cremnapa*, *Selidosema* 168  
*crenaeus*, *Orocrambus* 147  
*crepuscula*, *Scoparia* auct. 161  
*critica*, *Eudonia* 155  
*crociicapitella*, *Monopis* 63  
*Crocidosema* 115  
*crocostoma*, *Taoscelis* 89  
*Crocydopora* 141  
*crotala*, *Tingena* 101  
*crucifera*, *Calicotis* 109  
*cryodana*, *Heterocrossa* 130  
*crypsidora*, *Epichorista* (auct. 2) 122  
*crysipimima*, *Erechthias* 61  
*crysinoa*, *Eudonia* 155  
*Crypsitricha* 60  
*Cryptaspasma* 115  
*cryptica*, *Helastia* 181  
*Cryptolechia* auct. 87, 98  
*Cryptomima* 144  
*Ctenoplusia* 213  
*Ctenopseustis* 120  
**CTENUCHIDAE** 195  
**CUCULLIINAE** 198  
*cucullina*, *Aletia* 199  
*Culladia* 144  
*culta*, *Archyala* 59  
*cultus*, *Orocrambus* 147  
*cuneata*, *Aletia* 199  
*Ericodesma* 117  
*cuneiferana*, *Teras* 121  
*cuneigera*, *Teras* 124  
*cuprea*, *Orthenches* 74  
*curtisella*, *Prays* 75  
*curva*, "Eurythecta" auct. 123  
*Curvisacculus* 121  
*cyameuta*, *Scoparia* 159  
*cyanopetra*, *Aletia* 199  
*cyanospila*, *Conopomorpha* 72  
*cyclobathra*, *Maoritenes* 128  
*cyclopicus*, *Orocrambus* 147  
*Cydia* 115  
*cymatias*, *Eudonia* 155  
*cymatodes*, *Dasypodia* 197  
*cymodoce*, *Tinea* sensu Meyrick 65  
*cymosema*, *Eurydice* 182  
*cymozeucta*, *Helastia* 181  
*cynica*, *Dichromodes* 191  
*Cynthia* 137  
*cynthia*, *Samia* 140  
*cyphonias*, *Trachypepla* 106  
*cypracma*, *Stigmella* 53  
*cyptastis*, *Eudonia* 155
- D**
- daiclesalis*, *Mnesictena* 153  
*dalella*, *Rhigognostis* 76
- DANAINAE** 136  
*Danaus* 136  
*Daphnis* 140  
*Dasypodia* 197  
*Dasyuris* 177  
*deamatella*, *Pyroderces* auct. 86  
*Deana* 152  
*debilis*, *Hadena* 204  
*deceptura*, *Xylina* 201  
*Declana* 164  
*declarata*, *Asaphodes* 173  
*declavata*, *Scoparia* auct. 162  
*declivis*, *Scoparia* auct. 161  
*decora*, *Tingena* 101  
*decoranda*, *Erechthias* 61  
*decorata*, *Meterana* 206  
*defigurata*, *Bityla* 195  
*deflorata*, *Hypocala* 197  
*dejectaria*, *Gellonia* 165  
*delicatulata*, *Xanthorhoe* auct. 191  
*Delogenes* 141  
*deltoidata*, *Hydriomena* auct. 183  
*deltophora*, *Eudonia* 155  
*demiana*, *Tortrix* auct. 128  
*demissa*, *Sabatinca* 52  
*denotata*, *Scotosia* 177  
*dentata*, *Aletia* 199  
*dentigera*, *Leucania* 208  
**DEPRESSARIINAE** 87  
*deprivatalis*, *Platyptilia* 133  
*derogatella*, *Eschatotypa* 62  
*descendens*, *Porina* 57  
*descriptata*, *Cidaria* 183  
*desiccata*, *Hyperythra* 164  
*despecta*, *Hepialus* 58  
*detritana*, *Sciaphila* 119  
*Detunda* 164  
*Diarsia* 212  
*Diasemia* 152  
*diatmeta*, *Meterana* 206  
*dicharacta*, *Tinea* sensu Meyrick 65  
*dichorda*, *Glyphipterix* 77  
*Dichromodes* 191  
*dicrenellus*, *Orocrambus* 147  
*dictyarcha*, *Orthenches* 74  
*Dictyoploca* 140  
*diffusaria*, *Larentia* 181  
*dimorphella*, *Monopis* 63  
*dinacosma*, *Gymnobathra* 92  
*dinocosma*, *Gymnobathra* 92  
*dinodes*, *Aoraia* 56  
*Eudonia* 155  
*dione*, *Ichneutica* 205  
*dionysias*, *Asaphodes* 173  
*Dioxycanus* 57  
*Dipaustica* 201  
*diphtheralis*, *Scoparia* auct. 161

- diploorrhous*, *Crambus* 147  
*Diploseustis* 143  
*dipsasalis*, *Scopula?* 153  
*Dipterina* 128  
*discariana*, *Acroclita* 114  
*disjunctella*, *Vanicela* 70  
*disjungens*, *Graphania* 202  
*disparilis*, *Orthenches* 74  
*dissimilis*, *Aponotoreas* 171  
*dissociata*, *Cidaria* 181  
*distans*, *Sphinx* 140  
*distincta*, *Ctenopseustis* 126  
    *Stathmopoda* 108  
*distracta*, *Melanchra* 205  
*ditrocha*, *Circoxena* 86  
**DITRYSIA ENDOPORIA** 59  
**DITRYSIA EXOPORIA** 55  
*dives*, *Ichneutica* 206  
    *Melanchra* 202  
*dividua*, *Kiwaia* 82  
    *Tinea* sensu Meyrick 66  
*divulsana*, *Merophyas* 118  
*dochmia*, *Eudonia* 156  
*Dodonidia* 138  
*Dolichernis* 70  
*dolopaea*, *Strepsicrates* 116  
*Donacostola* 110  
*donovani*, *Elvia* 178  
*doroxena*, *Micropardalis* 52  
*dorsana*, *Grapholita* 115  
*dotata*, *Meterana* 206  
*doubledayi*, *Nyctemera* 194  
*Doxophyrtis* 73  
**DREPANOIDEA** 193  
*drosocalca*, *Orthenches* 74  
*drosophaea*, *Phryganostola* 76  
*drurella*, *Chrysoesthia* 81  
*Dryadaula* 60  
*dryas*, *Pasiphila* 187  
*dryocyma*, *Eucymatoge* 175  
*dryphactis*, *Scoparia* 159  
*dubia*, *Tanaoctena* 76  
*dubiella*, *Tinea* 65  
*Dumbletonius* 57  
*dunedensis*, *Leucania* 210
- E**
- Ecclitica* 121  
*Ectopatria* 212  
*edna*, *Lycaena* 135  
*egregia*, *Declana* 164  
*ejectana*, *Strepsicrates* 116  
*ejuncida*, *Scoparia* 159  
*Elachista* 111  
*Elachista* auct. 112  
**ELACHISTIDAE** 111
- elaeas*, *Caloptilia* 71  
*elaina*, *Glaucocharis* 144  
*elaphra*, *Antiscopa* 154  
*electilis*, *Kupea* 145  
*electrica*, *Hierodoris* 89  
*elegans*, *Cosmodes* 195  
*elephantina*, *Epichorista* (auct. 2) 122  
*elutella*, *Epehestia* 142  
*Elvia* 178  
*Elygea* 197  
*emphanes*, *Epichorista* (auct. 1) 122  
**Empista**  
*emplasta*, *Strepsicrates* 116  
*empyrea*, *Aletia* 199  
*encapna*, *Scoparia* 159  
*encausta*, *Amastris* 169  
    *Tortrix* 121  
*enchophorus*, *Orocrambus* 147  
*Endophthora* 61  
*Endotricha* 143  
*endotherma*, *Stathmopoda* 108  
*Endrosis* 91  
**ENNOMINAE** 163  
*enodis*, *Tingena* 101  
*enoplana*, *Cacoecia* 126  
*enysii*, *Aoraia* 56  
    *Dasyuris* 178  
    *Chrysophanus* 135  
*eodora*, *Sabatinca* 53  
*Epalxiphora* 122  
**EPERMENIIDAE** 132  
**EPERMENIOIDEA** 132  
*Ephesia* 142  
*ephorus*, *Orocrambus* 147  
*ephyraria*, *Panagra* 169  
*Epialus* 58  
*epiastra*, *Dipaustica* 201  
*Epicasis* 164  
*epicalca*, *Tingena* 101  
*epichlora*, *Stathmopoda* 109  
*Epichorista* (Meyrick) 122  
*epicomia*, *Antiscopa* 154  
*epicremna*, *Eudonia* 156  
*epicryptis*, *Microdes* 184  
*epicura*, *Harmologa* 122  
*Epicyme* 178  
*epimylia*, *Tingena* 101  
*Epiolus* 58  
*epiphaea*, *Glaucocharis* 144  
*epiphanes*, *Izatha* 95  
*Epiphryne* 179  
*Epiphthora*  
*Epiphyas* 117  
*episema*, *Xanthorhoe* 175  
*epistrota*, *Scythris* 110  
*Epitymbiini*  
*epixyla*, *Lysiphragma* 62

- epomiana*, *Heterocrossa* 130  
*epomiana* auct., *Carposina* 131  
*epotis*, *Stenoptilia* 134  
*Epyaxa* 179  
*Eraea* 201  
*erastis*, *Glyphipteric* 77  
*eratopis*, *Protosynaema* 75  
*erebia*, *Graphania* 202  
*erebinata*, *Scotosia* 165  
*Erebiola* 138  
*erebistis*, *Erechthias* 62  
*erebopsis*, *Heliothela* 132  
*Erechthias* 61  
*erechtitus*, *Stigmella* 53  
*eremana*, "Eurythecta" auct. 123  
*eremita*, *Hierodoris* 89  
*Eretria* 154  
*ergatis*, *Scoparia* auct. 161  
*eribola*, *Epichorista* (auct. 2) 122  
*erichrysa*, *Metacrias* 194  
*Ericodesma* 117  
*eriosoma*, *Chrysodeixis* 213  
*eriphaea*, *Tingena* 101  
*eriphylla*, *Heterocrossa* 130  
*erratica*, *Pasiphila* 187  
*Eschatotypa* 62  
*ethela*, *Gracilaria* 72  
*ethelella*, *Monopis* 63  
*Etiella* 142  
*euastera*, *Glyphipteric* 77  
*euboliaria*, *Aspilates* 170  
*eucalypti*, *Antheraea* 140  
*Euchersadaula* 91  
*eucola*, *Batrachedra* 84  
*eucrossa*, *Aletia* 201  
*eucyrtta*, *Epiphyas* 117  
*Eudonia* 154  
*eudorana*, *Pyrgotis* 127  
*Eugennaea* 62  
*Eulechria* 91  
*eumenopa*, *Tingena* 101  
*eumetalla*, *Acrocercops* 71  
*eupitheciaria*, *Helastia* 181  
*Euprocteodes* 111  
*curybathra*, *Kiwaia* 82  
*curychora*, *Elachista* 111  
*Eurydice* 182  
*curygrapha*, *Thallostoma* 65  
*curyleucota*, *Trachypepla* 106  
*Eurythecta* 118  
*Eurythecta* auct. 123  
*eustyla*, *Cateristis* 79  
*Euthictis* 92  
*euthrepta*, *Glaphyrarcha* 130  
*Eutorna* 87  
*Euxoa* 212  
*exanimis*, *Eulechria* 91  
*exarcha*, *Barea* 91  
*exaula*, *Cosmiotes* 111  
*excessana*, *Planotortrix* 125  
*exempta*, *Spodoptera* 196  
*exilis*, *Scoparia* auct. 161  
*eximia*, *Orocrambus clarkei* 147  
*exocha*, *Asterivora* 113  
*exochana*, *Heterocrossa* 130  
*Exoria* 115  
*exoriens*, *Asaphodes* 173  
*exospila*, *Erechthias* 61  
*expolita*, *Helastia* 181  
*exprompta*, *Boarmia* 165  
*exquisita*, *Meterana* 207  
*exsanguis*, *Sagephora* 64  
*exsiccata fallax*, *Tathorhynchus* 198  
*Exsiliracha* 159  
*exsommis*, *Atomotricha* 90  
*exsularis*, *Hypenodes* 212  
*extensalis*, *Daraba* 154  
*externella*, *Erechthias* 61
- F**
- fagana*, *Pseudoips* 198  
*fagicola*, *Tinea* sensu Meyrick 66  
*falcata*, *Homodotis* 182  
*falcatalis*, *Platyptilia* 133  
*falcatella*, *Samana* 192  
*fallax*, *Mnesarchaea* 55  
*Tathorhynchus* 198  
*falsa*, *Scoparia* auct. 161  
*falsidica*, *Aletia falsidica* 199  
*falsiloqua*, *Tingena* 101  
*famularis*, *Scoparia* auct. 161  
*farinalis*, *Pyralis* 143  
*farinaria*, *Patagoniodes* 142  
*farinata*, *Helastia* 181  
*fascialata*, *Pseudocoremia* 167  
*fascialis*, *Hyenia* 153  
*fasciata*, *Arctesthes* 171  
*Asterivora* 113  
*Horisme* 175  
*Tatosoma* 190  
*fastigata*, *Planotortrix* 125  
*felix*, *Gonodontis* 166  
*Sagephora* 64  
*fenerata*, *Pseudocoremia* 167  
*fenestrata*, *Tingena* 101  
*fenwicki*, *Graphania* 202  
*Mallabathra* 68  
*feredayi*, *Declana* 164  
*Eudonia* 156  
*Lycaena* 135  
*Feredaya* 201  
*ferox*, *Paranotoreas* 186  
*ferruginella*, *Monopis* 63  
*fervida*, *Tortrix* auct. 128

- festiva*, *Harmologa* 124  
*Trachypepla* 106  
*fibriata*, *Aletia* 199  
*figlinaria*, *Acidalia* 193  
*filata*, *Chloroclystis* auct. 176  
*filicicola*, *Batrachedra* 84  
*fimbriata*, *Scoparia* auct. 161  
*fischeri*, *Charagia* 56  
*flammea*, *Planotortrix* 126  
*flava*, *Anomis* 197  
*Pseudocoremia* 167  
*flavescens*, *Planotortrix* 126  
*flavidalis*, *Mnesictena* 153  
*flavidella*, *Gymnobathra* 92  
*flexata*, *Sestra* 169  
*flexivittana*, *Sciaphila* 120  
*flexuosellus*, *Orocrambus* 148  
*floccosa*, *Declana* 164  
*florida*, *Izatha* 95  
*fluminea*, *Pseudocoremia* 167  
*formosa*, *Macarostola* 72  
*fortinata*, *Ischalis* 166  
*fortis*, *Homohadena* 199  
*fragilis*, *Mallobathra* 69  
*fragosata*, *Selidosema?* 168  
*fraterna*, *Ctenopseustis* 120  
*fraudulenta*, *Epichorista* (auct. 2) 122  
*fraxinea*, *Anisoplaca* 80  
*freta*, *Borkhausenia* 103  
*frigida*, *Hierodoris* 89  
*Xanthorhoe* auct. 191  
*frischella*, *Coleophora* 85  
*frivola*, *Asaphodes* 173  
*frontella(?)*, *Graciliaria* 70  
*fugitivana*, *Protithona* 116  
*fugitivellus*, *Orocrambus* 148  
*fulguritella*, *Erechthias* 62  
*fuliginea*, *Wiseana* 58  
*fullonia*, *Othreis* 198  
*fulminea*, *Dasyuris* 178  
*fulva*, *Paranotoreas* 186  
*Stigmella* 53  
*fumata*, *Scoparia* auct. 161  
*fumipalpata*, *Pasiphila* 187  
*funerea*, *Aletia* 199  
*furcatalis* var.  $\beta$ ,  $\gamma$ , *Aciptilus* 133  
*Pterophorus* 132  
*furcillata*, *Tinea* sensu Meyrick 66  
*furtiva*, *Melanchra* 202  
*furva*, *Pasiphila* 187  
*fusca*, *Locheutis* 97  
*Mnesarchaea* 55  
*Porina* 57  
*fuscinata*, *Camptogramma* 170  
*fuscipunctella*, *Tinea* 63  
*fuscoflava*, *Reductoderces* 69  
*fucus*, *Dioxycanus* 57  
*fusiferana*, *Sciaphila* 128  
*fusilella*, *Mnesarchaea* 55  
*fusilis*, *Stathmopoda* 109  
*fusiplagiata*, *Sestra* 169
- G**
- Gadira* 143  
*gahniae*, *Proditrix* 75  
*galactilis*, *Scoparia* 158  
*galactina*, *Nymphostola* 88  
*galatheae*, *Elachista galathea* 112  
*galaxias*, *Notoreas* (Meyrick) 185  
*Trachypepla* 106  
*gallaria*, *Ischalis* 166  
*Galleria* 141  
**GALLERIINAE** 141  
*gallicolens*, *Cacoecia* 141  
*Gargaphia* 169  
*Gauna* 143  
*gavisana*, *Catamacta* 119  
*Gelechia* auct. 81  
**GELECHIIDAE** 80  
**GELECHIOIDEA** 80  
*Gellonia* 165  
*Gelonia* 165  
*Gelophaula* 123  
*generosa*, *Crypsitricha* 60  
*Gen. indet.* (Psychidae) 69  
**GEOMETRIDAE** 163  
**GEOMETROIDEA** 163  
*gerasmia*, *Elachista* auct. 112  
*Gingidiobora* 180  
*glacialis*, *Declana* 164  
*glaciata*, *Asaphodes* 173  
*Glaphyrracha* 130  
*glaucata*, *Elvia* 178  
*Glaucocharis* 144  
*glaucophanes*, *Tawhitia* 151  
*glaucoterma*, *Kiwaia* 82  
*globulosa*, *Mallobathra* 68  
*glyciniae*, *Phalaenodes* 196  
*glycinopa*, *Dryadaula* 60  
*glypharcha*, *Orthenches* 74  
**GLYPHIPTERIGIDAE** 76  
*Glyphipterix*  
*gobiata*, *Astrocidaria* 175  
*gonerilla*, *Bassaris gonerilla* 136  
*Gonophylla* 166  
*gonosemana*, *Heterocrossa* 131  
*gorgopis*, *Tauroscopa* 150  
*gourlayi*, *Aletia* 200  
*Porina* 57  
*gracilis*, *Crambus* 145  
*Scoparia* auct. 161  
**GRACILLARIIDAE** 70  
*graminea*, *Exsilirarcha* 159

*graminosa*, *Feredayia* 201  
*grammalis*, *Diasemia* 152  
*grammocosma*, *Prothinodes* 64  
*Grammodes* 197  
*grandiosa*, *Melanchra* 207  
*granella*, *Nemapogon* 63  
*Graphania* 201  
*Grapholita* 115  
*grata*, *Tingena* 101  
*gregalis*, *Helostibes* auct. 89  
*gressitti*, *Eudonia* 156  
*griseata*, *Adeixis* 191  
    *Declana* 164  
    *Izatha* 95  
    *Tingena* 101  
*grisescens*, *Dasyuris anceps* 177  
*griseipennis*, *Mamestra* 200  
*grisella*, *Achroia* 141  
*Grypotheca* 67  
*gurri*, *Pareromene* 145  
*Gymnobathra* 92  
*gypsotis*, *Dichromodes* 191  
*gyrotoma*, *Eudonia* 156

## H

*haastaria*, *Zanclopteryx?* 169  
*haasti*, *Platyptilia* 133  
***Habrophila* 62**  
*habropis*, *Gymnobathra* 93  
**HADENINAE 199**  
*haemophaea*, *Astrocidaria* 175  
*halianthes*, *Pasiphila* 188  
*halopis*, *Scoparia* 159  
*halosparta*, *Eschatotypa* 62  
*hamadelpha*, *Mnesarchaea* 55  
*hamatella*, *Gymnobathra* 92  
*hamatus*, *Tiramula hamatus* 136  
*hamiltoni*, *Aletia* 199  
*haplotomus*, *Orocrambus* 148  
*hapsimacha*, *Amphixystis* 59  
*hardenbergiella*, *Lithocolletis* 72  
*Harmologa* 124  
*harmonica*, *Glaucocharis* 144  
    *Hydriomena* 175  
*harpalaea*, *Scoparia* 159  
*Harpalyce* 182  
*harpophorus*, *Orocrambus* 148  
*harrisii*, *Argyrophenga* 138  
*hartii*, *Tmetolophota* 209  
*hastata*, *Tingena* 101  
*hawthornei*, *Hydriomena* auct. 183  
*Hectacma* 61  
*hectori*, *Aenetus* 56  
    *Dasyuris* 178  
*heightwayi*, *Pasiphila* 188  
    *Sabatinca* 52

*Helastia* 180  
*Helastiodes* 186  
*helias*, *Asaphodes* 173  
*heliastis*, *Platyptilia* 133  
*Helicoverpa* 211  
*heliocypa*, *Glaucocharis* 144  
*Helostibes* auct. 88, 89  
*helotes*, *Orocrambus* 148  
*Heliothela* 152  
**HELIOTHINAE 210**  
*Heliothis* 210  
*Helleia* 135  
*Hellula* 152  
*helmsi*, *Dodonidia* 138  
*helmsii*, *Dodonidia* 138  
*helonoma*, *Cosmiotes* 111  
*hemiclista*, *Ecclitica* 121  
*hemiclistra*, *Erechthias* 62  
*hemicycla*, *Eudonia* 156  
*hemimochla*, *Tingena* 102  
*hemionana*, *Epichorista* 122  
*hemiplaca*, *Eudonia* 156  
*hemizona*, *Hydriomena* auct. 183  
*Hendecasticha* 115  
**HEPIALIDAE 56**  
**HEPIALINAЕ 56**  
**HEPIALOIDEA 56**  
*heptacentra*, *Paramorpha* 131  
*herana*, *Teras* 121  
*hermanella*, *Chrysoesthia* 81  
*Hermione* 179  
*hermione*, *Declana* 165  
*heroica*, *Izatha* 95  
*heteranthes*, *Crambus* 146  
*heteraulus*, *Orocrambus* 148  
**Heterocrossa**  
*heterospora*, *Kiwaia* 82  
*hexaleuca*, *Notoreas* (Meyrick) 185  
*Hierodoris* auct. 73, 89  
*Hierodoris* Meyrick 89  
*hieropis*, *Trachypepla* 106  
*Hierohestis* 63  
*hippeis*, *Kiwaia* 82  
*Hippolyte* 178  
**Hippotion 140**  
*Hofmannophila* 94  
*hokowhitalis*, *Platyptilia* 133  
*holanthes*, *Glaucocharis* 145  
*hollandiae* auct., *Dasygaster* 201  
*holochalca*, *Pantosperma* 78  
*holochra*, *Stathmopoda* 108  
*holorphna*, *Cnephasia* auct. 120  
*homalocyma*, *Xanthorhoe* 180  
*homalopa*, *Mallobathra* 68  
*homerica*, *Ichneutica* 206  
    *Orthenches* 74  
*Homodotis* 182

- homodoxa*, *Tingena* 102  
*Homoeosoma* 142  
*Homohadena* 199  
*homomorpha*, *Stathmonyma* 178  
*homoscia*, *Graphania* 202  
*honesta*, *Tingena* 102  
*honorata*, *Tingena* 102  
*hookeri*, *Elachista* 112  
*hoplodesma*, *Tingena* 102  
*horaea*, *Tingena* 102  
*Horisme* 182  
*horistes*, *Orocrambus* 148  
*horningae*, *Grypotheca* 67  
*horticola*, *Stathmopoda* 108  
*howesi*, *Tauroscopa* 150  
*howesii*, *Lysiphragma* 62  
*hudsoni*, *Heterocrossa philpotti* 131  
    *Izatha* 95  
    *Metacrias* 194  
    *Pasiphila* 187  
    *Physetica* 209  
    *Xyridacma* 193  
*humeraria*, *Sestra* 169  
*humerata*, *Scotosia* 177  
*humilialis*, *Scoparia* auct. 161  
*humilis*, *Pasiphila* 188  
    *Pyrgotis* 127  
*humillima*, *Pseudocoremia*  
*huttoni*, *Izatha* 95  
    *Metacrias* 194  
*hyalinatus*, *Epiolus* 58  
*Hyalophora* 140  
*hybreasalis*, *Deana* 152  
*hydralis*, *Hellula* 152  
*Hydiromena* auct. 182  
*hydrocosma*, *Doxophyrtis* 73  
*hyetodes*, *Gymnobathra* 93  
*Hygraula* 151  
*Hymenia* 152  
*hymenopis*, *Protosynaema* 75  
**HYPENINAE** 211  
**HYPENODINAE** 211  
*Hyperaucha* 211  
*Hypeuryntis* 109  
*Hypocala* 197  
*Hypolimnas* 137
- I**
- ianthina*, *Lycaena boldenarum* 135  
    *Sabatinca* 52  
*Ichneutica* 205  
*ida*, *Asaphodes* 173  
    *Bassaris gonerilla* 137  
    *Chloroclystis* auct. 177  
    *Dichromodes* 192  
*idiogama*, *Tingena* 102
- ignobilis*, *Heterocrossa* 131  
*illecta*, *Platysenta* 196  
*illectella*, *Cosmiotes* 111  
*illita*, *Hierodoris* 89  
*illota*, *Scoparia* auct. 161  
*illistris*, *Reductoderces* 69  
*imbriferana*, *Dipterina* 128  
*immunis*, *Taeniocampa* 212  
*imperfecta*, *Asaphodes* 173  
*impletus*, *Hepialus* 57  
*implexa*, *Graphiphoga* 196  
*importuna*, *Trachypepla* 106  
*improba*, *Trithamnora* 66  
*impudicis*, *Chloroclystis* auct. 176  
*inaemonaria*, *Coremia* 180  
*inaequata*, *Gymnobathra* 93  
*inana*, *Ctenopseustis* 121  
*inaptana*, *Teras* 124  
*incendaria*, *Tortrix* auct. 128  
*inceptura*, *Xylina* 201  
*incessana*, "Cnephasia" auct. 120  
*inchoata*, *Meterana* 207  
*inclarata*, *Cidaria* 183  
*inclinataria*, *Cidaria* 190  
*incompta*, *Aponotoreas* 171  
*incongruella*, *Sabatinca* 52  
*inconspicua*, *Chersotis* 213  
*inconstans*, *Aletia* 200  
*incrassatellus*, *Crambus* 150  
**INCURVARIINA** 59  
**INCURVARIOIDEA** 59  
*indecorella*, *Sphyrelata* 98  
*indicans*, *Erechthias* 62  
*indicataria*, *Eupithecia* 176  
*indistincta*, *Pseudocoremia* 167  
*indistinctalis*, *Scoparia* auct. 161  
*indocilisaria*, *Zermizinga* 170  
*indoescens*, *Trachypepla* 106  
*indomita*, *Tortrix* 124  
*indubitata*, *Platyptilia* 133  
*inductata*, *Chloroclystis* auct. 176  
*inexpiata*, *Eupithecia* 193  
*infantaria*, *Larentia* 181  
*infensa*, *Graphania* 202  
    *Strepsicrates* 116  
*infimana*, *Sciaphila* 116  
*infusa*, *Agrotis* 212  
*infusata*, *Larentia* 180  
*ingenua*, *Trachypepla* 107  
*innominata*, *Agrotis* 212  
*innotata*, *Leucania* 209  
*innotatalis*, *Pterophorus* 133  
*innotatana*, *Tortrix?* 119  
*innotella*, *Tingena* 102, 103  
*inoperata*, *Larentia* 180  
*inopiata*, *Cidaria* 183  
*inornata*, *Eutorna* 87

- insignis*, *Aponotoreas* 171  
*Graphania* 203  
*Hierodoris* 81  
*Stigmella* 53  
*insignita*, *Astrogenes* 60  
*Pseudocoremia* 167  
*insolita*, *Lopharcha* 114  
*inspoliata*, *Asterivora* 113  
*intactella*, *Gelechia* 118  
*interclusa*, *Larentia* 184  
*intermixta*, *Diarsia* 212  
*interpunctella*, *Plodia* 142  
*interrupta*, *Glaucocharis* 145  
*intractana*, *Sperchia* 128  
*intrudens*, *Eunychia* 153  
*inusitata*, *Tortrix* auct. 126  
*invexata*, *Larentia* 180  
*involuta*, *Anomis* 197  
*iocheaera*, *Glyphipterix* 77  
*iocchondra*, *Asterivora* 113  
*iophaea*, *Heterocrossa* 131  
*iophanes*, *Hierodoris* 89  
*iota*, *Misella* 199  
*Ipana* 164  
*ipson*, *Agrotis* 212  
*irabunda*, *Chloroclystis* 188  
*Irenicodes* 111  
*iridia*, *Clepsicosma* 152  
*iridoxa*, *Charixena* 73  
*irregulata*, *Chloroclystis* auct. 177  
*Phriessogonus*  
*Ischalis* 166  
*ischnycyma*, *Notoreas* (Meyrick) 185  
*ischnomorpha*, *Homoeosoma* 142  
*Isochasta* 81  
*isochytus*, *Orocrambus* 148  
*isogama*, *Atomotricha* 90  
*isoleuca*, *Notoreas* (Meyrick) 185  
*isomoera*, *Notoreas* (Meyrick) 185  
*Isonomeutis* 129  
*isoterma*, *Platyptilia* 133  
*itea*, *Bassaris* 137  
*Izatha* 94
- J**  
*jacobaeae*, *Tyria* 194  
*jactatana*, "Cnephasia" auct. 120  
*janata*, *Achaea* 196  
*janitae*, *Argyrophenga* 138  
*jansoni*, *Orocrambus* 148  
*jeanae*, *Kiwaia* 82  
*jocosa*, *Wiseana* 58  
*jocularis*, *Sagephora* 64  
*jubata*, *Dolichernis* 93  
*Gymnobathra* 93  
*juncicolor*, *Nonagria* 210
- junctilinea*, *Declana* 165  
*Junonia* 137
- K**  
*kaikourensis*, *Arctesthes* 171  
*kerguelensis*, *Pringleophaga* 64  
*kershawi*, *Cynthia* 137  
*Kessleria* 73  
*Kiwaia* 81  
*kuehniella*, *Ephestia* 142  
*Kupea* 145
- L**  
*labradus*, *Zizina labradus* 136  
*Laciarella* 114  
*lacteella*, *Endrosis* 91  
*lactiflua*, *Pseudocoremia* 168  
*lacustris*, *Aletia* 200  
*Elachista* 110  
*Pasiphila* 187  
*laetifica*, *Sphyrelata* 98  
*Lampides* 135  
*lancealana*, *Bactra* 115  
*Lantanophaga* 132  
*lapidosa*, *Mallobathra* 68  
*lapillosa*, *Kiwaia* 82  
*laquearia*, *Eugennaea* 62  
*laqueorum*, *Cosmiotes* 111  
*Planotortrix syntona* 126  
*Stigmella* 53
- LARENTIINAE** 170  
*lassa*, *Tingena* 102  
*lata*, *Aletia?* 205  
*Melanchna* 204  
*latens*, *Borkhausenia* 99  
*Lathicrossa* 96  
*lathionella herceus*, *Oreixenica* 138  
*lathriopa*, *Euchersadaula* 91  
*laticostatus*, *Phriessogonus* 189  
*latomana*, *Cnephasia* auct. 120  
*laudata*, *Tingena* 102  
*Lecithocera* 87
- LECITHOCERIDAE** 87  
*lectus*, *Orocrambus* 148  
*leda bankia*, *Melanitis* 138  
*legionaria*, *Scoparia* 160  
*legnota*, *Eudonia* 156  
*lenis*, *Kiwaia* 82  
*lenita*, *Leptocroca* 96  
*leonina*, *Aoraia* 56  
*Tawhitia* 151  
*lepidella*, *Glaucocharis* 145  
*leptalaea*, *Eudonia* 156  
*Leptocroca* 96  
*leptomera*, *Declana* 165  
*leptomitella*, *Stegommata* 79

- leptophaea*, *Scoparia* 156  
*Leptosaces* 92  
*leptosema*, *Glyptipterix* 77  
*Leptosoma* 194  
*lestevata*, *Tatosoma* 190  
*letharga*, *Tingena* 102  
*leucanialis*, *Chilo* 149  
*leucaniana*, *Merophysas* 118  
*leucelaea*, *Pseudocoremia* 168  
*leucobathra*, *Dasyuris* 178  
*leucocentra*, *Lathicrossa* 96  
*leucocyma*, *Acrocercops* 71  
*leucogramma*, *Eudonia* 156  
*leucophthalma*, *Gadira* 144  
*leucoplanetis*, *Trachypepla* 107  
*Leucoptera* 79  
*leucospora*, *Paracystola* 97  
*leucotelus*, *Cebysa* 67  
*leucothrinca*, "Eurythecta" auct. 123  
*leucoxantha*, *Glaucocharis* 145  
*levicula*, *Tingena* 103  
*levigata*, *Gymnobathra* 93  
*levis*, *Meterana* 207  
*lewisi*, *Atomotricha* 90  
    *Orocrambus* 148  
*liara*, *Euphyia* 184  
*lichenella*, *Cryptolechia* 95  
*lichenodes*, *Chloroclystis* 177  
    *Trachypepla* 107  
*lignana*, *Graphania* 203  
*ligniferana*, *Conchylis* 116  
*lignifusca*, *Hadena* 203  
*lignisecta*, *Xylina* 205  
*lignosata*, *Scotosia* 165  
*ligniverin*, *Aenetus* 56  
*lilacina*, *Herminia* 211  
*limatula*, *Scoparia* auct. 161  
*limbata*, *Oecophora* 93  
*Limnaecia* 86  
*limodes*, *Microcolona* 86  
*limodoxa*, *Delogenes* 142  
*limonodes*, *Asaphodes* 173  
*Lindera* 64  
*lindsayae*, *Archyala* 98  
*lindsayi*, *Archyala* 59  
    *Epichorista* (auct. 2) 122  
    *Graphania* 203  
    *Ichneutica* 206  
    *Leptocroca* 96  
    *Orocrambus* 148  
*linealis*, *Eudonia* 156  
*linearis*, *Caloptilia* 72  
*liochroa*, *Phaeosaces* 98  
*Liothula* 67  
*lissoxyla*, *Tmetolophota* 209  
*literata*, *Cadmogenes* 73  
    *Heterocrossa* 131  
*lithias*, *Graphania* 203  
*Lithilaria* 211  
*Lithocolletis* 72  
*lithodes*, *Kiwaia* 82  
*lithoxesta*, *Stenoptilia* 134  
*lithurga*, *Astrocidaria* 175  
*litterata*, *Batrachedra* 84  
*litura*, *Spodoptera* 196  
*Locheutis* 97  
*locularis*, *Eudonia* 156  
*longstaffi*, *Aletia* 200  
*Lopharcha* 114  
*lophogramma*, *Xanthorhoe* auct. 191  
*loreymimma*, *Mythimna* 208  
*lotinana*, *Catamacta* 119  
*lotrix*, *Utetheisa lotrix* 195  
*loxias*, "Eurythecta" auct. 123  
*loxoscia*, *Mnesarchaea* 55  
*Loxostege* 153  
*loxotis*, *Tingena* 103  
*lucida*, *Stigmella* 54  
*lucidata*, *Epyaxa* 179  
*lucilia*, *Sabatinca* 53  
*luciplagana*, *Paedisca* 127  
*ludibunda*, *Melancha* 208  
*lugubris*, *Cryptaspasma* 115  
*luminatrix*, *Eudonia* 156  
*luminosa*, *Chloroclystis* 188  
*lunata*, *Pasiphila* 188  
*lupinata*, *Pseudocoremia* 168  
*luridus*, *Crambus* 147  
*lutata*, *Prothiodes* 64  
*lutea*, *Pseudocoremia*  
*luteicornella*, *Lecithocera* 87  
*Lycaena* 135  
**LYCAENIDAE** 135  
**LYCAENINAE** 135  
*lychnopa*, *Erechthias* 62  
*lychnophanes*, *Gelophaula* 123  
    *Scoparia* auct. 161  
*lycosema*, *Acipitilia* 133  
**LYMANTRIIDAE** 195  
**LYONETIDAE** 79  
*Lyrcea* 192  
*Lysiphragma* 62  
*Lysiphragma* auct. 214  
*Lythria* auct. 184  
  
**M**  
*macarella*, *Tingena* 103  
*Macarastola* 72  
*machaeristes*, *Orocrambus* 148  
*macropetana*, *Strepsicrates* 116  
*macrozyga*, *Erechthias* 62  
*maculata*, *Chloroclystis* 187  
*Zelleria* auct. 76

- maculipennis*, *Plutella* 75  
*maculosa*, *Heterocrossa* 131  
  *Pseudocoremia* 168  
  *Tortrix* 118  
*magnimaculata*, *Pasiphila* 188  
*mahanga*, *Botys* 144  
*mahiana*, *Paedisca* 125  
*mairi*, *Porina* 58  
*malacelloides*, *Angustalius* 143  
*malachita*, *Pasiphila* 188  
*Mallobathra* 67  
*manganeutis*, *Eudonia* 156  
*manubriata*, *Izatha* 95  
*manxifera*, ?*Chlenias* 164  
*maori*, *Mamestra* 208  
*Maoria* 201  
*maoriana*, *Teras* 128  
*maoriaria*, *Xanthorhoe* 181  
*maoriata*, *Scotopteryx* 165  
***Maoricrambus*** 145  
*maoriella*, *Nephopteryx* 158  
  *Stigmella* 54  
***Maoritenes*** 128  
*maranta*, *Tingena* 103  
*marcida*, *Tingena* 103  
*margarita*, *Orthosia* 196  
*margaritis*, *Tinea* sensu Meyrick 66  
*marginana*, *Conchylis* 119  
*marginata*, *Paramorpha* 131  
*marmarea*, *Asterivora* 113  
*marmorina*, *Mnesictena* 153  
*marmorata*, *Ichneutica* 206  
*matermea*, *Kiwaia* 82  
*materna*, *Elyaea* 197  
*matutina*, *Protosynaema* 75  
*maui*, *Chrysophanus* 136  
*mauritia acronyctoides*, *Spodoptera* 196  
*maya*, *Graphania* 203  
*mechanitis*, *Notoreas* (Meyrick) 185  
*Mecyna* sensu Doubleday 154  
  *sensu* Meyrick 153  
*megalyntha*, *Proditrix* 75  
*Megacraspedus* auct. 83  
*megaspilata*, *Homodotis* 182  
*melampetrus* *Orocrambus* 148  
  *Orocrambus* 148  
*melanaegis*, *Eudonia* 157  
*melanamma*, *Tingena* 103  
*melanephele*, *Artigisa* 197  
***Melanitis*** 138  
*melanocentra*, *Chloroclystis* 188  
*melanochra*, *Stathmopoda* 108  
*melanombra*, *Epiphthora* 81  
*melanophaea*, *Cnephasia* auct. 120  
*melanoplitha*, *Phthorimaea* 83  
*melanosperma*, *Ericodesma* 117  
*melanotricha*, *Erechthias* 61  
*melanura*, *Elachista* auct. 112  
*melicerta*, *Phalaena* 197  
*melichrysa*, *Eschatotypa* 62  
*melinata*, *Pseudocoremia* 168  
*melinella*, *Tingena* 103  
*melitastes*, *Orocrambus* 148  
*meliturga*, *Eudonia* 157  
*mellonella*, *Galleria* 141  
*melochlora*, *Pasiphila* 188  
*melographa*, *Proteodes* 88  
*memorabilis*, *Corocosma* 91  
*memotuina*, *Mallobathra* 68  
*menanaria*, *Angerona* 163  
*meridiana*, *Melancha* 206  
*meristes*, *Crambus* 146  
*meritus*, *Crambus* 146  
***Mermeritis*** 97  
*merope*, *Meterana* 207  
***Merophysa*** 118  
*merula*, *Percnodaimon* 139  
*mesoschista*, *Izatha* 95  
*mesotypa*, *Crypsitricha* 60  
*messaniella*, *Phyllonorycter* 72  
*metachrysa*, *Phycomorpha* 129  
***Metacrias*** 194  
*metadelta*, *Izatha* 95  
*metallifera*, *Glaucoccharis* 145  
  *Trachypepla* 101  
*metasticta*, *Glyphipteric* 77  
***Meterana*** 206  
*metrosema*, *Mallobathra* 68  
*meyricci*, *Meterana* 207  
*Meyrickia* 163  
*micalis*, *Tebenna* 114  
*micrastra*, *Tmetolophota* 209  
*microbathra*, *Cnephasia* auct. 120  
***Microcolona*** 86  
***Microdes*** 184  
*microdora*, *Glaucoccharis* 145  
*microlitha*, *Asterivora* 113  
*micromela*, *Lecithocera* 87  
***Micropardalis*** 52  
*microphanes*, *Reductoderces* 69  
*microphthalma*, *Eudonia* 157  
*micropis*, *Morrisonia* 204  
*micropolis*, *Dasyuris* 178  
***MICROPTERIGIDAE*** 52  
***MICROPTERIGOIDEA*** 52  
*microtheriella*, *Stigmella* 54  
*milligani*, *Izatha* 94  
*mimica*, *Epichorista* (auct. 2) 123  
  *Wiseana* 58  
*miniella*, *Macarostola* 72  
*minima*, *Chloroclystis* 177  
  *Diploseustis* 143  
*ministra*, *Asterivora* 114  
*minos*, *Cladoxyccanus* 57

*minualis*, *Eudonia* 157  
*minuscularis*, *Scoparia* auct. 161  
*minuta*, *Trachypepla* 107  
*mira*, *Izatha* 95  
*miraculosa*, *Ochetarcha* 125  
*mistata*, *Sauris* 190  
*mitis*, *Aletia* 200  
*mixochlora*, *Lysiphragma* 63  
*Mnesarchaea* 55  
**MNESARCHAEIDAE** 55  
**MNESARCHAOIDEA** 55  
*mnesichola*, *Asaphodes* 173  
*Mnesictena* 153  
*moanalis*, *Scoparia* 156  
*mochlophorana*, *Parienia* 116  
*mochlota*, *Tinea* sensu Meyrick 66  
*Mocis* 197  
*moderata*, *Aletia* 200  
*modesta*, *Chloroclystis* 186  
    *Maoritenes* 128  
*modica*, *Pseudocoremia* 168  
*molesta*, *Grapholita* 115  
*molifera*, *Scoparia* auct. 161  
*mollis*, *Graphania* 203  
*molybditis*, *Tortrix* auct. 128  
**MOMPHIDAE** 87  
*monacha*, *Pseudocoremia* 168  
*monastra*, *Decadarchis* 61  
*monochroma*, *Scoparia* auct. 161  
*monodonta*, *Borkhausenia* 103  
*monoliata*, *Cidaria* 183  
*monophragma*, *Kiwaia* 82  
*Monopis* 63  
*monospilalis*, *Aciptilus* 133  
*monospilella*, *Tingena* 102  
**MONOTRYNSIA** 53  
*monoviridisata*, *Tatosoma* 190  
*morangella*, *Glyphipterix* 77  
*morbida*, *Heterocrossa* 131  
*morella*, *Borkhausenia* 91  
*mori*, *Bombyx* 140  
*morosa*, *Graphania* 203  
    *Tingena* 103  
*morosana*, *Paedisca* 129  
**Morova** 141  
*mucidalis*, *Sceliodes* 154  
*nullata*, *Asthena* 179  
*mundula*, *Agrotis* 212  
    *Aletia* 200  
    *Oecophora* 52  
*munita*, *Tinea* sensu Meyrick 66  
*muriferata*, *Sarisa* 169  
*muscosata*, *Pasiphila* 188  
*Musotima* 151  
**MUSOTIMINAE** 151  
*mutabilis*, *Helastia* 181  
*mutans*, *Graphania* 203

*mylites*, *Orocrambus* 148  
*myrrhina*, *Dryadaula* 60  
*mysteriastis*, *Stathmopoda* 108  
*mysticopa*, *Tinea* 64  
*mystis*, *Semiocosma* 95  
*Mythimna* 208

**N**

*namuella*, *Blabophanes* 63  
*nana*, *Sufetula* 143  
*napaea*, *Elachista* 112  
*nebulosa*, *Gingidiobora* 180  
    *Pasiphila* 188  
    *Tauroscopa* 146  
*necopina*, *Glyphipterix* 77  
*neglecta*, *Kiwaia* 82  
*negligens*, *Steganoptycha* 118  
*nehata*, *Cidaria* 182  
*nelsonaria*, *Ischalis* 166  
*Nemapogon* 63  
*neoselena*, *Drepanodes* 169  
*nephelias*, *Asaphodes* 173  
*nephelomima*, *Prays* 75  
*nephelotana*, *Prothelymna* 129  
*nephoptera*, *Glyphipterix* 78  
*Nepticula* 53  
**NEPTICULIDAE** 53  
**NEPTICULOIDEA** 53  
*nereis*, *Chloroclystis* auct. 177  
*nerina*, *Hypolimnas bolina* 137  
*nervata*, *Hadena* 202  
*nervosa*, *Ichneutica* 206  
*Nesarcha* 152  
*neurae*, *Leucania* 210  
*nexalis*, *Crambus* 150  
*Niditinea* 63  
*niger*, *Dichromodes* 192  
*nigra*, *Cremnogenes* 103  
    *Gymnobathra* 93  
    *Scythris* 110  
    *Tatosoma* 190  
*nigrofasciata*, *Euphyia* 176  
*nigrosparsa*, *Declana* 164  
*nimbosa*, *Trachypepla* 107  
*niphocrena*, *Notoreas* (Meyrick) 185  
*niphospora*, *Scoparia* auct. 161  
*niphostrota*, *Prothelymna* 129  
*niphozela*, *Scythris* 110  
*nitens*, *Hygraula* 151  
*nitidalis*, *Musotima* 151  
*Nitocris* 196  
*nivalis*, *Orthenches* 73  
*nivea*, *Epiphthora* 81  
*niveata*, *Declana* 165  
*nivescens*, *Asterivora* 114  
*nobilia*, *Aletia* 200

- NOCTUIDAE 195  
 NOCTUINAE 212  
 NOCTUOIDEA 194  
*nocturna*, *Rhathamictis* 69  
 NOLINAE 213  
*nomeutis*, *Scoparia* 159  
*notabilis*, *Donacostola* 110  
     *Tauroscopa* 150  
*notata*, *Ichneutica* 206  
     *Mnesictena* 153  
*Noteraula* 115  
*noteraula*, *Bactra* 115  
*notophaea*, *Planotortrix* 126  
*Notoreas* Meyrick 184  
*novaezealandiae*, *Porina* 58  
*nullifera*, *Graphania* 204  
*Nyctemera* 194  
*nycteris*, *Tingena* 103  
*nyctopis*, *Trachypepla* 106  
**NYMPHALIDAE** 136  
**NYMPHALINAE** 136  
*Nymphostola* 88
- O**  
*obarata*, *Asaphodes* 174  
*obliqua*, *Leptocroca* 97  
*obliquana*, *Ctenopseustis* 120  
*oblongana*, *Harmologa* 124  
*obscura*, *Euxoa* 213  
     *Mallabathra* 68  
     *Xanthorhoe* 172  
*obsecrata*, *Aletia* 200  
*obsistalis*, *Eretria* 154  
*obsoleta*, *Leucania* 209  
     *Xanthorhoe* 172  
*obstructus*, *Crambus* 150  
*obtruncata*, *Cidaria?* 170  
*obtusaria*, *Lozogramma* 170  
*occulta*, *Xanthorhoe* auct. 191  
*oceania*, *Melanchra* 202  
*Ochetarcha* 125  
*ochnosema*, *Cnephasia* auct. 120  
*ochrea*, *Chalastra* 163  
*ochreifera*, *Euphyia* 184  
*ochrogastra*, *Chersadaula* 91  
*ochroleuca*, *Cosmiotes* 111  
*ochthistis*, *Meterana* 207  
*ocneropis*, *Trachypepla* 97, 107  
*octans*, *Dasyuris* 178  
     *Meterana* 207  
*octias*, *Hyperaucha* 211  
*octomaculata*, *Epirrhantis* 192  
*octonaria*, *Glyphipterix* 78  
*octophora*, *Eudonia* 157  
*oculata*, *Eudonia* 157  
*oculiferella*, *Oligochroa* 142  
*oeconomia*, *Atomotricha* 90
- OECOPHORIDAE 87  
 OECOPHORINAE 88, 90  
*Oegoconia* 85  
 OENOCHROMINAE 191  
*Oenetus* 56  
*oenospora*, *Sporophyla* 143  
*oestriformis*, *Synanthedon* 112  
*officiosa*, *Hydriomena* 179  
*ogygia*, *Stigmella* 54  
*ohauensis*, *Helastia* 181  
*Oiceticus* 67  
*oleariae*, *Asterivora* 114  
     *Izatha* 95  
     *Thiotricha* 83  
**OLETHREUTINAE** 114  
*Oligichroa* 142  
*olivea*, *Graphania* 204  
*oliveri*, *Graphania* 204  
*ombrodella*, *Tingena* 104  
*ombrodes*, *Pseudocoremia* 168  
*ombrodoeca*, *Cosmiotes* 111  
*omichleuta*, *Gymnobathra* 93  
*omichlias*, *Asaphodes* 174  
*omicron*, *Graphania* 204  
*ommatis*, *Atomotricha* 90  
*omnivora*, *Liothula* 67  
*omogramma*, *Endophthora* 61  
*omoplaca*, *Graphania* 204  
*omoscopa*, *Opogona* 63  
*omphalota*, *Gymnobathra* 93  
*oncobolus*, *Maoricrambus* 146  
*oncolobus*, *Crambus* 146  
*ondinata*, *Asthena* 189  
*ooptila*, *Tingena* 76  
*opaca*, *Tingena* 104  
*operculella*, *Phthorimaea* 83  
**OPHIDERINAE** 196  
*ophiodryas*, *Tingena* 104  
*ophiopa*, *Goniophylla* 166  
*opipara*, *Paranotoreas* 186  
*Opodiphthera* 140  
*Opogona* 63  
*oporaea*, *Tingena* 104  
     “*Opostega*” 79  
*oppositus*, *Orocrambus* 148  
*optanias*, *Bactra* 115  
*opulenta*, *Archyala* 59  
*oraria*, *Asaphodes* 174  
*ordishi*, *Orocrambus* 149  
*oreas*, *Cladoxycanus* 57  
     *Eudonia* 157  
*Oreixenica* 138  
*organaea*, *Eudonia* 157  
*oriasta*, *Stigmella* 53, 54  
*orichalcea*, *Thysanoplusia* 214  
*origenes*, *Gymnobathra* 93  
*orites*, *Stenoptilia* 134

- ornatipenella*, Coleophora 85  
*ornatus*, Orocrambus 149  
*ornithias*, Monopis 63  
*Orocrambus* 146  
*Orophora* 67  
*orophylla*, Xanthorhoe auct. 191  
*orophylloides*, Xanthorhoe auct. 191  
*orphnaea*, Aponotoreas 171  
*Orthenches* 73  
*Orthoclydon* 185  
*orthocopa*, Planotortrix 126  
*ortholeuca*, Notoreas (Meyrick) 185  
*orthophanes*, Schiffermuelleria 98  
*orthopris*, Planotortrix 126  
*otagalis*, Botys 153  
*othello*, Oreina? 139  
*Othreis* 198  
*oxleyi*, Zizina labradus 136  
**OXYCANINAE** 57  
*oxyina*, Tingena 99, 104  
*oxymachaera*, Glyptipterix 78  
*oxyptera*, Asaphodes 174  
*Oxythecta* 97
- P**
- pachyerga*, Eudonia 157  
*pachyscia*, Leucania 200  
*pactolia*, Dryadaula 60  
*pagaia*, Graphania 204  
*Palaeomicra* 52  
*pallacopis*, Endophthora 61  
*pallescens*, Maoria 204  
*pallescentella*, Tinea 65  
*palliata*, Gelophaula 123  
*pallida*, Orthosia 195  
*pallidula*, Scoparia auct. 161  
  *Tingena* 104.  
*palthidiata*, Euchlaena? 166  
*paltomacha*, Eudonia 157  
*panacicorticis*, Acrocercops 71  
*panacifinens*, Acrocercops 71  
*panacitorsens*, Acrocercops 71  
*panacivagans*, Acrocercops 71  
*panacivermiformis*, Acrocercops 71  
*panagrata*, Scotosia 163  
*Pancyma* 179  
*panda*, Aletia 200  
*pannularia*, Gellonia 165  
*Panopaea* 179  
*panopla*, Scoparia auct. 161  
*pansicolor*, Meterana 207  
*pantheropa*, Mnesictena 153  
*Pantosperma* 78  
**PAPILIONOIDEA** 135  
*paracausta*, Graphania 204  
*parachalca*, Scoparia 160
- paracosma*, Mnesarchaea 55  
*paradelpha*, Notoreas (Meyrick) 185  
*paradesma*, Isochasta 81  
*Paradetis* 186  
*paraglypta*, Archyala 59  
*Paragyrtis* 191  
*parallela*, Epagoge 127  
*paralodes*, Chloroclystis 187  
*paraloxa*, Merophysas 118  
*Paramorpha* 131  
*paraneura*, Semiocosma 94  
*Paranotoreas* 186  
*parapleura*, Kiwaia 82  
*Parapoynx* 151  
*paratrimma*, Tingena 104  
*paraxenus*, Orocrambus 149  
*paraxysta*, Tmetolophota 210  
*parca*, Gymnobathra 93  
  *Scoparia* auct. 162  
*Pareromene* 144  
*Parienia* 115  
*parmata*, Aletia 199  
*parmifera*, Scoparia 160  
*Parocystola* 97  
*paronalis*, Scopula? 152  
*parora*, Austrocidaria 175  
*parorma*, Glaucocharis 145  
*parthenia*, Strepsicrates 117  
*partheniata*, Dasyuris 178  
*parvitis*, Celama 213  
*parvula*, Kiwaia 82  
*parvulata*, Phibalapteryx 177  
*Parysatis* 186  
*pascoella*, Scoparia auct. 162  
*pascoi*, Meterana 207  
*Pasiphila* 186  
*Pasithea* 184  
*passa*, Mamestra 204  
*passalota*, Sabatinca 53  
*pastinaria*, Coremia 183  
*Patagoniodes* 142  
*paterna*, Cnephasia auct. 120  
*patruelis*, Aciptilia 133  
*patularia*, Tephrosia 165  
*pauca*, Meterana 208  
*paucita*, Hydriomena 182  
*pauculana*, Teras 118  
*paula*, Tingena 104  
*pedias*, Orocrambus 147  
*pedella*, Stathmopoda 106  
*pelanodes*, Graphania 204  
*pelistis*, Mamestra 203  
*pellionella*, Tinea 65  
*pellurgata*, Chalastra 163  
*pelurgata*, Chalastra 163  
*pentadactyla*, Tawhitia 151  
*pentazyga*, Archyala 60

- penthalea*, *Tingena* 104  
*peracuta*, *Morrisonia* 208  
*percnitis*, *Izatha* 95  
**Percnodaimon** 138  
*perductata*, *Cidaria* 183  
*pergrata*, *Selidosema* 167  
*perichlora*, *Tingena* 104  
*perieresalis*, *Diploseustis* 143  
*periphanes*, *Eudonia* 157  
*peripheraea*, *Asaphodes* 174  
*perisseuta*, *Mallobathra* 68  
*perissopa*, *Stigmella* 54  
*pernyi*, *Antherea* 140  
*peroneanella*, *Izatha* 95  
*perornata*, *Notoreas* (Meyrick) 185  
*persecta*, *Epichorista* (auct. 2) 123  
**Persectania** 208  
*perspera*, *Rathamictis* 69  
*pertinax*, *Grypotheca* 67  
*perversata*, *Cidaria* 183  
*pervius*, *Orocrambus* 147  
*pessota*, *Andesia* 198  
*Petasactis* 63  
*petraula*, *Gadira* 144  
*petrias*, *Harmologa* 124  
*petrina*, *Dichromodes* 192  
*Scoparia* 160  
*petrodoxa*, *Mallobathra* 68  
*petrograpta*, *Graphania* 204  
*petropola*, *Larentia* 180  
*phaeoptila*, *Izatha* 96  
*Phaeosaces* 98  
*phaeoxyla*, "Eurythecta" auct. 123  
*Phalaenodes* 196  
*phalerias*, *Scoparia* auct. 162  
*pharetria*, *Kiwaia* 82  
*pharmactis*, *Tingena* 104  
*pharotoma*, *Crypsitricha* 60  
*phaula*, *Tmetolophota* 209  
*phaulacosma*, *Eutorna* 88  
*phegophylla*, *Tingena* 104  
*philadelphia*, *Gymnobathra* 93  
*philerga*, *Eudonia* 157  
*philetaera*, *Eudonia* 157  
*Philobota* 96  
*philocapna*, *Loxostege* 153  
*philochora*, *Locheutis* 97  
*Philocryptica* 125  
*philopoana*, *Tortrix* 118  
*philpotti*, *Asaphodes* 174  
*Curvisacculus* 121  
*Heterocrossa philpotti* 131  
*Orocrambus* 149  
*Philpottia* Meyrick 73  
*Philpottia* Viette 57  
*phlegyra*, *Stathmopoda* 109  
*phortegella*, *Sagephora* 64  
*photinella*, *Trachypepla* 107  
*phragmitella*, *Limnaecia* 86  
*phricias*, *Graphania* 205  
*Phrissogonus* 189  
*phryganitis*, *Brachmia*, sp. nr 81  
*Phryganostola* 76  
*Phthorimaea* 83  
**PHYCITINAE** 141  
*Phycomorpha* 129  
*Phylacodes* 74  
*Phyllonorycter* 72  
*Physetica* 209  
*Phyzanica* 87  
*picarella*, *Izatha* 96  
*picarella* var.  $\beta$ , *Oecophora* 94  
*pictoriana*, *Planotortrix* 126  
*pictula*, *Meterana* 207  
**PIERIDAE** 139  
*Pieris* 139  
*placida*, *Daphnis* 140  
*plaesiosema*, *Symmetrischema* 83  
*plagiata*, *Pyrgotis* 127  
*plagiarella*, *Tingena* 104  
*plagiaula*, *Elachista* 112  
*plagifurcata*, *Cidaria* 183  
*plagigera*, *Glyphipteryx* 78  
*planetella*, *Izatha* 91  
*planetopa*, *Glaucocharis* 145  
*Planotortrix* 125  
*platyptera*, *Semiocosma* 94  
*Platytilla* 133  
*Platysenta* 196  
*plebejana*, *Crocidosema* 115  
*plemochoa*, *Kiwaia* 82  
*plena*, *Graphania* 205  
*plexippus*, *Danaus* 136  
*plinthina*, *Pasiphila* 188  
*plinthoglypta*, *Pyrgotis* 127  
*Plodia* 142  
*plumbea*, *Helastia* 181  
*plumbiflua*, *Stathmopoda* 108  
*plumbosa*, *Izatha* 96  
*plurilineata*, *Chlorochroma* 189  
*plurimata*, *Coremia* 179  
*plusiata*, *Hadena* 196  
**PLUSIINAE** 213  
*Plutella* auct. 75  
*Plutella Schrank* 74  
*pluto*, *Erebia* 139  
*pluviata*, *Dasyuris* 178  
*polias*, *Capua* 119  
*politata*, *Orthenches* 74  
*Politeia* 164  
*politis*, *Tingena* 102  
*polychroa*, *Mamestra* 202  
*polycymaria*, *Astheniodes* 189  
*polygonalis maorialis*, *Uresiphita* 154

- polyleuca*, *Trachypepla* 106  
*Polygonia* 166  
**Polyorthini** 114  
*polypodii*, *Philocryptica* 125  
*pomona*, *Catopsilia pomona* 139  
*pomonella*, *Cydia* 115  
*pongalis*, *Eudonia* 157  
*pontifica*, *Harmologa* 124  
*Porina* 56  
*porophora*, *Leptocroca* 97  
*porphyraula*, *Zelleria* auct. 76  
*porphyreana*, *Pyrgotis* 119  
*porphyrias*, *Paradetis* 186  
*porphyritis*, *Orthenches* 74  
*postvittana*, *Epiphyas* 117  
*potamias*, *Protithona* 116  
*practica*, *Xanthorhoe* 179  
*praecipitalis*, *Gelophaula* 124  
*praefactata*, *Orthoclydon* 185  
*praerupta*, *Astrocidaria* 176  
*praesignis*, *Meterana* 208  
*prasinias*, *Asaphodes* 174  
*prasinodes*, *Orthenches* 74  
*prasophyta*, *Izatha* 96  
*Prays* 75  
*primaria*, *Gymnobathra* 93  
*princeps*, *Venusia* 185  
*Pringleophaga* 64  
*prionistis*, *Graphania* 205  
*prionota*, *Astrocidaria* 176  
*priscana*, *Teras* 121  
*privatana*, *Paedisca* 120  
*probenota*, *Aletia* 200  
*Probolaea* 182  
*prochloraa*, *Ascerodes* 118  
*Proditrix* 75  
**PRODOXIDAE** 59  
*productata*, *Pseudocoremia* 168  
*profunda*, *Proteodes* 88  
*progama*, *Stigmella* 54  
*progonopis*, *Stigmella* 54  
*promelanaria*, *Panagra* 193  
*pronephela*, *Tingena* 105  
*propalaea*, *Stigmella* 54  
*prophetica*, *Lathicrossa* 96  
*propria*, *Tmetolophota* 210  
*prospiciens*, *Atomotricha* 90  
*proteastis*, *Mamestra* 208  
*Proteodes* 88  
*Proternia* 153  
*Proterodesma* 64  
*Proteroeca* 153  
*Prothelymna* 128  
*Prothinodes* 64  
*Protithona* 116  
*protochlora*, *Trachypepla* 107  
*Protosynaema* 75  
*prototoxa*, *Selidosema* 163  
**Protyparcha** 159  
*provida*, *Xylina* 212  
*prymnaea*, *Asaphodes* 174  
*psamathodes*, *Larentia* 180  
*psamminella*, *Bedellia* 79  
*psammitis*, *Eudonia* 158  
*psammochroa*, *Plutella* auct. 75  
*Pseudocoremia* 166  
*Pseudoips* 198  
*Pseudosestra* 169  
*pseudospretella*, *Hofmannophila* 94  
*pseudostinaria*, *Orthoclydon* 186  
*psithyra*, *Batrachedra* 84  
**PSYCHEODINAE** 67  
**PSYCHIDAE** 67  
*psychra*, *Izatha* 96  
*Pterophorus* 132  
**PTEROPHORIDAE** 132  
**PTEROPHOROIDEA** 132  
*ptyoptera*, *Anisoplaca* 80  
*pulchella*, *Deiopeia* 195  
*pulchelloides vaga*, *Utetheisa* 194  
*pulcherrima*, *Grammodes* 197  
*pulchraria*, *Poecilasthena* 189  
*pulla*, *Locheutis* 97  
*pulvrea*, *Phthorimaea* 83  
*pulverulenta*, *Platyptilia* 134  
*pumila*, *Elachista* 112  
*Kiwaia* 82  
*punana*, *Grapholita* 127  
*punctellus*, *Orocrambus* 149  
*punctigera*, *Heliothis (Helicoverpa)* 211  
*punctilineana*, *Teras* 170  
*punctilineata*, *Helastia* 181  
*pungata*, *Selidosema* 168  
*pnicea*, *Pasiphila* 188  
*pura*, *Scoparia* auct. 162  
*purdii*, *Tmetolophota* 210  
*purpurea*, *Austramathes*  
*Gracilaria* 71  
*purpurifera*, *Hydriomena* auct. 184  
*pusillidactyla*, *Lantanophaga* 132  
**PYRALIDAE** 141  
**PYRALINAE** 143  
*Pyralis* 143  
**PYRALOIDEA** 141  
*pyramaria*, *Cidaria* 173  
*pyramidiás*, *Pyrgotis* 127  
**PYRAUSTINAE** 152  
*Pyrgotis* 126  
*Pyroderces* auct. 86  
*pyrosalis*, *Endotricha* 143  
*pyrsophanes*, *Glaucocharis* 145

**Q**

- quadralis*, *Margarita* 153  
*quadrijuga*, *Sabatinca* 53  
*quadripuncta*, *Symmoca* 85  
*quadristrigata*, *Microdes* 184  
*quaestoria*, *Eudonia* 158  
*quaestuosa*, *Protosynaema* 75  
*querula*, *Cryptaspasma* 115  
*quieta*, *Phthorimaea* 83  
*quinquestriata*, *Lopharcha* 114

**R**

- radians*, *Thylacosceles* 119  
*Radinigoes* 195  
*rajella*, *Phyllonorycter* 92  
*rakaiaensis*, *Eudonia* 158  
*ramosellus*, *Orocrambus* 149  
*ranata*, *Sauris* 190  
*rangona*, *Crambus* 149  
*rapae*, *Pieris rapae* 139  
*Raumatia* 116  
*rauparaha*, *Lycaena* 135  
*recta*, *Asaphodes* 174  
*rectella*, *Tinea* 63  
*rectilineata*, *Chloroclystis* 184  
*recurvalis*, *Hymenia* 152  
*recusana*, *Conchylis* 127  
*Reductoderces* 69  
*regilla*, *Notoreas* (Meyrick) 185  
*repercussa*, *Scoparia* 156  
*repletalis aeolodes*, *Platyptilia* 134  
*repletearia*, *Acidalia* 193  
*repletatis repletatis*, *Platyptilia*  
*restincta*, *Isonomeutis* 129  
*reticularis*, *Harmologa* 124  
*Rhalsa* 211  
*Rhathamictis* 69  
*Rhigognostis* 76  
*rhodobapta*, *Cryptolechia* 87  
*rhodopleura*, *Mamestra* 207  
*Rictonis* 196  
*rigescens*, *Izatha* 96  
*risata*, *Asthena* 179  
*rivalis*, *Pasiphila* 189  
*rivularis*, *Phibalapteryx* 175  
*rixata*, *Hydriomena* auct. 184  
*robiniosa*, *Tingena* 105  
*robusta*, *Eurythecta* 118  
*robustaria*, *Coremia* 179  
**ROESLERSTAMMIIDAE** 70  
*ropeana*, *Teras* 121  
*rrorida*, *Zelleria* auct. 76  
*rosearia*, *Epyaxa* 180  
*roseata*, *Crypsitricha* 60  
    *Trachypepla* 107  
*rosicoma*, *Sabatinca* 53

- rotuella*, *Scoparia* 160  
*rubella*, *Pasiphila* 189  
*rubescens*, *Graphania* 205  
*rubiginella*, *Endophthora* 61  
*rubophaga*, *Heterocrossa* 131  
*rubraria*, *Scopula* 193  
*rubro viridans*, *Hepialus* 56  
*rubropunctaria*, *Epicyme* 179  
*rubroviridans*, *Charagia* 56  
*rudisata*, *Pseudocoremia rudisata* 169  
*rufescens*, *Larentia* 182  
*rufipellis*, *Chloroclystis* 188  
*rufopunctella*, *Gymnobathra* 93  
*rufosparsa*, *Proteodes* 88  
*rufulincta*, *Chloroclystis* 188  
*rugata*, *Glyphipteryx* 78  
*rureana*, *Catamacta* 119  
*rusticella*, *Tinea* 63  
*rutella*, *Setomorpha* 64  
*rutilans*, *Gracilaria* 71

**S**

- Sabatinca* 52  
*sabulosa*, *Borkhausenia* 103  
*sabulosella*, *Eudonia* 158  
*saeva*, *Melancha* 207  
*Sagephora* 64  
*sagittifera*, *Elachista* 112  
*Salebria* 142  
*salmachus*, *Sesia* 112  
*salmoni*, *Helastia* 181  
*salueta*, *Orthenches* 74  
*salustius*, *Lycaena* 135  
*Samana* 192  
*Samia* 140  
*sanctimonea*, *Heterocrossa* 131  
*sandycias*, *Pasiphila* 189  
*sanguinea*, *Harmologa* 124  
*sanguinolenta*, *Leptocroca* 96  
*sapiens*, *Aletia* 209  
*sarcanthes*, *Heterocrossa* 131  
*sarcitrella*, *Endrosis* 91  
*sarcoxantha*, *Gymnobathra* 93  
*Sarisa* 169  
*saristes*, *Crambus* 146  
**SATURNIIDAE** 140  
**SATYRINAE** 138  
*saxana*, *Sciaphila* 116  
*scabra*, *Argua* 164  
*scaphodes*, *Protyparcha* 159  
*scariphota*, *Selidosema* 168  
*Sceliodes* 153  
*Scenoploca* 144  
*schaefferella*, *Schiffermuelleria* 98  
*schedias*, *Crambus* 146  
*schematica*, *Kiwaia* 83

- Schiffermuelleria* 98  
*schistaria, Poecilasthena* 189  
*Schoenotenini* 128  
*scholaea, Leptocroca* 97  
*Schrankia* 211  
*Scieropepla* 110  
*scintellula, Glyphipterix* 78  
*scintilla, Glyphipterix* 78  
*scissa, Helastia* 181  
*scissaria, Theoxena* 192  
*scitulus, Crambus* 147  
*scoliastis, Harmologa* 124  
*Scoparia* 159  
*Scoparia* auct. 160  
**SCOPARIINAE** 154  
*scopariooides, Orocrambus* 149  
*Scopula* 193  
*Scoriodyta* 69  
*scoriota, Mallobathra* 68  
*scotosialis, Rapsa* 211  
*scripta, Scoparia* auct. 162  
*scriptaria, Cleora* 163  
*scruposa, Ericodesma* 118  
*scutata, Graphania* 205  
*scutatus, Orocrambus* 149  
**SCYTHRIDAE** 110  
*Scythris* 110  
*seclusa, Tingena* 105  
*sedata, Gelechia* 82  
*seelandica, Meserana* 67  
*selenaea, Glaucocharis* 145  
*selene, Actias* 140  
*selenitis, Caloptilia* 72  
*selenophora, Dasypodia* 197  
*Selidosema* 163, 166  
*semialbata, Eupithecia* 176  
*semicocta, Epichorista* (auct. 2) 123  
*semifasciata, Orthenches* 74  
 Tatosoma 190  
*semiferana, Capua* 119  
*semifissata, Xanthorhoe* auct. 191  
*semilauta, Trachypepla* 107  
*semilineata, Cidaria* 177  
*seminuda, Stathmopoda* 108  
*Semiocosma* 94  
*semisignata, Helastia* 181  
*semivittata, Tmetolophota* 210  
*semnodes, Cryptolechia* 87  
*semochlora, Pasiphila* 189  
*senex, Aoraia* 57  
*separata, Mythimna* 208  
*septentrionalis, Orthenches* 74  
*sequens, Graphania* 205  
*sera, Rhigognostis* 76  
*serena, Tingena* 105  
*sericata, Graphania* 205  
*sericea, Bityla* 195
- Chersotis* 213  
*Sericia* 198  
*sericodes, Asaphodes* 174  
*servana, Ctenopseustis* 121  
*servilisana, Sciaphila* 116  
*servularia, Fidonia?* 171  
**SESIIDAE** 112  
**SESIOIDEA** 112  
*Sestra* 169  
*Setomorpha* 64  
**SICULINAE** 141  
*sideraspis, Scoparia* 160  
*sideritis, Strepsicrates* 116  
*siderodeta, Tingena* 115  
*siderota, Tingena* 115  
*signata, Wiseana* 58  
*Simaethis* 113  
*similata, Austrocidaria* 176  
*similis, Glyphipterix* 78  
 Mnesarchaea 55  
*Orthenches* 74  
*Tmetolophota* 210  
*similisata, Cidaria* 182  
*simla, Dictyoploca* 140  
*simplex, Notoreas* (Meyrick) 185  
*Orocrambus* 149  
*simulans, Dichromodes* 192  
 Phibalapteryx 175  
*sinuata, Scoparia* auct. 162  
*sinuosa, Declana* 164  
 Tingena 105  
*siraea, Gelophaula* 124  
*sirenica, Bascantis* 60  
*siria, Cephalissa* 176  
*siriana, Epichorista* (auct. 2) 123  
*siriellus, Orocrambus* 149  
*siris, Arctesthes* 171  
 Helastia 182  
*sistens, Aletia* 200  
*sisyrana, Harmologa* 125  
*sisyrota, Titanomis* 214  
*Sitotroga* 83  
*skelloni, Hadena* 203  
*Stathmopoda* 108  
*sminthistis, Hysia* 202  
*smithi, Proteodes* 88  
*sollenis, Aletia* 201  
*somnulentella, Bedellia* 79  
*sophistes, Crambus* 147  
*sophorae, Stigmella* 54  
*sophronellus, Orocrambus* 149  
*sordida, Atomotricha* 90  
*Sorensenata* 128  
*sorenseni, Tinearupa sorenseni* 98  
 sp. Dumbleton, Aoraia 57  
 sp. indet., Coleophora 85  
 sp. nr *phryganitis*, Brachmia 81

- sp. nr *suffusa*, *Chloroclystis* 177  
*sparsa*, *Gelechia* 81  
*spartifoliella*, *Leucoptera* 79  
*spartodeta*, *Trachypepla* 107  
*spatiosa*, *Planotortrix* 126  
*specifica*, *Alysia* 204  
*speciosa*, *Harmologa* 125  
*spectans*, *Speiredonia* 198  
*Speiredonia* 198  
*Sperchia* 128  
*sphaeriata*, *Dichromodes* 192  
*sphagnea*, *Mamestra* 201  
*sphenias*, *Tortrix* auct. 128  
*sphenocosma*, *Tinea* sensu Meyrick 66  
*sphenota*, *Zelleria* auct. 76  
**SPHINGIDAE** 140  
*sphragitis*, *Chloroclystis* auct. 177  
*Sphyrelata* 98  
*spissicornis*, *Coleophora* 85  
*spodiaea*, *Mermeritis* 97  
*Spodoptera* 196  
*Spoladea* 153  
*spoliatana*, *Sciaphila* 129  
**Sporophyla** 142  
*spurcatana*, *Teras* 120  
*spurcata*, *Xylina* 204  
*squalida*, *Codaria* 184  
*squamea*, *Gymnobathra* 93  
*stella*, *Coridomorpha* 88  
     *Glaucocharis* 145  
*stellata*, *Hierodoris* auct. 73  
     *Zapryrastra* 87  
**STENOMATINAE** 109  
*stenopterella*, *Crocydopora* 141  
*Stenoptilia* 134  
*stephanitis*, *Asaphodes* 174  
*stereota*, *Crypsitricha* 60  
*steropaea*, *Eudonia* 158  
*steropastis*, *Sagephora* 64  
     *Tmetolophota* 210  
*steropucha*, *Protosynaema* 75  
**STERRHINAE** 193  
*stigmatica*, *Scotosia* 163  
*Stigmella* 53  
*stilbella*, *Erechthias* 62  
*stinaria*, *Asaphodes* 174  
*stipata*, *Meterana* 208  
*straminea*, *Noteraula* 115  
*strangulata*, ?*Camptogramma* 170  
*strategica*, *Dasyuris* 178  
     *Metacrias* 194  
*Stratocleis* 166  
*Stratonice* 171  
*Strepsiceros* 116  
**Strepsicrates** 116  
*streptophora*, *Stratocleis* 163  
*stricta*, *Austrocidaria* 176  
     *strigosus*, *Aphomia* 151  
*strigulata*, *Mallobathra* 68  
*strophaea*, *Culladia* 144  
*stulta*, *Tmetolophota* 210  
*suavis*, *Pseudocoremia* 169  
*subalpina*, *Mallobathra* 69  
*subantarctica*, *Xanthorhoe* 191  
*subcarinata*, *Sagephora* 64  
*subditella*, *Eudonia* 158  
     *Gelechia* 91  
*subdola*, *Tortrix* 117  
*subductata*, *Larentia* 180  
*subfasciata*, *Morova* 141  
*subferralis*, *Gauna* 143  
*subflava*, *Xanthorhoe* 174  
*subita*, *Scoparia* auct. 162  
*subitata*, *Scotosia* 176  
*sublicellus*, *Crambus* 150  
*sublignalis*, *Salebria* 142  
*sublitella*, *Thamnosara* 98  
*submarginalis*, *Eudonia* 158  
*subobscurata*, *Gingidiobora* 180  
*subochraria*, *Anachloris* 170  
*subpavonella*, *Elachista* 62  
*subpurpureata*, *Poecilasthena* 189  
*subrectaria*, *Hydriomena* auct. 184  
*subsecivella*, *Biloba* 80  
*subtentaria*, *Acidalia* 185  
*suffusa*, *Pasiphila* 189  
*sulcana*, *Tmetolophota* 210  
*sulpitiata*, *Hemerophila* 166  
*suppressaria*, *Horisme* 182  
*sylvestris*, *Scoparia* auct. 162  
*ylvicola*, *Dumbletonius* 57  
*symbolaea*, *Asterivora* 114  
*Symmetrischema* 83  
**SYMMOCINAE** 85  
*symmorphia*, *Eutorna* 88  
*Synanthedon* 112  
*synclinalis*, *Aponotoreas* 171  
*Syntomactis* 86  
*syntona*, *Planotortrix syntona*

## T

- taipana*, *Planotortrix* 126  
*tamarindi*, *Opsiphanes* 136  
*Tanaoctena* 76  
*Tanaoctenota* 76  
*taongella*, *Gelechia* 106  
*Taoscelis* 89  
*tapetrella*, *Trichophaga* 66  
*tarda*, *Blastobasis* 85  
*tartarea*, *Meterana* 208  
*Tathorhynchus* 198  
*Tatosoma* 190  
*Tauroscopa* 150

- Tawhitia* 151  
*Tebenna* 114  
*technica*, *Petasactis* 64  
*Teia* 195  
*temenaula*, *Leucania* 200  
*temperata*, *Aletia* 201  
*tenebrosa*, *Epichorista* (auct. 2) 123  
*tenuis*, *Athetis* 195  
*tephrophanes*, *Tingena* 105  
*Tephrosaria* 65  
*teras*, *Psecadia* 96  
*terminella*, *Erechthias* 62  
*terranea*, *Archyala* 60  
*terrena*, *Pseudocoremia* 169  
    *Tingena* 105  
*tessellatella*, *Lindera* 64  
*testulata*, *Chloroclystis* auct. 177  
*tetrachroa*, *Graphania* 205  
*tetracycla*, *Scoparia* auct. 162  
*tetragona*, *Proditrix* 75  
*tetraphala*, *Thiotricha* 83  
*texta*, *Tinea* sensu Meyrick 66  
*thalamota*, *Heterocrossa* 131  
*thalerodes*, *Tingena* 105  
*thallophora*, *Elachista* 112  
*Thallostoma* 65  
*Thambotricha* 132  
*Thamnosara* 98  
*theatralis*, *Epichorista* (auct. 1) 122  
*Thectophila* 86  
*Theoxena* 192  
*thermochromata*, *Ischalis* 166  
*thetodes*, *Gymnobathra* 92  
*Thiotricha* 83  
*Thistricha* 83  
*tholodella*, *Gymnobathra* 93  
*thoracica*, *Bityla* 195  
*thorybodes*, *Thiotricha* 83  
*thranias*, *Borkhausenia* 102  
*thrinodes*, *Crambus* 150  
*Thyatira* 193  
**THYATIRIDAE** 193  
*Thylacosceles* 109  
*thymiastes*, *Orocrambus* 149  
*Thyone* 171  
*thyraula*, *Kiwaia* 83  
*thyridias*, *Eudonia* 158  
**THYRIDIDAE** 141  
**THYRIDOIDEA** 141  
*Thysanoplusia* 214  
*tigrina*, *Archyala* 60  
*tigris*, *Planotortrix* 126  
*tillyardi*, *Asterivora* 114  
*timaralis*, *Diathrausta* 151  
*timarata*, *Cidaria* 176  
*timora*, *Tatosoma* 190  
*Tinea* sensu Meyrick 65  
*Tineola* 66  
*Tingena* 99  
*tipulata*, *Tatosoma* 190  
*tipuliformis*, *Synanthedon* 112  
*Tirumala* 136  
*Titanomis* 214  
*Tmetolophota* 209  
*tonnoiri*, *Mallobathra* 69  
*topia*, *Tatosoma* 190  
*toreuma*, *Izatha* 95  
*toreuta*, *Declana* 165  
*toriata*, *Microdes* 184  
*tornospila*, *Chloroclystis* 187  
*tornota*, *Pyrgotis* 127  
*torodes*, *Eudonia* 158  
*torogramma*, *Ecclitica* 121  
*toroneura*, *Tmetolophota* 210  
*toroterra*, *Harmologa* 125  
**TORTRICIDAE** 114  
**TORTRICINAE** 117  
*Tortricini* 129  
**TORTRICOIDEA** 114  
*Tortrix* auct. 128  
*toumatou*, *Orophora* 67  
*Trachybathra* 124  
*Trachypepla* 105  
*transaurea*, *Dasyuris* 178  
*transcissalis*, *Crambus* 150  
*transfixa*, *Pyrgotis* 127  
*transitaria*, *Tatosoma* 190  
*transtrigana*, *Sciaphila* 120  
*transversella*, *Argyresthia* 78  
    *Epithecis* 80  
*trapezitis*, *Tauroscopa* 151  
*trapezophora*, *Scoparia* auct. 162  
*traversii*, *Catocala* 197  
*triangularis*, *Grypotheeca* 67  
    *Scoparia* 212  
*triatma*, *Scythris* 110  
*tributaria*, *Gelophaula* 124  
*tricentra*, *Stigmella* 54  
*trichodeta*, *Anarsia* 80  
*Trichophaga* 66  
*trichroa*, *Catamacta* 127  
*triclera*, *Eudonia* 158  
*trifolii*, *Coleophora* 85  
*Trigonistis* 211  
*trimaculata*, *Eurythecta* 116  
*trimolybditis*, *Stathmopoda* 109  
*triorthota*, *Curvisacculus* 121  
*Trioxycanus* 56  
*triphragma*, *Helastia* 182  
*tripunctaria*, *Deilemera* 194

*triscelis*, *Scoparia*  
*triselena*, *Glyptipterix* 78  
*tristictica*, *Batrachedra* 84  
*tristis*, *Asterivora* 114  
    *Euchersadaula* 91  
*trisulca*, *Gelophaula* 124  
*Trithamnora* 66  
*tritochlora*, *Gelophaula* 119  
*tritonellus*, *Orocrambus* 149  
*trivirgata*, *Eudonia* 158  
*Tsochasta* 82  
*tuhualis*, *Orocrambus* 149  
*tuhuata*, *Acidalia* 189  
*tuicana*, *Scoparia* auct. 162  
*tungella*, *Glyptipterix* 78  
*turbida*, *Xylina* 202  
*turbotti*, *Proterodesma* 64  
*turbulentana*, *Sciaphila* 121  
*turicensis*, *Tinea* 65  
*turneri*, *Scoparia* auct. 162  
*tylogramma*, *Endophthora* 61  
*typhicola*, *Scieropepla* 110  
*typhlopa*, *Monopis* 63  
*Tyria* 194

## U

*umbra*, *Melanchra* 204  
*umbraculata*, *Wiseana* 58  
*umbrosa*, *Austrocidaria* 176  
*undalis*, *Hellula* 152  
*undosata*, *Epiphryne* 179  
*undulata*, *Xanthorhoe* 175  
*undulifera*, *Phibalapteryx* 175  
*unica*, *Tmetolophota* 210  
*unicolor*, *Orophora* 67  
*unilinea*, *Xyridacma* 193  
*unimaculatus*, *Dumbletonius* 57  
*urbana*, *Asterivora* 114  
*Uresiphita* 154  
*urticae*, *Pasiphila* 189  
*usitata*, *Pachycnemia* 169  
*ustaria*, *Xyridacma* 192  
*ustimacula*, *Scoparia* 160  
*ustiramis*, *Eudonia* 158  
*ustistriga*, *Graphania* 205  
*Utetheisa* 194  
*utuella*, *Oecophora* 92

## V

*vacua*, *Leptocroca* 97  
*vagata*, *Locheutis* 97  
*valenternota*, *Scoparia* auct. 162  
*vana*, *Gelophaula* 124  
*Vanicela* 70  
*vapidus*, *Crambus* 150

*varia*, *Eurythecta* 116  
    *Proteodes* 88  
*variabilis*, *Ischalis* 166  
    *Leptocroca* 97  
*varians*, *Lyrcea* 193  
*variegata*, *Capua* 127  
*variolarus*, *Pielus* 58  
*vates*, *Thambotricha* 132  
*veda*, *Agrotis* 213  
*Velasquez* 151  
*venipunctata*, *Epyaxa* 180  
*ventosus*, *Orocrambus* 150  
*venustatis*, *Austrocidaria* 176  
*venustula*, *Azelina* 166  
*veronicae*, *Xyridacma* 193  
*verriculata*, *Epiphryne* 179  
*verrucosa*, *Chlenias* 165  
*versarella*, *Coleophora* 85  
*versuta*, *Atomotricha* 90  
*vestita*, *Tingena* 105  
*vetustana*, *Olindia* 129  
*vexata*, *Porina* 58  
    *Xylina* 204  
*vibratrix*, *Helostibes* auct. 88  
*vieta*, *Pasiphila* 189  
*vigens*, *Erana*  
    *vigens* auct., *Stenoptilia* 134  
*vilis*, *Cacoecia* 124  
*villida calybe*, *Junonia* 137  
*villosa*, *Aponotoreas* 171  
*vinaria*, *Leptocroca* 97  
*vinitincta*, *Orthenches* 74  
*virescens*, *Aenetus* 56  
    *Aletia* 201  
*virgata*, *Orthenches* 74  
*viridis*, *Antichloris* 195  
    *Dianthoecia* 205  
*vitellus*, *Orocrambus* 150  
*vitiosa*, *Meterana* 208  
*voluta*, *Paedisca* 120  
*vulcanica*, *Pasithea* 186  
*vulgaris*, *Orocrambus* 150  
*vulpecula*, *Scoparia* auct. 162

## W

*watti*, *Cosmiotes*  
*Wiseana* 57  
*Witlesia* 154

## X

*xanthaspis*, *Epiphryne* 179  
*xanthialis*, *Adena* 152  
*xanthodelta*, *Euthictis* 92  
*xanthodesma*, *Tingena* 105  
*xanthogramma*, *Melanchra* 203

*xanthogrammus*, *Orocrambus* 150  
*xanthomicta*, *Tingena* 105  
*Xanthorhoe* auct. 190  
*Xeroscopa*  
*xestobela*, *Glyphipterix* 78  
*Xyleutes* 112  
*xylinana*, *Grapholita* 127  
XYLORYCTINAE 110  
*xylostella*, *Plutella* 75  
*Xynonia* 192  
*xyrias*, *Leptocroca* 97  
*Xyridacma* 192  
*xysmatias*, *Eudonia* 158  
*xystrota*, *Bactra* 115

Y

*ypsilonaria*, *Coremia* 191  
YPONOMEUTIDAE 73  
YPONOMEUTOIDEA 73

Z

*Zaprystra* 87  
*zatricha*, *Chloroclystis* 187

*zatrophana*, *Epichorista* (auct. 2) 123  
*Zeempista* 81  
*zelaea*, *Eurythecta* 118  
*Zelleria* auct. 76  
*zelota*, *Glyphipterix* 78  
*zephyrana*, *Gymnobathra* 65  
*Zermizinga* 170  
*zestodes*, *Tortrix* auct. 128  
ZEUGLOPTERA 52  
*ziczac*, *Caustaloma?* 166  
*zinckenella*, *Chrysoesthia* 81  
*zimmermanni*, *Kessleria* 73  
*Zirosaris* 94  
*Zizina* 136  
*zomeuta*, *Simaethis* 113  
*zonodoxa*, *Sabatinca* 53  
*zophodactyla*, *Stenoptilia* 134  
*zophochalca*, *Athrips* auct. 80  
*zophoclaena*, *Eudonia* 158  
*zophodactyla*, *Stenoptilia* 134  
*zophoessa*, *Eulechria* 91  
*zopyra*, *Paranotoreas* 186  
*zorianella*, *Acrocercops* 71  
*zygiana*, *Pyrgotis* 127  
*Zylobara* 166




ENVOI

*"A needless alexandrine ends my song,  
Which like a wounded snake drags its slow length  
along."*

—Alexander Pope





NATURAL  
HISTORY  
MUSEUM  
PUBLICATIONS

## ADVERTISEMENT

From the British Museum (Natural History):

### ***Occasional Papers on Systematic Entomology***

The objective of this new occasional series is to make available in hard copy some of the basic data that is essential to the preparation of comprehensive accounts of the world insect fauna. The papers have been fully researched bibliographically, and consist of checklists of nominal taxa, and faunal lists with information on host plants and localities, based mainly on the collections and libraries of the British Museum (Natural History).

Titles currently available:

No. 1 *Catalogue of the Neotropical Tigermoths.* A. Watson & D.T. Goodger, 1986. A4 paperback, 74 pp., 4 colour plates. 0 565 00994 X. £14.50.

No. 2. *An Annotated Checklist of the Carabidae (including Cicindelinae, Physodinae, and Paussinae) Recorded from Borneo.* N.E. Stork, 1986. A4 paperback, 24 pp., 1 map. 0 565 00995 8. £7.00.

No. 3. *A Catalogue of the Crotch Collection of Coccinellidae (Coleoptera).* R.D. Gordon, 1987. A4 paperback, 48 pp. 0 565 01016 6. £9.50.




No. 4. *An Annotated Checklist of Thysanoptera from Australia.* L.A. Mound & K.J. Houston, 1987. A4 paperback, 28 pp. 0 565 01036 0. £4.00.

Numbers are published at irregular intervals. Orders and inquiries should be sent to:

Publications Sales  
British Museum (Natural History)  
Cromwell Road  
London SW7 5BD  
ENGLAND

---

# TAXON :




This proforma data sheet may be photocopied as required; copyright is waived. Use the back for data such as host records.

JAN	<input type="checkbox"/>
FEB	<input type="checkbox"/>
MAR	<input type="checkbox"/>
APR	<input type="checkbox"/>
MAY	<input type="checkbox"/>
JUN	<input type="checkbox"/>
JUL	<input type="checkbox"/>
AUG	<input type="checkbox"/>
SEP	<input type="checkbox"/>
OCT	<input type="checkbox"/>
NOV	<input type="checkbox"/>
DEC	<input type="checkbox"/>

This base-map is used in the *Fauna of N.Z.* for recording the distribution of collection localities. The small grid divisions are 10 minutes of arc by latitude and longitude.

## DISTRIBUTION IN TIME

Suggested subdivisions:  
(A, adult; E, egg; J, juvenile;  
L, larva; P, pupa)



# Fauna of New Zealand




Number 14

Lepidoptera  
– annotated catalogue, and  
keys to family-group taxa

J. S. Dugdale

XXV • ANNIVERSARY ISSUE • XXV  
SYSTEMATICS GROUP, ENTOMOLOGY DIVISION, DSIR  
1963–1988




### **North Island**

AK – Auckland  
 BP – Bay of Plenty  
 CL – Coromandel  
 GB – Gisborne  
 HB – Hawkes Bay  
 ND – Northland  
 RI – Rangitikei  
 TK – Taranaki  
 TO – Taupo  
 WA – Wairarapa  
 WI – Wanganui  
 WN – Wellington  
 WO – Waikato

### **South Island**

BR – Buller  
 CO – Central Otago  
 DN – Dunedin  
 FD – Fiordland  
 KA – Kaikoura  
 NN – Nelson  
 MB – Marlborough  
 NC – North Canterbury  
 WD – Westland



SI – Stewart Island

MB – Marlborough  
 MC – Mid Canterbury  
 MK – Mackenzie  
 NC – North Canterbury  
 NN – Nelson  
 OL – Otago Lakes  
 SC – South Canterbury  
 SD – Marlborough Sounds  
 SL – Southland  
 WD – Westland

# Fauna of New Zealand

INTRODUCTION

KEYS TO TAXA

ILLUSTRATIONS

CATALOGUE

REFERENCES

INDEX TO TAXA

This series of refereed occasional publications has been established with two major objectives: to encourage those with expert knowledge of elements in the New Zealand fauna to publish concise yet comprehensive accounts; and to provide a means of identification accessible to the non-specialist. It will deal with non-marine invertebrates, since the vertebrates are well documented, and marine forms are covered by the series *Marine Fauna of New Zealand*.

Contributors should discuss their intentions with an appropriate member of the Fauna Advisory Group or with the Series Editor before commencing work (for names and addresses, see page ii). All necessary guidance will be given.

Persons wishing to receive issues of the *Fauna* should address inquiries to the Publications Officer, Science Information Publishing Centre, DSIR, P.O. Box 9741, Wellington, New Zealand, who will maintain standing orders in three categories, as follows. 'A' — an invoice will be sent for each number, as soon after publication as possible. 'B' — essentially as for 'A', but invoices will be sent only for those numbers in a nominated field of interest (e.g., beetles only, mites only). 'C' — updated catalogues and order forms will be sent from time to time. Orders should be accompanied by full payment; rates quoted are surface mail, New Zealand (GST included) and overseas. All overseas orders are charged at the US\$ rate.

## IN PRINT

No. 1 *Terebrantia* (Insecta: Thysanoptera), by Laurence A. Mound & Annette K. Walker. ISBN 0-477-06687-9. Published 23 December 1982. 120 p. Price NZ\$29.95 (overseas US\$29.95).

No. 2 *Osoriinae* (Insecta: Coleoptera: Staphylinidae), by H. Pauline McColl. ISBN 0-477-06688-7. Published 23 December 1982. Second impression May 1983. 96 p. Price NZ\$18.60 (overseas US\$18.60).

No. 3 *Anthribidae* (Insecta: Coleoptera), by B. A. Holloway. ISBN 0-477-06703-4. Published 23 December 1982. Second impression February 1985. 272 p. Price NZ\$41.00 (overseas US\$41.00).

No. 4 *Eriophyoidea* except *Eriophyinae* (Arachnida: Acari), by D. C. M. Manson. ISBN 0-477-06745-X. Published 12 November 1984. 144 p. Price NZ\$29.95 (overseas US\$29.95).

No. 5 *Eriophyinae* (Arachnida: Acari: Eriophyoidea), by D. C. M. Manson. ISBN 0-477-06746-8. Published 14 November 1984. 128 p. Price NZ\$29.95 (overseas US\$29.95).

No. 6 *Hydraenidae* (Insecta: Coleoptera), by R. G. Ordish. ISBN 0-477-06747-6. Published 12 November 1984. 64 p. Price NZ\$18.60 (overseas US\$18.60).

No. 7 *Cryptostigmata* (Arachnida: Acari) — a concise review, by M. Luxton. ISBN 0-477-06762-X. Published 8 December 1985. 112 p. Price NZ\$29.95 (overseas US\$29.95).

No. 8 *Calliphoridae* (Insecta: Diptera), by James P. Dear. ISBN 0-477-06764-6. Published 24 February 1986. 88 p. Price NZ\$18.60 (overseas US\$18.60).

No. 9 *Protura* (Insecta), by S. L. Tuxen. ISBN 0-477-06765-4. Published 24 February 1986. 52 p. Price NZ\$18.60 (overseas US\$18.60).

No. 10 *Tubulifera* (Insecta: Thysanoptera), by Laurence A. Mound & Annette K. Walker. ISBN 0-477-06784-0. Published 22 September 1986. 144 p. Price NZ\$34.65 (overseas US\$34.65).

No. 11 *Pseudococcidae* (Insecta: Hemiptera), by J. M. Cox. ISBN 0-477-06791-3. Published 7 April 1987. 232 p. Price NZ\$49.95 (overseas US\$49.95).

No. 12 *Pompilidae* (Insecta: Hymenoptera), by A. C. Harris. ISBN 0-477-02501-3. Published 13 November 1987. 160 p. Price NZ\$39.95 (overseas US\$39.95).

No. 13 *Encyrtidae* (Insecta: Hymenoptera), by J. S. Noyes. ISBN 0-477-02517-X. Published 9 May 1988. 192 p. Price NZ\$44.95 (overseas US\$44.95).

No. 14 *Lepidoptera* — annotated catalogue, and keys to family-group taxa, by J. S. Dugdale. ISBN 0-477-02518-8. 264 p. Publication date and price to be announced.

## IN PREPARATION (and scheduled for early publication)

Insecta Ambositrinae, by I. Naumann. Nepticulidae, by H. Donner & C. Wilkinson. Chalcidoidea (part), by J. S. Noyes & E. W. Valentine.

This is a PDF facsimile of the printed publication, and is fully searchable. It is supplied for individual use only and is not to be posted on websites (links should be made to the page from which it was downloaded).

No part of this work covered by copyright may be reproduced or copied in any form or by any means (graphic, electronic, or mechanical, including photocopying, recording, taping, information retrieval systems, or otherwise) without the written permission of the publisher.

*Fauna of New Zealand* website copy 2008,  
[www.landcareresearch.co.nz](http://www.landcareresearch.co.nz)

Dugdale, J. S. 1988: Lepidoptera – annotated catalogue, and keys to family-group taxa. *Fauna of New Zealand* 14, 264 pp.

Date of publication: 23 September 1988  
*Fauna of New Zealand*, ISSN 0111-5383; 14  
ISBN 0-477-02518-8

New Zealand Lepidoptera: catalogue. Scanned images from BUGZ project ([www.bugz.org.nz](http://www.bugz.org.nz)) provided by Stephen Pawson for OCR. Text OCRed and corrected for this searchable PDF by Trevor Crosby, FNZ series editor, 29 July 2008. Users may extract text from this PDF for their own use, but must check it against the original document for text sequence, accuracy, and formatting.